



88117302



**MATHEMATICS**  
**STANDARD LEVEL**  
**PAPER 2**

Candidate session number

0	0								
---	---	--	--	--	--	--	--	--	--

Thursday 3 November 2011 (morning)

Examination code

1 hour 30 minutes

8	8	1	1	-	7	3	0	2
---	---	---	---	---	---	---	---	---

## INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions on the answer sheets provided. Write your session number on each answer sheet, and attach them to this examination paper and your cover sheet using the tag provided.
- At the end of the examination, indicate the number of sheets used in the appropriate box on your cover sheet.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.



0112

Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, e.g. if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

**SECTION A**

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 7]

Let  $f(x) = 2x + 4$  and  $g(x) = 7x^2$ .


- (a) Find  $f^{-1}(x)$ . [3 marks]
- (b) Find  $(f \circ g)(x)$ . [2 marks]
- (c) Find  $(f \circ g)(3.5)$ . [2 marks]

A large rectangular box containing horizontal dotted lines for writing answers. A large, faint watermark logo for 'SAT PREP AHMEDABAD' with 'SP' in the center is overlaid on the box.



2. [Maximum mark: 6]

The cumulative frequency curve below represents the heights of 200 sixteen-year-old boys.



Use the graph to answer the following.

- (a) Write down the median value. [1 mark]
- (b) A boy is chosen at random. Find the probability that he is shorter than 161 cm. [2 marks]
- (c) Given that 82 % of the boys are taller than  $h$  cm, find  $h$ . [3 marks]

.....

.....

.....

.....

.....

.....


.....

.....



3. [Maximum mark: 6]

Consider the following circle with centre O and radius 6.8 cm.



*diagram  
not to scale*

The length of the arc PQR is 8.5 cm.

- (a) Find the value of  $\theta$ . [2 marks]
- (b) Find the area of the shaded region. [4 marks]

Handwriting practice area with horizontal dotted lines. A large watermark for 'SAT PREP SP AHMEDABAD' is visible in the background.





4. [Maximum mark: 6]

Consider the triangle ABC, where  $AB = 10$ ,  $BC = 7$  and  $\hat{C}AB = 30^\circ$ .

(a) Find the two possible values of  $\hat{A}CB$ . [4 marks]

(b) Hence, find  $\hat{A}BC$ , given that it is acute. [2 marks]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



5. [Maximum mark: 5]

Consider the expansion of  $(3x^2 + 2)^9$ .

(a) Write down the number of terms in the expansion.

[1 mark]

(b) Find the term in  $x^4$ .

[4 marks]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



6. [Maximum mark: 8]

Jose takes medication. After  $t$  minutes, the concentration of medication left in his bloodstream is given by  $A(t) = 10(0.5)^{0.014t}$ , where  $A$  is in milligrams per litre.

- (a) Write down  $A(0)$ . [1 mark]
- (b) Find the concentration of medication left in his bloodstream after 50 minutes. [2 marks]
- (c) At 13:00, when there is no medication in Jose's bloodstream, he takes his first dose of medication. He can take his medication again when the concentration of medication reaches 0.395 milligrams per litre. What time will Jose be able to take his medication again? [5 marks]

The answer area contains a large watermark logo for 'SAT PREP AHMEDABAD' in the center. The logo consists of a circular border with the text 'SAT PREP' at the top and 'AHMEDABAD' at the bottom, surrounding the letters 'SP' in a stylized font. Below the logo, there are approximately 15 horizontal dotted lines for writing the answers to the questions.



7. [Maximum mark: 7]

Let  $f(t) = 2t^2 + 7$ , where  $t > 0$ . The function  $v$  is obtained when the graph of  $f$  is transformed by

a stretch by a scale factor of  $\frac{1}{3}$  parallel to the  $y$ -axis,  
followed by a translation by the vector  $\begin{pmatrix} 2 \\ -4 \end{pmatrix}$ .

(a) Find  $v(t)$ , giving your answer in the form  $a(t-b)^2 + c$ . [4 marks]

(b) A particle moves along a straight line so that its velocity in  $\text{ms}^{-1}$ , at time  $t$  seconds, is given by  $v$ . Find the distance the particle travels between  $t = 5.0$  and  $t = 6.8$ . [3 marks]

The answer area contains a large watermark logo for 'SAT PREP SP AHMEDABAD' centered over a grid of horizontal dotted lines. The logo consists of a circular border with the text 'SAT PREP' at the top and 'AHMEDABAD' at the bottom, and the letters 'SP' in the center with a green leaf-like graphic below them.



Do **NOT** write solutions on this page. Any working on this page will **NOT** be marked.

### SECTION B

Answer **all** the questions on the answer sheets provided. Please start each question on a new page.

8. [Maximum mark: 14]

(a) Consider an infinite geometric sequence with  $u_1 = 40$  and  $r = \frac{1}{2}$ .

(i) Find  $u_4$ .

(ii) Find the sum of the infinite sequence.

[4 marks]

Consider an arithmetic sequence with  $n$  terms, with first term  $(-36)$  and eighth term  $(-8)$ .

(b) (i) Find the common difference.

(ii) Show that  $S_n = 2n^2 - 38n$ .

[5 marks]

(c) The sum of the infinite geometric sequence is equal to twice the sum of the arithmetic sequence. Find  $n$ .

[5 marks]



Do **NOT** write solutions on this page. Any working on this page will **NOT** be marked.

9. [Maximum mark: 16]


A company produces a large number of water containers. Each container has two parts, a bottle and a cap. The bottles and caps are tested to check that they are not defective.

A cap has a probability of 0.012 of being defective. A random sample of 10 caps is selected for inspection.

- (a) Find the probability that exactly one cap in the sample will be defective. [2 marks]
- (b) The sample of caps passes inspection if at most one cap is defective. Find the probability that the sample passes inspection. [2 marks]

The heights of the bottles are normally distributed with a mean of 22 cm and a standard deviation of 0.3 cm.

- (c) (i) **Copy** and complete the following diagram, shading the region representing where the heights are less than 22.63 cm.



- (ii) Find the probability that the height of a bottle is less than 22.63 cm. [5 marks]
- (d) (i) A bottle is accepted if its height lies between 21.37 cm and 22.63 cm. Find the probability that a bottle selected at random is accepted.
- (ii) A sample of 10 bottles passes inspection if all of the bottles in the sample are accepted. Find the probability that the sample passes inspection. [5 marks]
- (e) The bottles and caps are manufactured separately. A sample of 10 bottles and a sample of 10 caps are randomly selected for testing. Find the probability that both samples pass inspection. [2 marks]



Do **NOT** write solutions on this page. Any working on this page will **NOT** be marked.

10. [Maximum mark: 15]

Let  $f(x) = \frac{20x}{e^{0.3x}}$ , for  $0 \leq x \leq 20$ .

- (a) Sketch the graph of  $f$ . [3 marks]
- (b) (i) Write down the  $x$ -coordinate of the maximum point on the graph of  $f$ .  
(ii) Write down the interval where  $f$  is increasing. [3 marks]
- (c) Show that  $f'(x) = \frac{20 - 6x}{e^{0.3x}}$ . [5 marks]
- (d) Find the interval where the rate of change of  $f$  is increasing. [4 marks]





Please **do not** write on this page.  
Answers written on this page will  
not be marked.







22127304



**MATHEMATICS**  
**STANDARD LEVEL**  
**PAPER 2**

Candidate session number

0	0								
---	---	--	--	--	--	--	--	--	--

Friday 4 May 2012 (morning)

Examination code

1 hour 30 minutes

2	2	1	2	-	7	3	0	4
---	---	---	---	---	---	---	---	---

## INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions on the answer sheets provided. Write your session number on each answer sheet, and attach them to this examination paper and your cover sheet using the tag provided.
- At the end of the examination, indicate the number of sheets used in the appropriate box on your cover sheet.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL information booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].



0116



Please **do not** write on this page.

Answers written on this page  
will not be marked.



Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, e.g. if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

### SECTION A

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 6]

The first three terms of an arithmetic sequence are 36, 40, 44, ...

- (a) (i) Write down the value of  $d$ .
- (ii) Find  $u_8$ . [3 marks]
- (b) (i) Show that  $S_n = 2n^2 + 34n$ .
- (ii) Hence, write down the value of  $S_{14}$ . [3 marks]



2. *[Maximum mark: 7]*

Let  $f(x) = 2x^2 - 8x - 9$ .

(a) (i) Write down the coordinates of the vertex.

(ii) Hence or otherwise, express the function in the form  $f(x) = 2(x - h)^2 + k$ . *[4 marks]*


(b) Solve the equation  $f(x) = 0$ . *[3 marks]*



3. [Maximum mark: 6]

$$\text{Let } \mathbf{M} = \begin{pmatrix} x & 2x \\ x^2 & 1 \end{pmatrix} \text{ and } \mathbf{N} = \begin{pmatrix} 2 & 3 & -1 \\ 4 & 2 & 0 \\ 1 & 5 & 1 \end{pmatrix}.$$

- (a) Find  $\det \mathbf{M}$ . [2 marks]
- (b) Write down  $\det \mathbf{N}$ . [1 mark]
- (c) Find the value of  $x$  for which  $\det \mathbf{M} = \det \mathbf{N}$ . [3 marks]




The answer area contains a large watermark logo for SAT PREP AHMEDABAD. The logo is circular with 'SAT PREP' at the top and 'AHMEDABAD' at the bottom. In the center, there are the letters 'SP' in a large, stylized font, with a green leaf-like shape behind the 'P'.



4. [Maximum mark: 7]

The graph of  $y = (x-1)\sin x$ , for  $0 \leq x \leq \frac{5\pi}{2}$ , is shown below.



The graph has  $x$ -intercepts at 0, 1,  $\pi$  and  $k$ .

(a) Find  $k$ . [2 marks]

The shaded region is rotated  $360^\circ$  about the  $x$ -axis. Let  $V$  be the volume of the solid formed.

(b) Write down an expression for  $V$ . [3 marks]

(c) Find  $V$ . [2 marks]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....




5. [Maximum mark: 6]

Let  $M = \begin{pmatrix} p & -1 & -2 \\ 1 & 1 & -2 \\ 1 & q & -1 \end{pmatrix}$  and  $M^{-1} = \frac{1}{2} \begin{pmatrix} 3 & -5 & 4 \\ -1 & 1 & 0 \\ 1 & -3 & 2 \end{pmatrix}$ .

- (a) Find the value of  $p$  and of  $q$ . [3 marks]
- (b) Solve the system of linear equations.

$$\begin{aligned} px - y - 2z &= 7 \\ x + y - 2z &= 2 \\ x + qy - z &= -3 \end{aligned}$$

[3 marks]



6. [Maximum mark: 6]

Consider the expansion of  $\left(2x^3 + \frac{b}{x}\right)^8 = 256x^{24} + 3072x^{20} + \dots + kx^0 + \dots$

(a) Find  $b$ .

[3 marks]

(b) Find  $k$ .

[3 marks]

A large rectangular area with horizontal dotted lines for writing answers. A watermark logo is visible in the center, featuring the text "SAT PREP" at the top, "AHMEDABAD" at the bottom, and "SP" in large letters in the middle, with a green leaf-like graphic element.





7. *[Maximum mark: 7]*

The probability of obtaining “tails” when a biased coin is tossed is 0.57. The coin is tossed ten times. Find the probability of obtaining

- (a) **at least** four tails; *[4 marks]*
- (b) the fourth tail on the tenth toss. *[3 marks]*

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Turn over


Do **NOT** write solutions on this page.

**SECTION B**

Answer **all** questions on the answer sheets provided. Please start each question on a new page.

8. [Maximum mark: 13]

The histogram below shows the time  $T$  seconds taken by 93 children to solve a puzzle.



The following is the frequency distribution for  $T$ .

Time	$45 \leq T < 55$	$55 \leq T < 65$	$65 \leq T < 75$	$75 \leq T < 85$	$85 \leq T < 95$	$95 \leq T < 105$	$105 \leq T < 115$
Frequency	7	14	$p$	20	18	$q$	6

(a) (i) Write down the value of  $p$  and of  $q$ .

(ii) Write down the median class.

[3 marks]

(b) A child is selected at random. Find the probability that the child takes less than 95 seconds to solve the puzzle.

[2 marks]

(This question continues on the following page)



Do **NOT** write solutions on this page.

(Question 8 continued)

Consider the class interval  $45 \leq T < 55$ .

(c) (i) Write down the interval width.

(ii) Write down the mid-interval value.

[2 marks]

(d) Hence find an estimate for the

(i) mean;


(ii) standard deviation.

[4 marks]

John assumes that  $T$  is normally distributed and uses this to estimate the probability that a child takes less than 95 seconds to solve the puzzle.

(e) Find John's estimate.


[2 marks]



Do **NOT** write solutions on this page.

9. [Maximum mark: 15]

The following diagram shows a triangle ABC.




$BC = 6$ ,  $\hat{CAB} = 0.7$  radians,  $AB = 4p$ ,  $AC = 5p$ , where  $p > 0$ .

(a) (i) Show that  $p^2(41 - 40\cos 0.7) = 36$ .

(ii) Find  $p$ .

[4 marks]

Consider the circle with centre B that passes through the point C. The circle cuts the line CA at D, and  $\hat{ADB}$  is obtuse. Part of the circle is shown in the following diagram.



(b) Write down the length of BD.

[1 mark]

(c) Find  $\hat{ADB}$ .

[4 marks]

(d) (i) Show that  $\hat{CBD} = 1.29$  radians, correct to 2 decimal places.

(ii) Hence, find the area of the shaded region.


[6 marks]



Do **NOT** write solutions on this page.

10. [Maximum mark: 17]

The following diagram shows two ships A and B. At noon, ship A was 15 km due north of ship B. Ship A was moving south at  $15 \text{ km h}^{-1}$  and ship B was moving east at  $11 \text{ km h}^{-1}$ .



(a) Find the distance between the ships

(i) at 13:00;

(ii) at 14:00.

[5 marks]

Let  $s(t)$  be the distance between the ships  $t$  hours after noon, for  $0 \leq t \leq 4$ .

(b) Show that  $s(t) = \sqrt{346t^2 - 450t + 225}$ .

[6 marks]

(c) Sketch the graph of  $s(t)$ .

[3 marks]

(d) Due to poor weather, the captain of ship A can only see another ship if they are less than 8 km apart. Explain why the captain cannot see ship B between noon and 16:00.

[3 marks]





Please **do not** write on this page.  
Answers written on this page  
will not be marked.





Please **do not** write on this page.  
Answers written on this page  
will not be marked.





Please **do not** write on this page.  
Answers written on this page  
will not be marked.







22127306



**MATHEMATICS**  
**STANDARD LEVEL**  
**PAPER 2**

Candidate session number

0	0								
---	---	--	--	--	--	--	--	--	--

Friday 4 May 2012 (morning)

Examination code

2	2	1	2	-	7	3	0	6
---	---	---	---	---	---	---	---	---

1 hour 30 minutes

## INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions on the answer sheets provided. Write your session number on each answer sheet, and attach them to this examination paper and your cover sheet using the tag provided.
- At the end of the examination, indicate the number of sheets used in the appropriate box on your cover sheet.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL information booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].



0112


Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, e.g. if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

**SECTION A**

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 6]

The following diagram shows  $\Delta PQR$ , where  $RQ = 9$  cm,  $\hat{P}RQ = 70^\circ$  and  $\hat{P}QR = 45^\circ$ .



- (a) Find  $\hat{R}PQ$ . [1 mark]
- (b) Find PR. [3 marks]
- (c) Find the area of  $\Delta PQR$ . [2 marks]

Working area with horizontal dotted lines for writing answers.




2. [Maximum mark: 6]

Let  $f(x) = \cos(e^x)$ , for  $-2 \leq x \leq 2$ .

(a) Find  $f'(x)$ . [2 marks]

.....  
.....  
.....  
.....

(b) On the grid below, sketch the graph of  $f'(x)$ . [4 marks]



3. [Maximum mark: 6]

The first term of a geometric sequence is 200 and the sum of the first four terms is 324.8.

(a) Find the common ratio. [4 marks]

(b) Find the tenth term. [2 marks]

The answer area contains horizontal dotted lines for writing. A watermark logo is centered over the lines, featuring the text "SAT PREP" at the top, "AHMEDABAD" at the bottom, and a large "SP" in the middle with a green leaf-like shape.



4. [Maximum mark: 6]

The heights of a group of seven-year-old children are normally distributed with mean 117 cm and standard deviation 5 cm. A child is chosen at random from the group.

- (a) Find the probability that this child is taller than 122.5 cm. [3 marks]
- (b) The probability that this child is shorter than  $k$  cm is 0.65. Find the value of  $k$ . [3 marks]



5. [Maximum mark: 6]

A particle moves in a straight line with velocity  $v = 12t - 2t^3 - 1$ , for  $t \geq 0$ , where  $v$  is in centimetres per second and  $t$  is in seconds.

(a) Find the acceleration of the particle after 2.7 seconds. [3 marks]

(b) Find the displacement of the particle after 1.3 seconds. [3 marks]



6. [Maximum mark: 7]

Let  $A = \begin{pmatrix} -1 & -1 & 2 \\ -1 & 0 & 1 \\ 2 & 1 & -2 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 0 & 2 \\ 0 & -1 & 0 \\ 3 & 1 & 2 \end{pmatrix}$ .

(a) Write down  $A^{-1}$ .

[2 marks]

(b) Let  $C$  be a  $3 \times 3$  matrix such that  $ACA^{-1} = B$ . Find  $C$ .

[5 marks]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



7. [Maximum mark: 8]

A factory makes lamps. The probability that a lamp is defective is 0.05. A random sample of 30 lamps is tested.

- (a) Find the probability that there is at least one defective lamp in the sample. [4 marks]
- (b) Given that there is at least one defective lamp in the sample, find the probability that there are at most two defective lamps. [4 marks]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....






Do **NOT** write solutions on this page.

**SECTION B**

Answer **all** questions on the answer sheets provided. Please start each question on a new page.

8. [Maximum mark: 16]

The following diagram shows the cuboid (rectangular solid) OABCDEFG, where O is the origin, and  $\vec{OA} = 4\mathbf{i}$ ,  $\vec{OC} = 3\mathbf{j}$ ,  $\vec{OD} = 2\mathbf{k}$ .



- (a) (i) Find  $\vec{OB}$ .
- (ii) Find  $\vec{OF}$ .
- (iii) Show that  $\vec{AG} = -4\mathbf{i} + 3\mathbf{j} + 2\mathbf{k}$ . [5 marks]
- (b) Write down a vector equation for
  - (i) the line OF;
  - (ii) the line AG. [4 marks]
- (c) Find the obtuse angle between the lines OF and AG. [7 marks]



Do **NOT** write solutions on this page.

9. [Maximum mark: 13]

Let  $f(x) = ax^3 + bx^2 + c$ , where  $a$ ,  $b$  and  $c$  are real numbers. The graph of  $f$  passes through the point  $(2, 9)$ .

(a) Show that  $8a + 4b + c = 9$ . [2 marks]

The graph of  $f$  has a local minimum at  $(1, 4)$ .

(b) Find two other equations in  $a$ ,  $b$  and  $c$ , giving your answers in a similar form to part (a). [7 marks]


(c) Find the value of  $a$ , of  $b$  and of  $c$ . [4 marks]



Do **NOT** write solutions on this page.

10. [Maximum mark: 16]

Consider the following circle with centre  $O$  and radius  $r$ .



The points  $P$ ,  $R$  and  $Q$  are on the circumference,  $\widehat{POQ} = 2\theta$ , for  $0 < \theta < \frac{\pi}{2}$ .

(a) Use the cosine rule to show that  $PQ = 2r \sin \theta$ . [4 marks]

Let  $l$  be the length of the arc  $PRQ$ .

(b) Given that  $1.3PQ - l = 0$ , find the value of  $\theta$ . [5 marks]

Consider the function  $f(\theta) = 2.6 \sin \theta - 2\theta$ , for  $0 < \theta < \frac{\pi}{2}$ .

(c) (i) Sketch the graph of  $f$ . [4 marks]

(ii) Write down the root of  $f(\theta) = 0$ . [4 marks]

(d) Use the graph of  $f$  to find the values of  $\theta$  for which  $l < 1.3PQ$ . [3 marks]





Please **do not** write on this page.  
Answers written on this page  
will not be marked.





88127302



**MATHEMATICS**  
**STANDARD LEVEL**  
**PAPER 2**

Candidate session number

0	0								
---	---	--	--	--	--	--	--	--	--

Wednesday 7 November 2012 (morning)

Examination code

1 hour 30 minutes

8	8	1	2	-	7	3	0	2
---	---	---	---	---	---	---	---	---

INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions on the answer sheets provided. Write your session number on each answer sheet, and attach them to this examination paper and your cover sheet using the tag provided.
- At the end of the examination, indicate the number of sheets used in the appropriate box on your cover sheet.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL information booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].



0112

Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, e.g. if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

**SECTION A**

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 6]

The first three terms of an arithmetic sequence are 5, 6.7, 8.4.

- (a) Find the common difference. [2 marks]
- (b) Find the 28<sup>th</sup> term of the sequence. [2 marks]
- (c) Find the sum of the first 28 terms. [2 marks]



2. [Maximum mark: 5]

$$\text{Let } A = \begin{pmatrix} 2 & 2 & 1 \\ 3 & 1 & 0 \\ 4 & 2 & 1 \end{pmatrix}.$$

(a) Write down  $A^{-1}$ . [2 marks]

(b) Hence or otherwise, find  $B$ , given that  $AB = \begin{pmatrix} -1 & 6 & -1 \\ 5 & -1 & 3 \\ 5 & 2 & 7 \end{pmatrix}$ . [3 marks]

.....

.....

.....

.....

.....

.....


.....

.....

.....


.....

.....



3. [Maximum mark: 6]

Let  $f(x) = x^3 - 2x - 4$ . The following diagram shows part of the curve of  $f$ .



The curve crosses the  $x$ -axis at the point P.

- (a) Write down the  $x$ -coordinate of P. [1 mark]
- (b) Write down the gradient of the curve at P. [2 marks]
- (c) Find the equation of the normal to the curve at P, giving your equation in the form  $y = ax + b$ . [3 marks]

.....

.....

.....

.....

.....

.....

.....

.....





4. [Maximum mark: 7]


The third term in the expansion of  $(2x + p)^6$  is  $60x^4$ . Find the possible values of  $p$ .

A large rectangular box with a black border, containing horizontal dotted lines for writing. In the center of the box is a large, faint watermark logo for 'SAT PREP AHMEDABAD'. The logo features the letters 'SP' in a stylized font with a green leaf-like shape behind them, surrounded by a circular border with the text 'SAT PREP' at the top and 'AHMEDABAD' at the bottom.



5. [Maximum mark: 6]

Let  $f(x) = a \cos(b(x-c))$ . The diagram below shows part of the graph of  $f$ , for  $0 \leq x \leq 10$ .



The graph has a local maximum at  $P(3, 5)$ , a local minimum at  $Q(7, -5)$ , and crosses the  $x$ -axis at  $R$ .

- (a) Write down the value of
  - (i)  $a$ ;
  - (ii)  $c$ . [2 marks]
- (b) Find the value of  $b$ . [2 marks]
- (c) Find the  $x$ -coordinate of  $R$ . [2 marks]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



6. *[Maximum mark: 8]*

In a large city, the time taken to travel to work is normally distributed with mean  $\mu$  and standard deviation  $\sigma$ . It is found that 4 % of the population take less than 5 minutes to get to work, and 70 % take less than 25 minutes.

Find the value of  $\mu$  and of  $\sigma$ .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....




7. [Maximum mark: 7]

A particle's displacement, in metres, is given by  $s(t) = 2t \cos t$ , for  $0 \leq t \leq 6$ , where  $t$  is the time in seconds.

(a) On the grid below, sketch the graph of  $s$ .

[4 marks]



(This question continues on the following page)



*(Question 7 continued)*

(b) Find the maximum velocity of the particle.

[3 marks]

A large rectangular area for writing, containing several horizontal dotted lines. In the center of this area is a large, faint watermark logo for "SAT PREP AHMEDABAD" featuring the letters "SP" in a stylized font with a green leaf-like graphic.



0912

Turn over


Do **NOT** write solutions on this page.

**SECTION B**

Answer **all** questions on the answer sheets provided. Please start each question on a new page.

8. [Maximum mark: 15]

The following diagram shows a circular play area for children.



The circle has centre O and a radius of 20 m, and the points A, B, C and D lie on the circle. Angle AOB is 1.5 radians.

- (a) Find the length of the chord [AB]. [3 marks]
- (b) Find the area of triangle AOB. [2 marks]

Angle BOC is 2.4 radians.

- (c) Find the length of arc ADC. [3 marks]
- (d) Find the area of the shaded region. [3 marks]
- (e) The shaded region is to be painted red. Red paint is sold in cans which cost \$32 each. One can covers  $140 \text{ m}^2$ . How much does it cost to buy the paint? [4 marks]



Do **NOT** write solutions on this page.

9. [Maximum mark: 15]

Consider the function  $f(x) = x^2 - 4x + 1$ .

(a) Sketch the graph of  $f$ , for  $-1 \leq x \leq 5$ . [4 marks]

This function can also be written as  $f(x) = (x - p)^2 - 3$ .

(b) Write down the value of  $p$ . [1 mark]

The graph of  $g$  is obtained by reflecting the graph of  $f$  in the  $x$ -axis, followed by a translation of  $\begin{pmatrix} 0 \\ 6 \end{pmatrix}$ .

(c) Show that  $g(x) = -x^2 + 4x + 5$ . [4 marks]

The graphs of  $f$  and  $g$  intersect at two points.

(d) Write down the  $x$ -coordinates of these two points. [3 marks]

Let  $R$  be the region enclosed by the graphs of  $f$  and  $g$ .

(e) Find the area of  $R$ . [3 marks]



Do **NOT** write solutions on this page.

10. [Maximum mark: 15]

At a large school, students are required to learn at least one language, Spanish or French. It is known that 75 % of the students learn Spanish, and 40 % learn French.

- (a) Find the percentage of students who learn **both** Spanish and French. [2 marks]
- (b) Find the percentage of students who learn Spanish, but not French. [2 marks]

At this school, 52 % of the students are girls, and 85 % of the girls learn Spanish.

- (c) A student is chosen at random. Let  $G$  be the event that the student is a girl, and let  $S$  be the event that the student learns Spanish.
- (i) Find  $P(G \cap S)$ .
- (ii) Show that  $G$  and  $S$  are **not** independent. [5 marks]
- (d) A boy is chosen at random. Find the probability that he learns Spanish. [6 marks]







22137304



**MATHEMATICS**  
**STANDARD LEVEL**  
**PAPER 2**

Candidate session number

0	0								
---	---	--	--	--	--	--	--	--	--

Friday 10 May 2013 (morning)

Examination code

2	2	1	3	-	7	3	0	4
---	---	---	---	---	---	---	---	---

1 hour 30 minutes

**INSTRUCTIONS TO CANDIDATES**

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL information booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].



0112

Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

**SECTION A**

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 7]

An arithmetic sequence is given by 5, 8, 11, ....

- (a) Write down the value of  $d$ . [1 mark]
- (b) Find
  - (i)  $u_{100}$ ;
  - (ii)  $S_{100}$ . [4 marks]
- (c) Given that  $u_n = 1502$ , find the value of  $n$ . [2 marks]

Working area with horizontal dotted lines for writing answers.



2. [Maximum mark: 6]

Consider the following cumulative frequency table.

$x$	Frequency	Cumulative frequency
5	2	2
15	10	12
25	14	26
35	$p$	35
45	6	41

- (a) Find the value of  $p$ . [2 marks]
  
- (b) Find
  - (i) the mean;
  - (ii) the variance. [4 marks]



3. [Maximum mark: 5]

In the expansion of  $(3x - 2)^{12}$ , the term in  $x^5$  can be expressed as  $\binom{12}{r} \times (3x)^p \times (-2)^q$ .

(a) Write down the value of  $p$ , of  $q$  and of  $r$ .

[3 marks]

(b) Find the coefficient of the term in  $x^5$ .

[2 marks]

The answer area is a large rectangle containing 15 horizontal dotted lines for writing. A watermark logo is centered over the lines. The logo consists of a circular border with the text 'SAT PREP' at the top and 'AHMEDABAD' at the bottom. In the center of the circle, the letters 'SP' are written in a large, blue, serif font, with a green leaf-like graphic element behind the 'P'.



4. [Maximum mark: 6]

$$-x - y + z = 2.5$$

Consider the system of equations  $x + y = 1$

$$-2x - y + 2z = -3$$

This system can be represented by the matrix equation  $AX = B$ , where  $X = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ .

(a) (i) Write down the matrix  $A$ .

(ii) Write down the matrix  $A^{-1}$ . [3 marks]


(b) Hence, find  $X$ . [3 marks]



5. [Maximum mark: 8]

The velocity of a particle in  $\text{ms}^{-1}$  is given by  $v = e^{\sin t} - 1$ , for  $0 \leq t \leq 5$ .

(a) On the grid below, sketch the graph of  $v$ . [3 marks]



- (b) (i) Write down the positive  $t$ -intercept.
- (ii) Find the total distance travelled by the particle in the first five seconds. [5 marks]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



6. [Maximum mark: 6]

Let  $f$  and  $g$  be functions such that  $g(x) = 2f(x+1) + 5$ .

(a) The graph of  $f$  is mapped to the graph of  $g$  under the following transformations:

vertical stretch by a factor of  $k$ , followed by a translation  $\begin{pmatrix} p \\ q \end{pmatrix}$ .

Write down the value of

(i)  $k$ ;

(ii)  $p$ ;

(iii)  $q$ .

[3 marks]

(b) Let  $h(x) = -g(3x)$ . The point  $A(6, 5)$  on the graph of  $g$  is mapped to the point  $A'$  on the graph of  $h$ . Find  $A'$ .

[3 marks]



7. [Maximum mark: 7]

A random variable  $X$  is normally distributed with  $\mu = 150$  and  $\sigma = 10$ .

Find the interquartile range of  $X$ .

A large rectangular area containing horizontal dotted lines for writing the answer. A large, faint watermark logo is centered in this area. The logo is circular and contains the text "SAT PREP" at the top and "AHMEDABAD" at the bottom. In the center of the logo are the letters "SP" in a large, stylized font, with a green leaf-like shape behind the "P".






Do **NOT** write solutions on this page.

**SECTION B**

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 14]

The diagram shows a circle of radius 8 metres. The points ABCD lie on the circumference of the circle.



$BC = 14 \text{ m}$ ,  $CD = 11.5 \text{ m}$ ,  $AD = 8 \text{ m}$ ,  $\hat{ADC} = 104^\circ$ , and  $\hat{BCD} = 73^\circ$


- (a) Find AC. [3 marks]
- (b) (i) Find  $\hat{ACD}$ .
- (ii) Hence, find  $\hat{ACB}$ . [5 marks]
- (c) Find the area of triangle ADC. [2 marks]
- (d) Hence or otherwise, find the total area of the shaded regions. [4 marks]



Do **NOT** write solutions on this page.

9. [Maximum mark: 15]

Let  $f(x) = \frac{100}{(1 + 50e^{-0.2x})}$ . Part of the graph of  $f$  is shown below.




- (a) Write down  $f(0)$ . [1 mark]
- (b) Solve  $f(x) = 95$ . [2 marks]
- (c) Find the range of  $f$ . [3 marks]
- (d) Show that  $f'(x) = \frac{1000e^{-0.2x}}{(1 + 50e^{-0.2x})^2}$ . [5 marks]
- (e) Find the maximum rate of change of  $f$ . [4 marks]



Do **NOT** write solutions on this page.

10. [Maximum mark: 16]

A Ferris wheel with diameter 122 metres rotates clockwise at a constant speed. The wheel completes 2.4 rotations every hour. The bottom of the wheel is 13 metres above the ground.



A seat starts at the bottom of the wheel.

(a) Find the maximum height above the ground of the seat. [2 marks]

After  $t$  minutes, the height  $h$  metres above the ground of the seat is given by

$$h = 74 + a \cos bt.$$

(b) (i) Show that the period of  $h$  is 25 minutes.

(ii) Write down the **exact** value of  $b$ . [2 marks]

(c) Find the value of  $a$ . [3 marks]

(d) Sketch the graph of  $h$ , for  $0 \leq t \leq 50$ . [4 marks]

(e) In one rotation of the wheel, find the probability that a randomly selected seat is at least 105 metres above the ground. [5 marks]





Please **do not** write on this page.  
Answers written on this page  
will not be marked.





22137306



**MATHEMATICS**  
**STANDARD LEVEL**  
**PAPER 2**

Candidate session number

0	0								
---	---	--	--	--	--	--	--	--	--

Friday 10 May 2013 (morning)

Examination code

1 hour 30 minutes

2	2	1	3	-	7	3	0	6
---	---	---	---	---	---	---	---	---

## INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL information booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].



0112

Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

**SECTION A**

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 5]

Let  $A = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 1 \\ -1 & 1 & 1 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix}$ .

(a) Write down  $A^{-1}$ . [2 marks]

(b) Solve  $AX = B$ . [3 marks]

Working area with horizontal dotted lines for writing.



2. [Maximum mark: 6]

The random variable  $X$  is normally distributed with mean 20 and standard deviation 5.

(a) Find  $P(X \leq 22.9)$ . [3 marks]

(b) Given that  $P(X < k) = 0.55$ , find the value of  $k$ . [3 marks]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....



3. [Maximum mark: 6]

The following diagram shows a triangle ABC.



*diagram  
not to scale*

The area of triangle ABC is  $80 \text{ cm}^2$ ,  $AB = 18 \text{ cm}$ ,  $AC = x \text{ cm}$  and  $\hat{BAC} = 50^\circ$ .

- (a) Find  $x$ . [3 marks]
- (b) Find BC. [3 marks]

Area for student response with a large watermark reading "SAT PREP SP AHMEDABAD".






4. [Maximum mark: 7]

Line  $L_1$  has equation  $\mathbf{r}_1 = \begin{pmatrix} 10 \\ 6 \\ -1 \end{pmatrix} + s \begin{pmatrix} 2 \\ -5 \\ -2 \end{pmatrix}$  and line  $L_2$  has equation  $\mathbf{r}_2 = \begin{pmatrix} 2 \\ 1 \\ -3 \end{pmatrix} + t \begin{pmatrix} 3 \\ 5 \\ 2 \end{pmatrix}$ .

Lines  $L_1$  and  $L_2$  intersect at point A. Find the coordinates of A.





5. *[Maximum mark: 6]*

The sum of the first three terms of a geometric sequence is 62.755, and the sum of the infinite sequence is 440. Find the common ratio.



6. [Maximum mark: 7]

The constant term in the expansion of  $\left(\frac{x}{a} + \frac{a^2}{x}\right)^6$ , where  $a \in \mathbb{Z}$ , is 1280. Find  $a$ .



7. [Maximum mark: 8]

The following diagram shows a circle with centre O and radius  $r$  cm.




diagram  
not to scale

Points A and B are on the circumference of the circle and  $\hat{AOB} = 1.4$  radians.

The point C is on [OA] such that  $\hat{BCO} = \frac{\pi}{2}$  radians.

(a) Show that  $OC = r \cos 1.4$ .

[1 mark]

(b) The area of the shaded region is  $25 \text{ cm}^2$ . Find the value of  $r$ .

[7 marks]

Area for student response with horizontal dotted lines.



Do **NOT** write solutions on this page.

### SECTION B

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 15]

Consider the points  $A(5, 2, 1)$ ,  $B(6, 5, 3)$ , and  $C(7, 6, a+1)$ , where  $a \in \mathbb{R}$ .

(a) Find

(i)  $\vec{AB}$ ;

(ii)  $\vec{AC}$ .

[3 marks]

Let  $\alpha$  be the angle between  $\vec{AB}$  and  $\vec{AC}$ .

(b) Find the value of  $a$  for which  $\alpha = \frac{\pi}{2}$ .

[4 marks]

(c) (i) Show that  $\cos \alpha = \frac{2a+14}{\sqrt{14a^2+280}}$ .

(ii) Hence, find the value of  $a$  for which  $\alpha = 1.2$ .

[8 marks]



Do **NOT** write solutions on this page.

9. [Maximum mark: 15]

A bag contains four gold balls and six silver balls.

(a) Two balls are drawn at random from the bag, with replacement. Let  $X$  be the number of gold balls drawn from the bag.

(i) Find  $P(X = 0)$ .

(ii) Find  $P(X = 1)$ .

(iii) Hence, find  $E(X)$ .

[8 marks]

Fourteen balls are drawn from the bag, with replacement.

(b) Find the probability that exactly five of the balls are gold.

[2 marks]

(c) Find the probability that at most five of the balls are gold.

[2 marks]

(d) Given that at most five of the balls are gold, find the probability that exactly five of the balls are gold. Give the answer correct to two decimal places.

[3 marks]



Do **NOT** write solutions on this page.

10. [Maximum mark: 15]

Let  $f(x) = e^{\frac{x}{4}}$  and  $g(x) = mx$ , where  $m \geq 0$ , and  $-5 \leq x \leq 5$ . Let  $R$  be the region enclosed by the  $y$ -axis, the graph of  $f$ , and the graph of  $g$ .

(a) Let  $m = 1$ .


(i) Sketch the graphs of  $f$  and  $g$  on the same axes.

(ii) Find the area of  $R$ .

[7 marks]

(b) Consider all values of  $m$  such that the graphs of  $f$  and  $g$  intersect. Find the value of  $m$  that gives the greatest value for the area of  $R$ .

[8 marks]





Please **do not** write on this page.

Answers written on this page  
will not be marked.







88137302



**MATHEMATICS**  
**STANDARD LEVEL**  
**PAPER 2**

Candidate session number

0	0								
---	---	--	--	--	--	--	--	--	--

Tuesday 12 November 2013 (morning)

Examination code

1 hour 30 minutes

8	8	1	3	-	7	3	0	2
---	---	---	---	---	---	---	---	---

## INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL information booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].



12EP01

Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

**SECTION A**

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 5]

Let  $A = \begin{pmatrix} 0 & 5 & 4 \\ 1 & 2 & 1 \\ 2 & 2 & 0 \end{pmatrix}$ , and  $B = \begin{pmatrix} 11 \\ 7 \\ 10 \end{pmatrix}$ .

- (a) Write down  $A^{-1}$ . [2]
- (b) Hence or otherwise, solve the equation  $AX = B$ . [3]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



2. [Maximum mark: 6]

Let  $f(x) = (x-1)(x-4)$ .

- (a) Find the  $x$ -intercepts of the graph of  $f$ . [3]
- (b) The region enclosed by the graph of  $f$  and the  $x$ -axis is rotated  $360^\circ$  about the  $x$ -axis. Find the volume of the solid formed. [3]

A large rectangular box containing horizontal dotted lines for writing answers. A watermark logo for 'SAT PREP SP AHMEDABAD' is centered in the background.



3. [Maximum mark: 6]

$$\text{Let } f(x) = \sqrt[3]{x^4} - \frac{1}{2}.$$

(a) Find  $f'(x)$ . [2]

(b) Find  $\int f(x)dx$ . [4]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



4. [Maximum mark: 6]

Two events  $A$  and  $B$  are such that  $P(A) = 0.2$  and  $P(A \cup B) = 0.5$ .

(a) Given that  $A$  and  $B$  are mutually exclusive, find  $P(B)$ . [2]

(b) Given that  $A$  and  $B$  are independent, find  $P(B)$ . [4]


A large rectangular box containing a faint watermark logo for 'SAT PREP AHMEDABAD' and a large 'SP' emblem. Below the watermark are several horizontal dotted lines intended for writing the solution.



5. [Maximum mark: 8]

A particle moves along a straight line such that its velocity,  $v\text{ms}^{-1}$ , is given by  $v(t) = 10te^{-1.7t}$ , for  $t \geq 0$ .

- (a) On the grid below, sketch the graph of  $v$ , for  $0 \leq t \leq 4$ . [3]



- (b) Find the distance travelled by the particle in the first three seconds. [2]

- (c) Find the velocity of the particle when its acceleration is zero. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



6. *[Maximum mark: 7]*

The time taken for a student to complete a task is normally distributed with a mean of 20 minutes and a standard deviation of 1.25 minutes.


(a) A student is selected at random. Find the probability that the student completes the task in less than 21.8 minutes. [2]

(b) The probability that a student takes between  $k$  and 21.8 minutes is 0.3. Find the value of  $k$ . [5]



7. [Maximum mark: 7]

Consider the graph of the semicircle given by  $f(x) = \sqrt{6x - x^2}$ , for  $0 \leq x \leq 6$ . A rectangle PQRS is drawn with upper vertices R and S on the graph of  $f$ , and PQ on the  $x$ -axis, as shown in the following diagram.



- (a) Let  $OP = x$ .
  - (i) Find PQ, giving your answer in terms of  $x$ .
  - (ii) Hence, write down an expression for the area of the rectangle, giving your answer in terms of  $x$ . [3]
- (b) (i) Find the rate of change of area when  $x = 2$ .
  - (ii) The area is decreasing for  $a < x < b$ . Find the value of  $a$  and of  $b$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....






Do **NOT** write solutions on this page.

**SECTION B**

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 14]

Consider a circle with centre  $O$  and radius  $7$  cm. Triangle  $ABC$  is drawn such that its vertices are on the circumference of the circle.



$AB = 12.2$  cm,  $BC = 10.4$  cm and  $\hat{ACB} = 1.058$  radians.

- (a) Find  $\hat{BAC}$ . [3]
- (b) Find  $AC$ . [5]
- (c) Hence or otherwise, find the length of arc  $ABC$ . [6]




Do **NOT** write solutions on this page.

9. [Maximum mark: 17]

Consider the lines  $L_1$  and  $L_2$  with equations  $L_1: \mathbf{r} = \begin{pmatrix} 11 \\ 8 \\ 2 \end{pmatrix} + s \begin{pmatrix} 4 \\ 3 \\ -1 \end{pmatrix}$  and  $L_2: \mathbf{r} = \begin{pmatrix} 1 \\ 1 \\ -7 \end{pmatrix} + t \begin{pmatrix} 2 \\ 1 \\ 11 \end{pmatrix}$ .

The lines intersect at point P.

- (a) Find the coordinates of P. [6]
- (b) Show that the lines are perpendicular. [5]
- (c) The point  $Q(7, 5, 3)$  lies on  $L_1$ . The point R is the reflection of Q in the line  $L_2$ . Find the coordinates of R. [6]



Do **NOT** write solutions on this page.

10. [Maximum mark: 14]

Samantha goes to school five days a week. When it rains, the probability that she goes to school by bus is 0.5. When it does not rain, the probability that she goes to school by bus is 0.3. The probability that it rains on any given day is 0.2.

- (a) On a randomly selected school day, find the probability that Samantha goes to school by bus. [4]
- (b) Given that Samantha went to school by bus on Monday, find the probability that it was raining. [3]
- (c) In a randomly chosen school week, find the probability that Samantha goes to school by bus on exactly three days. [2]
- (d) After  $n$  school days, the probability that Samantha goes to school by bus at least once is greater than 0.95. Find the smallest value of  $n$ . [5]
- 






Please **do not** write on this page.

Answers written on this page  
will not be marked.



12EP12



22147304



**MATHEMATICS**  
**STANDARD LEVEL**  
**PAPER 2**

Candidate session number

--	--	--	--	--	--	--	--	--	--

Wednesday 14 May 2014 (morning)

Examination code

1 hour 30 minutes

2	2	1	4	–	7	3	0	4
---	---	---	---	---	---	---	---	---

## INSTRUCTIONS TO CANDIDATES


- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].



12EP01



Please **do not** write on this page.  
Answers written on this page  
will not be marked.




Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

**SECTION A**

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 6]

The following diagram shows triangle ABC.



$AB = 6\text{ cm}$ ,  $BC = 10\text{ cm}$ , and  $\hat{A}BC = 100^\circ$ .

(a) Find AC. [3]

(b) Find  $\hat{B}CA$ . [3]

A large rectangular box containing ten horizontal dotted lines for writing the student's solution.







3. [Maximum mark: 7]

The following table shows the average weights ( $y$  kg) for given heights ( $x$  cm) in a population of men.

<b>Heights (<math>x</math> cm)</b>	165	170	175	180	185
<b>Weights (<math>y</math> kg)</b>	67.8	70.0	72.7	75.5	77.2

(a) The relationship between the variables is modelled by the regression equation  $y = ax + b$ .

(i) Write down the value of  $a$  and of  $b$ .

(ii) Hence, estimate the weight of a man whose height is 172 cm. [4]

(b) (i) Write down the correlation coefficient.

(ii) State which **two** of the following describe the correlation between the variables. [3]

- strong
- zero
- positive
- negative
- no correlation
- weak

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....



4. [Maximum mark: 6]

The following diagram shows two perpendicular vectors  $u$  and  $v$ .



(a) Let  $w = u - v$ . Represent  $w$  on the diagram above. [2]

(b) Given that  $u = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}$  and  $v = \begin{pmatrix} 5 \\ n \\ 3 \end{pmatrix}$ , where  $n \in \mathbb{Z}$ , find  $n$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5. [Maximum mark: 6]

The population of deer in an enclosed game reserve is modelled by the function  $P(t) = 210\sin(0.5t - 2.6) + 990$ , where  $t$  is in months, and  $t = 1$  corresponds to 1 January 2014.

(a) Find the number of deer in the reserve on 1 May 2014. [3]

(b) (i) Find the rate of change of the deer population on 1 May 2014.

(ii) Interpret the answer to part (i) with reference to the deer population size on 1 May 2014. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



6. [Maximum mark: 8]

Ramiro and Lautaro are travelling from Buenos Aires to El Moro.


Ramiro travels in a vehicle whose velocity in  $\text{ms}^{-1}$  is given by  $V_R = 40 - t^2$ , where  $t$  is in seconds.

Lautaro travels in a vehicle whose displacement from Buenos Aires in metres is given by  $S_L = 2t^2 + 60$ .

When  $t = 0$ , both vehicles are at the same point.

Find Ramiro's displacement from Buenos Aires when  $t = 10$ .


A large rectangular area containing horizontal dotted lines for writing. A large, faint watermark is centered in the background, featuring the text "SAT PREP" at the top, "SP" in the middle, and "AHMEDABAD" at the bottom, all within a circular border.



7. [Maximum mark: 7]

Let  $f(x) = \frac{g(x)}{h(x)}$ , where  $g(2) = 18$ ,  $h(2) = 6$ ,  $g'(2) = 5$ , and  $h'(2) = 2$ . Find the equation of the normal to the graph of  $f$  at  $x = 2$ .

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



Turn over


Do **NOT** write solutions on this page.

**SECTION B**

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 16]

The weights in grams of 80 rats are shown in the following cumulative frequency diagram.



(This question continues on the following page)



Do **NOT** write solutions on this page.

(Question 8 continued)

- (a) (i) Write down the median weight of the rats.
- (ii) Find the percentage of rats that weigh 70 grams or less. [4]

The same data is presented in the following table.

<b>Weights <math>w</math> grams</b>	$0 \leq w \leq 30$	$30 < w \leq 60$	$60 < w \leq 90$	$90 < w \leq 120$
<b>Frequency</b>	$p$	45	$q$	5

- (b) (i) Write down the value of  $p$ .
- (ii) Find the value of  $q$ . [4]
- (c) Use the values from the table to estimate the mean and standard deviation of the weights. [3]

Assume that the weights of these rats are normally distributed with the mean and standard deviation estimated in part (c).

- (d) Find the percentage of rats that weigh 70 grams or less. [2]
- (e) A sample of five rats is chosen at random. Find the probability that at most three rats weigh 70 grams or less. [3]



Do **NOT** write solutions on this page.

9. [Maximum mark: 15]

Let  $f(x) = \cos\left(\frac{\pi}{4}x\right) + \sin\left(\frac{\pi}{4}x\right)$ , for  $-4 \leq x \leq 4$ .

(a) Sketch the graph of  $f$ . [3]

(b) Find the values of  $x$  where the function is decreasing. [5]

(c) The function  $f$  can also be written in the form  $f(x) = a \sin\left(\frac{\pi}{4}(x+c)\right)$ , where  $a \in \mathbb{R}$ , and  $0 \leq c \leq 2$ . Find the value of

(i)  $a$ ;

(ii)  $c$ . [7]

10. [Maximum mark: 14]

Let  $f(x) = \frac{3x}{x-q}$ , where  $x \neq q$ .


(a) Write down the equations of the vertical and horizontal asymptotes of the graph of  $f$ . [2]

The vertical and horizontal asymptotes to the graph of  $f$  intersect at the point  $Q(1, 3)$ .

(b) Find the value of  $q$ . [2]

(c) The point  $P(x, y)$  lies on the graph of  $f$ . Show that  $PQ = \sqrt{(x-1)^2 + \left(\frac{3}{x-1}\right)^2}$ . [4]

(d) Hence find the coordinates of the points on the graph of  $f$  that are closest to  $(1, 3)$ . [6]







22147306



**MATHEMATICS**  
**STANDARD LEVEL**  
**PAPER 2**

Candidate session number

--	--	--	--	--	--	--	--	--	--

Wednesday 14 May 2014 (morning)

Examination code

1 hour 30 minutes

2	2	1	4	-	7	3	0	6
---	---	---	---	---	---	---	---	---

## INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].



12EP01


Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

**SECTION A**

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 6]

The following diagram shows a circle with centre O and radius 5 cm.



The points A, B and C lie on the circumference of the circle, and  $\widehat{AOC} = 0.7$  radians.


- (a) (i) Find the length of the arc ABC.
- (ii) Find the perimeter of the shaded sector. [4]
- (b) Find the area of the shaded sector. [2]

*(This question continues on the following page)*



*(Question 1 continued)*

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....




12EP03

**Turn over**

2. [Maximum mark: 6]

Let  $f(x) = 5 - x^2$ . Part of the graph of  $f$  is shown in the following diagram.



The graph crosses the  $x$ -axis at the points A and B.

- (a) Find the  $x$ -coordinate of A and of B. [3]
- (b) The region enclosed by the graph of  $f$  and the  $x$ -axis is revolved  $360^\circ$  about the  $x$ -axis. Find the volume of the solid formed. [3]

Area for student response with horizontal dotted lines.



3. [Maximum mark: 5]

The following table shows the amount of fuel ( $y$  litres) used by a car to travel certain distances ( $x$  km).

<b>Distance</b> ( $x$ km)	40	75	120	150	195
<b>Amount of fuel</b> ( $y$ litres)	3.6	6.5	9.9	13.1	16.2

This data can be modelled by the regression line with equation  $y = ax + b$ .

- (a) (i) Write down the value of  $a$  and of  $b$ .  
  
(ii) Explain what the gradient  $a$  represents. [3]
- (b) Use the model to estimate the amount of fuel the car would use if it is driven 110 km. [2]

Area with horizontal dotted lines for student answers.




4. [Maximum mark: 7]

Let  $A$  and  $B$  be independent events, where  $P(A) = 0.3$  and  $P(B) = 0.6$ .

(a) Find  $P(A \cap B)$ . [2]

(b) Find  $P(A \cup B)$ . [2]

(c) (i) On the following Venn diagram, shade the region that represents  $A \cap B'$ .



(ii) Find  $P(A \cap B')$ . [3]

A large rectangular area containing horizontal dotted lines for writing the answer to part (c)(ii).



5. [Maximum mark: 7]

In triangle ABC,  $AB = 6\text{ cm}$  and  $AC = 8\text{ cm}$ . The area of the triangle is  $16\text{ cm}^2$ .

(a) Find the two possible values for  $\hat{A}$ .

[4]

(b) Given that  $\hat{A}$  is obtuse, find BC.


[3]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



6. [Maximum mark: 8]

Let  $f(x) = p \cos(q(x+r)) + 10$ , for  $0 \leq x \leq 20$ . The following diagram shows the graph of  $f$ .



The graph has a maximum at  $(4, 18)$  and a minimum at  $(16, 2)$ .

- (a) Write down the value of  $r$ . [2]
- (b) (i) Find  $p$ .  
(ii) Find  $q$ . [4]
- (c) Solve  $f(x) = 7$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



7. [Maximum mark: 7]

Consider the expansion of  $x^2 \left( 3x^2 + \frac{k}{x} \right)^8$ . The constant term is 16 128.

Find  $k$ .



Do **NOT** write solutions on this page.

### SECTION B

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 15]

The number of bacteria in two colonies, A and B, starts increasing at the same time.

The number of bacteria in colony A after  $t$  hours is modelled by the function  $A(t) = 12e^{0.4t}$ .

- (a) Find the initial number of bacteria in colony A. [2]
- (b) Find the number of bacteria in colony A after four hours. [3]
- (c) How long does it take for the number of bacteria in colony A to reach 400? [3]

The number of bacteria in colony B after  $t$  hours is modelled by the function  $B(t) = 24e^{kt}$ .

- (d) After four hours, there are 60 bacteria in colony B. Find the value of  $k$ . [3]
- (e) The number of bacteria in colony A first exceeds the number of bacteria in colony B after  $n$  hours, where  $n \in \mathbb{Z}$ . Find the value of  $n$ . [4]



Do **NOT** write solutions on this page.

9. [Maximum mark: 15]

A particle moves in a straight line. Its velocity,  $v \text{ ms}^{-1}$ , at time  $t$  seconds, is given by

$$v = (t^2 - 4)^3, \text{ for } 0 \leq t \leq 3.$$

- (a) Find the velocity of the particle when  $t = 1$ . [2]
- (b) Find the value of  $t$  for which the particle is at rest. [3]
- (c) Find the total distance the particle travels during the first three seconds. [3]
- (d) Show that the acceleration of the particle is given by  $a = 6t(t^2 - 4)^2$ . [3]
- (e) Find all possible values of  $t$  for which the velocity and acceleration are both positive or both negative. [4]




Do **NOT** write solutions on this page.

10. [Maximum mark: 14]

A forest has a large number of tall trees. The heights of the trees are normally distributed with a mean of 53 metres and a standard deviation of 8 metres. Trees are classified as giant trees if they are more than 60 metres tall.

- (a) A tree is selected at random from the forest.
- (i) Find the probability that this tree is a giant.
- (ii) Given that this tree is a giant, find the probability that it is taller than 70 metres. [6]
- (b) Two trees are selected at random. Find the probability that they are both giants. [2]
- (c) 100 trees are selected at random.
- (i) Find the expected number of these trees that are giants.
- (ii) Find the probability that at least 25 of these trees are giants. [6]
- 





88147302



**MATHEMATICS**  
**STANDARD LEVEL**  
**PAPER 2**

Candidate session number

--	--	--	--	--	--	--	--	--	--

Thursday 13 November 2014 (morning)

Examination code

1 hour 30 minutes

8	8	1	4	-	7	3	0	2
---	---	---	---	---	---	---	---	---

## INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].



16EP01



Please **do not** write on this page.

Answers written on this page  
will not be marked.



Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

**SECTION A**

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 5]

Let  $f(x) = 2x + 3$  and  $g(x) = x^3$ .

(a) Find  $(f \circ g)(x)$ . [2]

(b) Solve the equation  $(f \circ g)(x) = 0$ . [3]



2. [Maximum mark: 6]

The following table shows the Diploma score  $x$  and university entrance mark  $y$  for seven IB Diploma students.

Diploma score ( $x$ )	28	30	27	31	32	25	27
University entrance mark ( $y$ )	73.9	78.1	70.2	82.2	85.5	62.7	69.4

(a) Find the correlation coefficient. [2]

The relationship can be modelled by the regression line with equation  $y = ax + b$ .

(b) Write down the value of  $a$  and of  $b$ . [2]

Rita scored a total of 26 in her IB Diploma.

(c) Use your regression line to estimate Rita's university entrance mark. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....





3. [Maximum mark: 7]

The following diagram shows a circle with centre O and radius 8 cm.



The points A, B and C are on the circumference of the circle, and  $\hat{AOB} = 1.2$  radians.

- (a) Find the length of arc ACB. [2]
- (b) Find AB. [3]
- (c) Hence, find the perimeter of the shaded segment ABC. [2]

Handwritten answer area with horizontal dotted lines for writing.




4. [Maximum mark: 8]

Let  $f(x) = -x^4 + 2x^3 - 1$ , for  $0 \leq x \leq 2$ .

(a) Sketch the graph of  $f$  on the following grid.

[3]



(b) Solve  $f(x) = 0$ .

[2]

(c) The region enclosed by the graph of  $f$  and the  $x$ -axis is rotated  $360^\circ$  about the  $x$ -axis. Find the volume of the solid formed.


[3]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



5. [Maximum mark: 7]

The following diagram shows part of the graph of  $y = p \sin(qx) + r$ .



The point  $A\left(\frac{\pi}{6}, 2\right)$  is a maximum point and the point  $B\left(\frac{\pi}{2}, 1\right)$  is a minimum point.  
Find the value of

- (a)  $p$ ; [2]
- (b)  $r$ ; [2]
- (c)  $q$ . [3]

Area for student response with horizontal dotted lines.



6. [Maximum mark: 6]

Consider the expansion of  $\left(\frac{x^3}{2} + \frac{p}{x}\right)^8$ . The constant term is 5103. Find the possible values of  $p$ .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....



7. [Maximum mark: 6]

A particle starts from point A and moves along a straight line. Its velocity,  $v \text{ ms}^{-1}$ , after  $t$  seconds is given by  $v(t) = e^{\frac{1}{2}\cos t} - 1$ , for  $0 \leq t \leq 4$ . The particle is at rest when  $t = \frac{\pi}{2}$ .

The following diagram shows the graph of  $v$ .



- (a) Find the distance travelled by the particle for  $0 \leq t \leq \frac{\pi}{2}$ . [2]
- (b) Explain why the particle passes through A again. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....




Do **NOT** write solutions on this page.

**SECTION B**

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 15]

The following cumulative frequency graph shows the monthly income,  $I$  dollars, of 2000 families.



(This question continues on the following page)



Do **NOT** write solutions on this page.

(Question 8 continued)

- (a) Find the median monthly income. [2]
- (b) (i) Write down the number of families who have a monthly income of 2000 dollars or less.
- (ii) Find the number of families who have a monthly income of more than 4000 dollars. [4]

The 2000 families live in two different types of housing. The following table gives information about the number of families living in each type of housing and their monthly income  $I$ .

	$1000 < I \leq 2000$	$2000 < I \leq 4000$	$4000 < I \leq 5000$
Apartment	436	765	28
Villa	64	$p$	122

- (c) Find the value of  $p$ . [2]
- (d) A family is chosen at random.
- (i) Find the probability that this family lives in an apartment.
- (ii) Find the probability that this family lives in an apartment, given that its monthly income is greater than 4000 dollars. [4]
- (e) Estimate the mean monthly income for families living in a villa. [3]



Do **NOT** write solutions on this page.

9. [Maximum mark: 14]

The first two terms of a geometric sequence  $u_n$  are  $u_1 = 4$  and  $u_2 = 4.2$ .

(a) (i) Find the common ratio.

(ii) Hence or otherwise, find  $u_5$ .

[5]

Another sequence  $v_n$  is defined by  $v_n = an^k$ , where  $a, k \in \mathbb{R}$ , and  $n \in \mathbb{Z}^+$ , such that  $v_1 = 0.05$  and  $v_2 = 0.25$ .


(b) (i) Find the value of  $a$ .

(ii) Find the value of  $k$ .

[5]

(c) Find the smallest value of  $n$  for which  $v_n > u_n$ .

[4]





Do **NOT** write solutions on this page.

10. [Maximum mark: 16]

The weights of fish in a lake are normally distributed with a mean of 760 g and standard deviation  $\sigma$ . It is known that 78.87% of the fish have weights between 705 g and 815 g.

- (a) (i) Write down the probability that a fish weighs more than 760 g.
- (ii) Find the probability that a fish weighs less than 815 g. [4]
- (b) (i) Write down the standardized value for 815 g.
- (ii) Hence or otherwise, find  $\sigma$ . [4]

A fishing contest takes place in the lake. Small fish, called tiddlers, are thrown back into the lake. The maximum weight of a tiddler is 1.5 standard deviations below the mean.

- (c) Find the maximum weight of a tiddler. [2]
- (d) A fish is caught at random. Find the probability that it is a tiddler. [2]
- (e) 25% of the fish in the lake are salmon. 10% of the salmon are tiddlers. Given that a fish caught at random is a tiddler, find the probability that it is a salmon. [4]





Please **do not** write on this page.  
Answers written on this page  
will not be marked.





Please **do not** write on this page.  
Answers written on this page  
will not be marked.





Please **do not** write on this page.  
Answers written on this page  
will not be marked.




16EP16

ISBN 978-1-78638-568-0



International Baccalaureate®  
Baccalauréat International  
Bachillerato Internacional

# Mathematics\_paper\_2\_\_T Z1\_SL



**Mathematics**  
**Standard level**  
**Paper 2**

Wednesday 13 May 2015 (afternoon)

Candidate session number

1 hour 30 minutes

--	--	--	--	--	--	--	--	--	--

**Instructions to candidates**

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[90 marks]**.



Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

### Section A

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 7]

The following table shows the average number of hours per day spent watching television by seven mothers and each mother’s youngest child.

<b>Hours per day that a mother watches television (<i>x</i>)</b>	2.5	3.0	3.2	3.3	4.0	4.5	5.8
<b>Hours per day that her child watches television (<i>y</i>)</b>	1.8	2.2	2.6	2.5	3.0	3.2	3.5

The relationship can be modelled by the regression line with equation  $y = ax + b$ .

- (a) (i) Find the correlation coefficient.
- (ii) Write down the value of  $a$  and of  $b$ . [4]

Elizabeth watches television for an average of 3.7 hours per day.

- (b) Use your regression line to predict the average number of hours of television watched per day by Elizabeth’s youngest child. Give your answer correct to one decimal place. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





2. [Maximum mark: 5]

Consider the expansion of  $(2x + 3)^8$ .

- (a) Write down the number of terms in this expansion. [1]
  
- (b) Find the term in  $x^3$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Turn over



3. [Maximum mark: 6]

In an arithmetic sequence  $u_{10} = 8$ ,  $u_{11} = 6.5$ .

(a) Write down the value of the common difference. [1]

(b) Find the first term. [3]

(c) Find the sum of the first 50 terms of the sequence. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



A large, light grey watermark logo is centered on the page. It consists of a circular border containing the text 'SAT PREP' at the top and 'AHMEDABAD' at the bottom. In the center of the circle are the letters 'SP' in a large, stylized font, with a green leaf-like shape behind the 'P'.



4. [Maximum mark: 7]

Let  $f(x) = \frac{2x-6}{1-x}$ , for  $x \neq 1$ .

(a) For the graph of  $f$

(i) find the  $x$ -intercept;

(ii) write down the equation of the vertical asymptote;

(iii) find the equation of the horizontal asymptote.

[5]

(b) Find  $\lim_{x \rightarrow \infty} f(x)$ .

[2]

A large rectangular area for writing answers, containing horizontal dotted lines and a large watermark logo for 'SAT PREP AHMEDABAD' with 'SP' in the center.




5. [Maximum mark: 6]

Let  $G(x) = 95e^{(-0.02x)} + 40$ , for  $20 \leq x \leq 200$ .

(a) On the following grid, sketch the graph of  $G$ .

[3]



(b) Robin and Pat are planning a wedding banquet. The cost per guest,  $G$  dollars, is modelled by the function  $G(n) = 95e^{(-0.02n)} + 40$ , for  $20 \leq n \leq 200$ , where  $n$  is the number of guests.

Calculate the **total** cost for 45 guests.


[3]

(This question continues on the following page)



(Question 5 continued)

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



6. [Maximum mark: 7]

Let  $f(x) = \frac{\ln(4x)}{x}$ , for  $0 < x \leq 5$ .

Points P(0.25, 0) and Q are on the curve of  $f$ . The tangent to the curve of  $f$  at P is perpendicular to the tangent at Q. Find the coordinates of Q.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....



7. [Maximum mark: 7]

The following diagram shows part of the graph of  $f(x) = -2x^3 + 5.1x^2 + 3.6x - 0.4$ .



(a) Find the coordinates of the local minimum point. [2]

(b) The graph of  $f$  is translated to the graph of  $g$  by the vector  $\begin{pmatrix} 0 \\ k \end{pmatrix}$ . Find all values of  $k$  so that  $g(x) = 0$  has exactly one solution. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



16EP09


Do **not** write solutions on this page.

**Section B**

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 13]


The following diagram shows a straight shoreline, with a supply store at S, a town at T, and an island L.



A boat delivers supplies to the island. The boat leaves S, and sails to the island. Its path makes an angle of  $20^\circ$  with the shoreline.

(a) The boat sails at 6 km per hour, and arrives at L after 1.5 hours. Find the distance from S to L. [2]

It is decided to change the position of the supply store, so that its distance from L is 5 km. The following diagram shows the two possible locations P and Q for the supply store.



(b) Find the size of  $\hat{SPL}$  and of  $\hat{SQL}$ . [5]

(c) The town wants the new supply store to be as near as possible to the town.

(i) State which of the points P or Q is chosen for the new supply store.

(ii) Hence find the distance between the old supply store and the new one. [6]



Do **not** write solutions on this page.

9. [Maximum mark: 16]

A company makes containers of yogurt. The volume of yogurt in the containers is normally distributed with a mean of 260 ml and standard deviation of 6 ml.

A container which contains less than 250 ml of yogurt is **underfilled**.

(a) A container is chosen at random. Find the probability that it is underfilled. [2]

The company decides that the probability of a container being underfilled should be reduced to 0.02. It decreases the standard deviation to  $\sigma$  and leaves the mean unchanged.

(b) Find  $\sigma$ . [4]

The company changes to the new standard deviation,  $\sigma$ , and leaves the mean unchanged. A container is chosen at random for inspection. It passes inspection if its volume of yogurt is between 250 and 271 ml.

(c) (i) Find the probability that it passes inspection.  
(ii) Given that the container is **not** underfilled, find the probability that it passes inspection. [6]

(d) A sample of 50 containers is chosen at random. Find the probability that 48 or more of the containers pass inspection. [4]






Do **not** write solutions on this page.

10. [Maximum mark: 16]

Consider a function  $f$ , for  $0 \leq x \leq 10$ . The following diagram shows the graph of  $f'$ , the derivative of  $f$ .



The graph of  $f'$  passes through  $(2, -2)$  and  $(5, 1)$ , and has  $x$ -intercepts at 0, 4 and 6.

(a) The graph of  $f$  has a local maximum point when  $x = p$ . State the value of  $p$ , and justify your answer. [3]

(b) Write down  $f'(2)$ . [1]

Let  $g(x) = \ln(f(x))$  and  $f(2) = 3$ .

(c) Find  $g'(2)$ . [4]

(d) Verify that  $\ln 3 + \int_2^a g'(x) dx = g(a)$ , where  $0 \leq a \leq 10$ . [4]


(This question continues on the following page)



Do **not** write solutions on this page.

**(Question 10 continued)**

(e) The following diagram shows the graph of  $g'$ , the derivative of  $g$ .



The shaded region  $A$  is enclosed by the curve, the  $x$ -axis and the line  $x = 2$ , and has area  $0.66 \text{ units}^2$ .

The shaded region  $B$  is enclosed by the curve, the  $x$ -axis and the line  $x = 5$ , and has area  $0.21 \text{ units}^2$ .

Find  $g(5)$ .

[4]





Please **do not** write on this page.  
Answers written on this page  
will not be marked.





Please **do not** write on this page.  
Answers written on this page  
will not be marked.





Please **do not** write on this page.  
Answers written on this page  
will not be marked.



16EP16


# Markscheme

**May 2015**

**Mathematics**

**Standard level**

**Paper 2**



This markscheme is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of the IB Assessment Centre.



## Instructions to Examiners

### Abbreviations

- M** Marks awarded for attempting to use a valid **Method**; working must be seen.
- (M)** Marks awarded for a valid **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

### Using the markscheme

#### 1 General

Mark according to RM assessor instructions and the document “**Mathematics SL: Guidance for e-marking May 2015**”. It is **essential** that you read this document before you start marking. In particular, please note the following. Marks must be recorded using the annotation stamps, using the RM assessor tool. Please check that you are entering marks for the right question. All the marks will be added and recorded by RM assessor.

If a part is **completely correct**, (and gains all the “must be seen” marks), use the ticks with numbers to stamp full marks. Do **not** use the ticks with numbers for anything else.

- If a part is completely wrong, stamp **A0** by the final answer.
- If a part gains anything else, all the working **must** have annotations stamped to show what marks are awarded. This includes any zero marks.

#### 2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award **M0** followed by **A1**, as **A** mark(s) depend on the preceding **M** mark(s), if any. An exception to this rule is when work for **M1** is missing, as opposed to incorrect (see point 4).
- Where **M** and **A** marks are noted on the same line, eg **M1A1**, this usually means **M1** for an **attempt** to use an appropriate method (eg substitution into a formula) and **A1** for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **AOA1A1**.
- Where the markscheme specifies (**M2**), **N3**, etc., do **not** split the marks, unless there is a note.
- Most **M** marks are for a **valid** method, ie a method which can lead to the answer: it must indicate some form of progress towards the answer.
- Once a correct answer to a question or part-question is seen, ignore further correct working. However, if further working indicates a lack of mathematical understanding do not award the final **A1**. An exception to this may be in numerical answers, where a correct exact value is followed by an incorrect decimal (see examples on next page).



## Examples

	Correct answer seen	Further working seen	Action
1.	$8\sqrt{2}$	5.65685 (incorrect decimal value)	Award the final <b>A1</b> (ignore the further working)
2.	$\frac{1}{4}\sin 4x$	$\sin x$	Do not award the final <b>A1</b>
3.	$\log a - \log b$	$\log(a - b)$	Do not award the final <b>A1</b>

### 3 N marks

If **no working shown**, award **N** marks for **correct** answers – this includes acceptable answers (see accuracy booklet). In this case, ignore mark breakdown (**M**, **A**, **R**). Where a student only shows a final incorrect answer with no working, even if that answer is a correct intermediate answer, award **NO**.

- Do **not** award a mixture of **N** and other marks.
- There may be fewer **N** marks available than the total of **M**, **A** and **R** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.
- There may not be a direct relationship between the **N** marks and the implied marks. There are times when all the marks are implied, but the **N** marks are not the full marks: this indicates that we want to see some of the working, without specifying what.
- For consistency within the markscheme, **N** marks are noted for every part, even when these match the mark breakdown.
- If a candidate has incorrect working, which somehow results in a correct answer, do **not** award the **N** marks for this correct answer. However, if the candidate has indicated (usually by crossing out) that the working is to be ignored, award the **N** marks for the correct answer.

### 4 Implied and must be seen marks

*Implied marks appear in **brackets** eg (M1).*

- Implied marks can only be awarded if the work is seen or if implied in subsequent working (a correct final answer does not necessarily mean that the implied marks are all awarded). There are questions where some working is required, but as it is accepted that not everyone will write the same steps, all the marks are implied, but the **N** marks are not the full marks for the question.
- Normally the correct work is seen in the next line.
- Where there is an (**M1**) followed by **A1** for each correct answer, if no working shown, one correct answer is sufficient evidence to award the (**M1**).

*Must be seen marks appear without **brackets** eg M1.*

- Must be seen marks can only be awarded if the work is seen.
- If a must be seen mark is not awarded because work is missing (as opposed to **M0** or **A0** for incorrect work) all subsequent marks may be awarded if appropriate.

### 5 Follow through marks (only applied after an error is made)

*Follow through (FT) marks are awarded where an incorrect answer (final or intermediate) from one **part** of a question is used correctly in **subsequent** part(s) or subpart(s). Usually, to award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if the only marks awarded in a subpart are for the **final** answer, then **FT** marks should be awarded if appropriate. Examiners are expected to check student work in order to award **FT** marks where appropriate.*

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** and **R** marks may be awarded if appropriate. (However, as noted above, if an **A** mark is not awarded because work is missing, all subsequent marks may be awarded if appropriate).
- Exceptions to this rule will be explicitly noted on the markscheme.
- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (eg probability greater than 1, use of  $r > 1$  for the sum of an infinite GP,  $\sin \theta = 1.5$ , non integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word “**their**” in a description, to indicate that candidates may be using an incorrect value.
- If a candidate makes an error in one part, but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the question says hence. It is often possible to use a different approach in subsequent parts that does not depend on the answer to previous parts.
- In a “show that” question, if an error in a previous subpart leads to not showing the required answer, do not award the final **A1**. Note that if the error occurs within the same subpart, the **FT** rules may result in further loss of marks.
- Where there are anticipated common errors, the **FT** answers are often noted on the markscheme, to help examiners. It should be stressed that these are not the only **FT** answers accepted, neither should **N** marks be awarded for these answers.

## 6 Mis-read

*If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). A candidate should be penalized only once for a particular mis-read. Use the **MR** stamp to indicate that this is a misread. Do not award the first mark in the question, even if this is an **M** mark, but award all others (if appropriate) so that the candidate only loses one mark for the misread.*

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (eg probability greater than 1, use of  $r > 1$  for the sum of an infinite GP,  $\sin \theta = 1.5$ , non integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates’ own work does **not** constitute a misread, it is an error.

## 7 Discretionary marks (**d**)

*An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. In such cases the annotation **DM** should be used and a brief **note** written next to the mark explaining this decision.*

## 8 Alternative methods

*Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.*

- Alternative methods for complete parts are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for parts of questions are indicated by **EITHER . . . OR**. Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

## 9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

## 10 Calculators

A GDC is required for paper 2, but calculators with symbolic manipulation features (eg TI-89) are not allowed.

**Calculator notation** The mathematics SL guide says:

*Students must always use correct mathematical notation, not calculator notation.*

Do **not** accept final answers written using calculator notation. However, do not penalize the use of calculator notation in the working.

## 11 Style

The markscheme aims to present answers using good communication, eg if the question asks to find the value of  $k$ , the markscheme will say  $k = 3$ , but the marks will be for the correct value 3 – there is usually no need for the “ $k =$ ”. In these cases, it is also usually acceptable to have another variable, as long as there is no ambiguity in the question, eg if the question asks to find the value of  $p$  and of  $q$ , then the student answer needs to be clear. Generally, the only situation where the full answer is required is in a question which asks for equations – in this case the markscheme will say “must be an equation”. Accept sloppy notation in the working, where this is followed by correct working eg  $-2^2 = 4$  where they should have written  $(-2)^2 = 4$ .

The markscheme often uses words to describe what the marks are for, followed by examples, using the eg notation. These examples are not exhaustive, and examiners should check what candidates have written, to see if they satisfy the description. Where these marks are **M** marks, the examples may include ones using poor notation, to indicate what is acceptable. A valid method is one which will allow candidate to proceed to the next step eg if a quadratic function is given in factorised form, and the question asks for the zeroes, then multiplying the factors does not necessarily help to find the zeros, and would not on its own count as a valid method.

## 12 Candidate work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

Candidates are meant to write their answers to Section A on the question paper (QP), and Section B on answer booklets. Sometimes, they need more room for Section A, and use the booklet (and often comment to this effect on the QP), or write outside the box. That is fine, and this work should be marked.

The instructions tell candidates not to write on Section B of the QP. Thus they may well have done some rough work here which they assume will be ignored. If they have solutions on the answer booklets, there is no need to look at the QP. However, if there are whole questions or whole part solutions missing on answer booklets, please check to make sure that they are not on the QP, and if they are, mark those whole questions or whole part solutions that have not been written on answer booklets.

### 13. Diagrams

The notes on how to allocate marks for sketches usually refer to passing through particular points or having certain features. These marks can only be awarded if the sketch is approximately the correct shape. All values given will be an approximate guide to where these points/features occur. In some questions, the first **A1** is for the shape, in others, the marks are only for the points and/or features. In both cases, unless the shape is approximately correct, no marks can be awarded (unless otherwise stated). However, if the graph is based on previous calculations, **FT** marks should be awarded if appropriate.

### 14. Accuracy of Answers

*If the level of accuracy is specified in the question, a mark will be allocated for giving the final answer to the required accuracy. When this is not specified in the question, all numerical answers should be given exactly or correct to three significant figures.*

Do not accept unfinished numerical final answers such as  $3/0.1$  (unless otherwise stated). As a rule, numerical answers with more than one part (such as fractions) should be given using integers (eg  $6/8$ ). Calculations which lead to integers should be completed, with the exception of fractions which are not whole numbers.

Intermediate values do not need to be given to the correct three significant figures. But, if candidates work with rounded values, this could lead to an incorrect answer, in which case award **A0** for the final answer.

Where numerical answers are required as the **final** answer to a part of a question in the markscheme, the markscheme will show

a truncated 6 sf value, the exact value if applicable, and the correct 3 sf answer.

Units (which are generally not required) will appear in brackets at the end.

**Section A**

1. (a) (i) evidence of valid approach **(M1)**  
 eg 1 correct value for  $r$ , (or for  $a$  or  $b$ , seen in (ii))  
 $0.946591$   
 $r = 0.947$  **A1 N2**
- (ii)  $a = 0.500957, b = 0.803544$   
 $a = 0.501, b = 0.804$  **A1A1 N2**  
**[4 marks]**
- (b) substituting  $x = 3.7$  into **their** equation **(M1)**  
 eg  $0.501(3.7) + 0.804$   
 $2.65708$  (2 hours 39.4252 minutes) **(A1)**  
 $y = 2.7$  (hours)(**must** be correct 1 dp, accept 2 hours 39.4 minutes) **A1 N3**  
**[3 marks]**
- Total [7marks]**
2. (a) 9 terms **A1 N1**  
**[1 mark]**
- (b) valid approach to find the required term **(M1)**  
 eg  $\binom{8}{r}(2x)^{8-r}(3)^r, (2x)^8(3)^0 + (2x)^7(3)^1 + \dots$ , Pascal's triangle to  
 8<sup>th</sup> row  
 identifying correct term (may be indicated in expansion) **(A1)**  
 eg 6th term,  $r = 5, \binom{8}{5}, (2x)^3(3)^5$   
 correct working (may be seen in expansion) **(A1)**  
 eg  $\binom{8}{5}(2x)^3(3)^5, 56 \times 2^3 \times 3^5$   
 $108864x^3$  (accept  $109000x^3$ ) **A1 N3**  
**[4 marks]**


**Notes:** Do not award any marks if there is clear evidence of adding instead of multiplying.  
 Do not award final **A1** for a final answer of  $108864$ , even if  $108864x^3$  is seen previously.  
 If no working shown award **N2** for  $108864$ .

**Total [5 marks]**



3. (a)  $d = -1.5$  A1 N1  
[1 mark]
- (b) **METHOD 1**
- valid approach (M1)  
 eg  $u_{10} = u_1 + 9d$ ,  $8 = u_1 - 9(-1.5)$
- correct working (A1)  
 eg  $8 = u_1 + 9d$ ,  $6.5 = u_1 + 10d$ ,  $u_1 = 8 - 9(-1.5)$
- $u_1 = 21.5$  A1 N2
- METHOD 2**
- attempt to list 3 or more terms in either direction (M1)  
 eg 9.5, 11, 12.5, ...; 5, 3.5, 2, ... ..
- correct list of 4 or more terms in **correct** direction (A1)  
 eg 9.5, 11, 12.5, 14
- $u_1 = 21.5$  A1 N2  
[3 marks]
- (c) correct expression (A1)  
 eg  $\frac{50}{2}(2(21.5) + 49(-1.5))$ ,  $\frac{50}{2}(21.5 - 52)$ ,  $\sum_{k=1}^{50} 21.5 + (k-1)(-1.5)$
- sum = -762.5 (exact) A1 N2  
[2 marks]
- Total [ 6 marks]**
4. (a) (i) valid approach (M1)  
 eg sketch,  $f(x) = 0$ ,  $0 = 2x - 6$   
 $x = 3$  or (3, 0) A1 N2
- (ii)  $x = 1$  (must be equation) A1 N1
- (iii) valid approach (M1)  
 eg sketch,  $\frac{2x}{-1x}$ , inputting large values of  $x$ , L'Hopital's rule  
 $y = -2$  (must be equation) A1 N2  
[5 marks]
- (b) valid approach (M1)  
 eg recognizing that  $\lim_{x \rightarrow \infty}$  is related to the horizontal asymptote,  
 table with large values of  $x$ , their  $y$  value from (a)(iii), L'Hopital's rule
- $\lim_{x \rightarrow \infty} f(x) = -2$  A1 N2  
[2 marks]
- Total [ 7 marks]**

5. (a)



A1A1A1

N3

**Note:** Curve must be approximately correct exponential shape (concave up and decreasing). Only if the shape is approximately correct, award the following:  
**A1** for left endpoint in circle,  
**A1** for right endpoint in circle,  
**A1** for asymptotic to  $y = 40$  (must not go below  $y = 40$ ).

[3 marks]

(b) attempt to find  $G(45)$

(M1)

eg 78.6241, value read from **their** graph

multiplying cost times number of people

(M1)

eg  $45 \times 78.6241$ ,  $G(45) \times 45$

3538.08

3540 (dollars)

A1

N2

[3 marks]

Total [ 6 marks]

6. recognizing that the gradient of tangent is the derivative (M1)  
 eg  $f'$
- finding the gradient of  $f$  at P (A1)  
 eg  $f'(0.25) = 16$
- evidence of taking negative reciprocal of **their** gradient at P (M1)  
 eg  $\frac{-1}{m}, -\frac{1}{f'(0.25)}$
- equating derivatives M1
- eg  $f'(x) = \frac{-1}{16}, f' = -\frac{1}{m}, \frac{x\left(\frac{1}{x}\right) - \ln(4x)}{x^2} = 16$
- finding the  $x$ -coordinate of Q,  $x = 0.700750$   
 $x = 0.701$  A1 N3
- attempt to substitute **their**  $x$  into  $f$  to find the  $y$ -coordinate of Q (M1)  
 eg  $f(0.7)$
- $y = 1.47083$   
 $y = 1.47$  A1 N2  
[7 marks]
7. (a)  $(-0.3, -0.967)$   
 $x = -0.3$  (exact),  $y = -0.967$  (exact) A1A1 N2  
[2 marks]
- (b)  $y$ -coordinate of local maximum is  $y = 11.2$  (A1)
- negating the  $y$ -coordinate of one of the max/min (M1)  
 eg  $y = 0.967, y = -11.2$
- recognizing that the solution set has two intervals R1  
 eg two answers,
- $k < -11.2, k > 0.967$  A1A1 N3N2  
[5 marks]

**Notes:** If working shown, do not award the final mark if strict inequalities are not used.  
 If no working shown, award **N2** for  $k \leq -11.2$  or **N1** for  $k \geq 0.967$


**Total [ 7 marks]**



**Section B**

8. (a) valid approach (M1)  
 eg  $\text{speed} = \frac{\text{distance}}{\text{time}}, 6 \times 1.5$   
 SL = 9 (km) A1 N2  
[2 marks]
- (b) evidence of choosing sine rule (M1)  
 eg  $\frac{\sin A}{a} = \frac{\sin B}{b}, \sin \theta = \frac{(SL) \sin 20^\circ}{5}$   
 correct substitution (A1)  
 eg  $\frac{\sin \theta}{9} = \frac{\sin 20^\circ}{5}$   
 37.9981  
 $\hat{S}PL = 38.0^\circ$  A1 N2  
 recognition that second angle is the supplement of first (M1)  
 eg  $180 - x$   
 142.001  
 $\hat{S}QL = 142^\circ$  A1 N2  
[5 marks]

continued...



Question 8 continued

(c) (i)	new store is at Q	<b>A1</b>	<b>N1</b>
(ii)	<b>METHOD 1</b>		
	attempt to find third angle	<b>(M1)</b>	
	eg $\hat{S}LP = 180 - 20 - 38$ , $\hat{S}LQ = 180 - 20 - 142$		
	$\hat{S}LQ = 17.998^\circ$ (seen anywhere)	<b>A1</b>	
	evidence of choosing sine rule or cosine rule	<b>(M1)</b>	
	correct substitution into sine rule or cosine rule	<b>(A1)</b>	
	eg $\frac{x}{\sin 17.998} = \frac{5}{\sin 20} \left( = \frac{9}{\sin 142} \right)$ , $9^2 + 5^2 - 2(9)(5)\cos 17.998^\circ$		
	4.51708 km		
	<b>4.52</b> (km)	<b>A1</b>	<b>N3</b>
	<b>METHOD 2</b>		
	evidence of choosing cosine rule	<b>(M1)</b>	
	correct substitution into cosine rule	<b>A1</b>	
	eg $9^2 = x^2 + 5^2 - 2(x)(5)\cos 142^\circ$		
	attempt to solve	<b>(M1)</b>	
	eg sketch; setting quadratic equation equal to zero;		
	$0 = x^2 + 7.88x - 56$		
	one correct value for $x$	<b>(A1)</b>	
	eg $x = -12.3973$ , $x = 4.51708$		
	4.51708		
	<b>4.52</b> (km)	<b>A1</b>	<b>N3</b>
			<b>[6 marks]</b>

**Total [13 marks]**

9. (a) 0.0477903  
probability = **0.0478** **A2 N2**  
**[2 marks]**
- (b)  $P(\text{volume} < 250) = 0.02$  **(M1)**  
 $z = -2.05374$  (may be seen in equation) **A1**  
attempt to set up equation with  $z$  **(M1)**  
eg  $\frac{\mu - 260}{\sigma} = z, 260 - 2.05(\sigma) = 250$   
 $4.86914$   
 $\sigma = 4.87$  (ml) **A1 N3**  
**[4 marks]**
- (c) (i) 0.968062  
 $P(250 < \text{Vol} < 271) = 0.968$  **A2 N2**
- (ii) recognizing conditional probability (seen anywhere, including in correct working) **R1**  
eg  $P(A|B), \frac{P(A \cap B)}{P(B)}, P(A \cap B) = P(A|B)P(B)$   
correct value or expression for  $P(\text{not underfilled})$  **(A1)**  
eg  $0.98, 1 - 0.02, 1 - P(X < 250)$   
 $\text{probability} = \frac{0.968}{0.98}$  **A1**  
 $0.987818$   
probability = **0.988** **A1 N2**  
**[6 marks]**

continued...

Question 9 continued

(d) **METHOD 1**

evidence of recognizing binomial distribution (seen anywhere) (M1)

eg  $X \sim B(50, 0.968)$ , binomial cdf,  $p = 0.968$ ,  $r = 47$

$P(X \leq 47) = 0.214106$  (A1)

evidence of using complement (M1)

eg  $1 - P(X \leq 47)$

0.785894

probability = **0.786** A1 N3

**METHOD 2**

evidence of recognizing binomial distribution (seen anywhere) (M1)

eg  $X \sim B(50, 0.968)$ , binomial cdf,  $p = 0.968$ ,  $r = 47$

$P(\text{not pass}) = 1 - P(\text{pass}) = 0.0319378$  (A1)

evidence of attempt to find P (2 or fewer fail) (M1)

eg 0, 1, or 2 not pass,  $B(50, 2)$

0.785894

probability = **0.786** A1 N3

**METHOD 3**

evidence of recognizing binomial distribution (seen anywhere) (M1)

eg  $X \sim B(50, 0.968)$ , binomial cdf,  $p = 0.968$ ,  $r = 47$

evidence of summing probabilities (M1)

eg  $P(X = 48) + P(X = 49) + P(X = 50)$

correct working

eg  $0.263088 + 0.325488 + 0.197317$  (A1)

0.785894

probability = **0.786** A1 N3

[4 marks]

Total [16 marks]

10. (a)  $p = 6$  **A1** **N1**  
 recognising that turning points occur when  $f'(x) = 0$  **R1** **N1**  
 eg correct sign diagram  
 $f'$  changes from positive to negative at  $x = 6$  **R1** **N1**  
**[3 marks]**
- (b)  $f'(2) = -2$  **A1** **N1**  
**[1 mark]**
- (c) attempt to apply chain rule **(M1)**  
 eg  $\ln(x)' \times f'(x)$   
 correct expression for  $g'(x)$  **(A1)**  
 eg  $g'(x) = \frac{1}{f(x)} \times f'(x)$   
 substituting  $x = 2$  into **their**  $g'$  **(M1)**  
 eg  $\frac{f'(2)}{f(2)}$   
 $-0.666667$   
 $g'(2) = -\frac{2}{3}$  (**exact**),  $-0.667$  **A1** **N3**  
**[4 marks]**
- (d) evidence of integrating  $g'(x)$  **(M1)**  
 eg  $g(x)|_2^a$ ,  $g(x)|_a^2$   
 applying the fundamental theorem of calculus (seen anywhere) **R1**  
 eg  $\int_2^a g'(x) = g(a) - g(2)$   
 correct substitution into integral **(A1)**  
 eg  $\ln 3 + g(a) - g(2)$ ,  $\ln 3 + g(a) - \ln(f(2))$   
 $\ln 3 + g(a) - \ln 3$  **A1**  
 $\ln 3 + \int_2^a g'(x) = g(a)$  **AG** **N0**  
**[4 marks]**

continued...

Question 10 continued

(e) **METHOD 1**

substituting  $a = 5$  into the formula for  $g(a)$  (M1)

eg  $\int_2^5 g'(x) dx$ ,  $g(5) = \ln 3 + \int_2^5 g'(x) dx$  (do not accept only  $g(5)$ )

attempt to substitute areas (M1)

eg  $\ln 3 + 0.66 - 0.21$ ,  $\ln 3 + 0.66 + 0.21$

correct working

eg  $g(5) = \ln 3 + (-0.66 + 0.21)$  (A1)

0.648612

$g(5) = \ln 3 - 0.45$  (exact), **0.649** A1 N3

**METHOD 2**

attempt to set up an equation for one shaded region (M1)

eg  $\int_4^5 g'(x) dx = 0.21$ ,  $\int_2^4 g'(x) dx = -0.66$ ,  $\int_2^5 g'(x) dx = -0.45$

two correct equations

eg  $g(5) - g(4) = 0.21$ ,  $g(2) - g(4) = 0.66$  (A1)

combining equations to eliminate  $g(4)$

eg  $g(5) - [\ln 3 - 0.66] = 0.21$  (M1)

0.648612

$g(5) = \ln 3 - 0.45$  (exact), **0.649** A1 N3

**METHOD 3**

attempt to set up a definite integral (M1)

eg  $\int_2^5 g'(x) dx = -0.66 + 0.21$ ,  $\int_2^5 g'(x) dx = -0.45$

correct working

eg  $g(5) - g(2) = -0.45$  (A1)

correct substitution

eg  $g(5) - \ln 3 = -0.45$  (A1)

0.648612

$g(5) = \ln 3 - 0.45$  (exact), **0.649** A1 N3

[4 marks]

**Total [16 marks]**

**Mathematics**  
**Standard level**  
**Paper 2**

Wednesday 13 May 2015 (afternoon)

Candidate session number

1 hour 30 minutes

--	--	--	--	--	--	--	--	--	--

**Instructions to candidates**

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[90 marks]**.



Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

### Section A

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 6]

The following diagram shows triangle ABC.




diagram not to scale

- (a) Find AC. [3]
- (b) Find the area of triangle ABC. [3]

(This question continues on the following page)





(Question 1 continued)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



2. [Maximum mark: 7]

Let  $\mathbf{u} = 6\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}$  and  $\mathbf{v} = 2\mathbf{i} + 2\mathbf{j} + \mathbf{k}$ .

(a) Find

(i)  $\mathbf{u} \cdot \mathbf{v}$ ;

(ii)  $|\mathbf{u}|$ ;

(iii)  $|\mathbf{v}|$ .

[5]

(b) Find the angle between  $\mathbf{u}$  and  $\mathbf{v}$ .

[2]



3. [Maximum mark: 6]

The following table shows the sales,  $y$  millions of dollars, of a company,  $x$  years after it opened.

<b>Time after opening (<math>x</math> years)</b>	2	4	6	8	10
<b>Sales (<math>y</math> millions of dollars)</b>	12	20	30	36	52

The relationship between the variables is modelled by the regression line with equation  $y = ax + b$ .

(a) (i) Find the value of  $a$  and of  $b$ .

(ii) Write down the value of  $r$ .

[4]

(b) Hence estimate the sales in millions of dollars after seven years.

[2]

A large rectangular area for writing answers, featuring horizontal dotted lines. A large, faint watermark logo is centered in the background, consisting of a circular emblem with the letters 'SP' in the middle and the text 'SAT PAPER' and 'AHMEDABAD' around the perimeter.



4. [Maximum mark: 5]

The third term in the expansion of  $(x+k)^8$  is  $63x^6$ . Find the possible values of  $k$ .


.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



5. [Maximum mark: 6]

Let  $f(x) = e^{x+1} + 2$ , for  $-4 \leq x \leq 1$ .

(a) On the following grid, sketch the graph of  $f$ . [3]



(b) The graph of  $f$  is translated by the vector  $\begin{pmatrix} 3 \\ -1 \end{pmatrix}$  to obtain the graph of a function  $g$ .

Find an expression for  $g(x)$ . [3]

.....  
.....  
.....  
.....  
.....  
.....  
.....



6. [Maximum mark: 7]

Ramiro walks to work each morning. During the first minute he walks 80 metres. In each subsequent minute he walks 90% of the distance walked during the previous minute. The distance between his house and work is 660 metres. Ramiro leaves his house at 08:00 and has to be at work by 08:15.

Explain why he will not be at work on time.

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



7. [Maximum mark: 8]

Let  $f(x) = kx^2 + kx$  and  $g(x) = x - 0.8$ . The graphs of  $f$  and  $g$  intersect at two distinct points. Find the possible values of  $k$ .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....




Do **not** write solutions on this page.

### Section B

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 13]

Let  $f(x) = \frac{9}{x+2}$  and  $g(x) = 3x^2$ , for  $x \geq 0$ . Parts of the graphs of  $f$  and  $g$  are shown in the following diagram.



The graphs of  $f$  and  $g$  intersect at the point  $P(p, q)$ .

(a) Find the value of  $p$  and of  $q$ . [3]

(b) Write down  $f'(p)$ . [2]

Let  $L$  be the normal to the graph of  $f$  at  $P$ .

(c) (i) Find the equation of  $L$ , giving your answer in the form  $y = ax + b$ . [5]

(ii) Write down the  $y$ -intercept of  $L$ . [5]

(d) Let  $R$  be the region enclosed by the  $y$ -axis, the graph of  $g$  and the line  $L$ . Find the area of  $R$ . [3]





Do **not** write solutions on this page.


9. [Maximum mark: 16]

A machine manufactures a large number of nails. The length,  $L$  mm, of a nail is normally distributed, where  $L \sim N(50, \sigma^2)$ .

- (a) Find  $P(50 - \sigma < L < 50 + 2\sigma)$ . [3]
- (b) The probability that the length of a nail is less than 53.92 mm is 0.975. Show that  $\sigma = 2.00$  (correct to three significant figures). [2]

All nails with length at least  $t$  mm are classified as large nails.


- (c) A nail is chosen at random. The probability that it is a large nail is 0.75. Find the value of  $t$ . [3]
- (d) (i) A nail is chosen at random from the large nails. Find the probability that the length of this nail is less than 50.1 mm.
- (ii) Ten nails are chosen at random from the large nails. Find the probability that at least two nails have a length that is less than 50.1 mm. [8]



Do **not** write solutions on this page.

10. [Maximum mark: 16]

The following diagram shows a square ABCD, and a sector OAB of a circle centre O, radius  $r$ . Part of the square is shaded and labelled  $R$ .



$\hat{A}OB = \theta$ , where  $0.5 \leq \theta < \pi$ .

- (a) Show that the area of the square ABCD is  $2r^2(1 - \cos\theta)$ . [4]
- (b) When  $\theta = \alpha$ , the area of the square ABCD is equal to the area of the sector OAB .
  - (i) Write down the area of the sector when  $\theta = \alpha$  .
  - (ii) Hence find  $\alpha$  . [4]
- (c) When  $\theta = \beta$ , the area of  $R$  is more than twice the area of the sector. Find all possible values of  $\beta$  . [8]



**Mathematics**  
**Standard level**  
**Paper 2**

Thursday 12 November 2015 (afternoon)

Candidate session number

1 hour 30 minutes

--	--	--	--	--	--	--	--	--	--

**Instructions to candidates**

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[90 marks]**.



Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

### Section A

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 6]

The following diagram shows a circle with centre  $O$  and radius 3 cm.




diagram not to scale

Points A, B, and C lie on the circle, and  $\widehat{AOC} = 1.3$  radians.

- (a) Find the length of arc ABC. [2]
- (b) Find the area of the shaded region. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



2. [Maximum mark: 5]

The following table shows the probability distribution of a discrete random variable  $X$ .

$x$	0	1	2	3
$P(X = x)$	0.15	$k$	0.1	$2k$

(a) Find the value of  $k$ . [3]

(b) Find  $E(X)$ . [2]

.....

.....

.....

.....

.....

.....


.....

.....

.....


.....

.....



3. [Maximum mark: 7]

Let  $f(x) = 2 \ln(x - 3)$ , for  $x > 3$ . The following diagram shows part of the graph of  $f$ .



- (a) Find the equation of the vertical asymptote to the graph of  $f$ . [2]
- (b) Find the  $x$ -intercept of the graph of  $f$ . [2]
- (c) The region enclosed by the graph of  $f$ , the  $x$ -axis and the line  $x = 10$  is rotated  $360^\circ$  about the  $x$ -axis. Find the volume of the solid formed. [3]

(This question continues on the following page)



(Question 3 continued)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



12EP05

Turn over

4. [Maximum mark: 7]

The first three terms of a geometric sequence are  $u_1 = 0.64$ ,  $u_2 = 1.6$ , and  $u_3 = 4$ .

(a) Find the value of  $r$ . [2]

(b) Find the value of  $S_6$ . [2]

(c) Find the least value of  $n$  such that  $S_n > 75\,000$ . [3]





5. [Maximum mark: 7]

Let  $C$  and  $D$  be independent events, with  $P(C) = 2k$  and  $P(D) = 3k^2$ , where  $0 < k < 0.5$ .

(a) Write down an expression for  $P(C \cap D)$  in terms of  $k$ . [2]

(b) Given that  $P(C \cap D) = 0.162$ , find  $k$ . [2]

(c) Find  $P(C' | D)$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



12EP07


Turn over

6. [Maximum mark: 6]

The velocity  $v \text{ ms}^{-1}$  of a particle after  $t$  seconds is given by

$$v(t) = (0.3t + 0.1)^t - 4, \text{ for } 0 \leq t \leq 5.$$

The following diagram shows the graph of  $v$ .



- (a) Find the value of  $t$  when the particle is at rest. [3]
- (b) Find the value of  $t$  when the acceleration of the particle is 0. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



7. [Maximum mark: 8]

Let  $f(x) = \ln(x^2)$ , for  $x \neq 0$ .

(a) Show that  $f'(x) = \frac{2}{x}$ . [2]

(b) The tangent to the graph of  $f$  at a point  $P(d, f(d))$  passes through another point  $Q(1, -3)$ . Find the value of  $d$ . [6]

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....




Do **not** write solutions on this page.

**Section B**

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 14]

The following diagram shows the quadrilateral ABCD.



$AD = 6 \text{ cm}$ ,  $AB = 15 \text{ cm}$ ,  $\hat{A}BC = 44^\circ$ ,  $\hat{A}CB = 83^\circ$  and  $\hat{D}AC = \theta$

(a) Find AC. [3]

(b) Find the area of triangle ABC. [3]

The area of triangle ACD is half the area of triangle ABC.

(c) Find the possible values of  $\theta$ . [5]

(d) Given that  $\theta$  is obtuse, find CD. [3]



Do **not** write solutions on this page.

9. [Maximum mark: 16]

An environmental group records the numbers of coyotes and foxes in a wildlife reserve after  $t$  years, starting on 1 January 1995.

Let  $c$  be the number of coyotes in the reserve after  $t$  years. The following table shows the number of coyotes after  $t$  years.

number of years ( $t$ )	0	2	10	15	19
number of coyotes ( $c$ )	115	197	265	320	406

The relationship between the variables can be modelled by the regression equation  $c = at + b$ .

- (a) Find the value of  $a$  and of  $b$ . [3]
- (b) Use the regression equation to estimate the number of coyotes in the reserve when  $t = 7$ . [3]

Let  $f$  be the number of foxes in the reserve after  $t$  years. The number of foxes can be modelled by the equation  $f = \frac{2000}{1 + 99e^{-kt}}$ , where  $k$  is a constant.

- (c) Find the number of foxes in the reserve on 1 January 1995. [3]
- (d) After five years, there were 64 foxes in the reserve. Find  $k$ . [3]
- (e) During which year were the number of coyotes the same as the number of foxes? [4]



Do **not** write solutions on this page.

10. [Maximum mark: 14]

The masses of watermelons grown on a farm are normally distributed with a mean of 10 kg. The watermelons are classified as small, medium or large.

A watermelon is small if its mass is less than 4 kg. Five percent of the watermelons are classified as small.

(a) Find the standard deviation of the masses of the watermelons. [4]

The following table shows the percentages of small, medium and large watermelons grown on the farm.

small	medium	large
5%	57%	38%

A watermelon is large if its mass is greater than  $w$  kg.

(b) Find the value of  $w$ . [2]

All the medium and large watermelons are delivered to a grocer.

(c) The grocer selects a watermelon at random from **this** delivery. Find the probability that it is medium. [3]

(d) The grocer sells all the medium watermelons for \$1.75 each, and all the large watermelons for \$3.00 each. His costs on this delivery are \$300, and his total profit is \$150. Find the number of watermelons in the delivery. [5]



**Mathematics**  
**Standard level**  
**Paper 2**

Wednesday 11 May 2016 (morning)

Candidate session number

1 hour 30 minutes

--	--	--	--	--	--	--	--	--	--

**Instructions to candidates**

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[90 marks]**.





Please **do not** write on this page.  
Answers written on this page will not  
be marked.





Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.


### Section A

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 6]

A random variable  $X$  is distributed normally with a mean of 20 and standard deviation of 4.

(a) On the following diagram, shade the region representing  $P(X \leq 25)$ . [2]



(b) Write down  $P(X \leq 25)$ , correct to two decimal places. [2]

(c) Let  $P(X \leq c) = 0.7$ . Write down the value of  $c$ . [2]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



2. [Maximum mark: 6]

Let  $f(x) = x^2$  and  $g(x) = 3 \ln(x + 1)$ , for  $x > -1$ .

(a) Solve  $f(x) = g(x)$ . [3]


(b) Find the area of the region enclosed by the graphs of  $f$  and  $g$ . [3]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



3. [Maximum mark: 7]

The following diagram shows three towns A, B and C. Town B is 5 km from Town A, on a bearing of  $070^\circ$ . Town C is 8 km from Town B, on a bearing of  $115^\circ$ .



- (a) Find  $\hat{A}BC$ . [2]
- (b) Find the distance from Town A to Town C. [3]
- (c) Use the sine rule to find  $\hat{A}CB$ . [2]

SAHIBGAT PREP  
 SP  
 AHMEDABAD

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Turn over

4. [Maximum mark: 6]

(a) Find the term in  $x^6$  in the expansion of  $(x + 2)^9$ . [4]

(b) Hence, find the term in  $x^7$  in the expansion of  $5x(x + 2)^9$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....



5. [Maximum mark: 6]

The mass  $M$  of a decaying substance is measured at one minute intervals. The points  $(t, \ln M)$  are plotted for  $0 \leq t \leq 10$ , where  $t$  is in minutes. The line of best fit is drawn. This is shown in the following diagram.



The correlation coefficient for this linear model is  $r = -0.998$ .

- (a) State **two** words that describe the linear correlation between  $\ln M$  and  $t$ . [2]
- (b) The equation of the line of best fit is  $\ln M = -0.12t + 4.67$ . Given that  $M = a \times b^t$ , find the value of  $b$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



6. [Maximum mark: 6]

In a geometric sequence, the fourth term is 8 times the first term. The sum of the first 10 terms is 2557.5. Find the 10th term of this sequence.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



7. [Maximum mark: 8]

**Note: One decade is 10 years**

A population of rare birds,  $P_t$ , can be modelled by the equation  $P_t = P_0 e^{kt}$ , where  $P_0$  is the initial population, and  $t$  is measured in decades. After one decade, it is estimated that  $\frac{P_1}{P_0} = 0.9$ .

- (a) (i) Find the value of  $k$ .  
  
(ii) Interpret the meaning of the value of  $k$ . [3]
  
- (b) Find the least number of **whole** years for which  $\frac{P_t}{P_0} < 0.75$ . [5]



Do **not** write solutions on this page.

### Section B

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 15]

A factory has two machines, A and B. The number of breakdowns of each machine is independent from day to day.

Let  $A$  be the number of breakdowns of Machine A on any given day. The probability distribution for  $A$  can be modelled by the following table.

$a$	0	1	2	3
$P(A = a)$	0.55	0.3	0.1	$k$

- (a) Find  $k$ . [2]
- (b) (i) A day is chosen at random. Write down the probability that Machine A has no breakdowns.
- (ii) Five days are chosen at random. Find the probability that Machine A has no breakdowns on exactly four of these days. [3]

Let  $B$  be the number of breakdowns of Machine B on any given day. The probability distribution for  $B$  can be modelled by the following table.

$b$	0	1	2	3
$P(B = b)$	0.7	0.2	0.08	0.02

- (c) Find  $E(B)$ . [2]

On Tuesday, the factory uses both Machine A and Machine B. The variables  $A$  and  $B$  are independent.

- (d) (i) Find the probability that there are exactly two breakdowns on Tuesday.
- (ii) Given that there are exactly two breakdowns on Tuesday, find the probability that both breakdowns are of Machine A. [8]






Do **not** write solutions on this page.

9. [Maximum mark: 14]

A particle P moves along a straight line so that its velocity,  $v \text{ ms}^{-1}$ , after  $t$  seconds, is given by  $v = \cos 3t - 2 \sin t - 0.5$ , for  $0 \leq t \leq 5$ . The initial displacement of P from a fixed point O is 4 metres.

(a) Find the displacement of P from O after 5 seconds. [5]

The following sketch shows the graph of  $v$ .



(b) Find when P is first at rest. [2]

(c) Write down the number of times P changes direction. [2]

(d) Find the acceleration of P after 3 seconds. [2]

(e) Find the maximum speed of P. [3]



Do **not** write solutions on this page.

10. [Maximum mark: 16]

The points A and B lie on a line  $L$ , and have position vectors  $\begin{pmatrix} -3 \\ -2 \\ 2 \end{pmatrix}$  and  $\begin{pmatrix} 6 \\ 4 \\ -1 \end{pmatrix}$  respectively.

Let O be the origin. This is shown on the following diagram.

**diagram not to scale**



(a) Find  $\vec{AB}$ . [2]

The point C also lies on  $L$ , such that  $\vec{AC} = 2\vec{CB}$ .


(b) Show that  $\vec{OC} = \begin{pmatrix} 3 \\ 2 \\ 0 \end{pmatrix}$ . [3]

Let  $\theta$  be the angle between  $\vec{AB}$  and  $\vec{OC}$ .

(c) Find  $\theta$ . [5]

Let D be a point such that  $\vec{OD} = k\vec{OC}$ , where  $k > 1$ . Let E be a point on  $L$  such that  $\hat{C}ED$  is a right angle. This is shown on the following diagram.

**diagram not to scale**



(d) (i) Show that  $|\vec{DE}| = (k-1)|\vec{OC}| \sin \theta$ .

(ii) The distance from D to line  $L$  is less than 3 units. Find the possible values of  $k$ . [6]



**Mathematics**  
**Standard level**  
**Paper 2**

Wednesday 11 May 2016 (morning)

Candidate session number

1 hour 30 minutes

--	--	--	--	--	--	--	--	--	--

**Instructions to candidates**


- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[90 marks]**.





2. [Maximum mark: 6]

The following diagram shows a quadrilateral ABCD.



$AD = 7 \text{ cm}$ ,  $BC = 8 \text{ cm}$ ,  $CD = 12 \text{ cm}$ ,  $\hat{DAB} = 1.75 \text{ radians}$ ,  $\hat{ABD} = 0.82 \text{ radians}$ .

- (a) Find  $BD$ . [3]
- (b) Find  $\hat{DBC}$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



3. [Maximum mark: 7]

Let  $f(x) = e^{0.5x} - 2$ .

(a) For the graph of  $f$

- (i) write down the  $y$ -intercept;
- (ii) find the  $x$ -intercept;
- (iii) write down the equation of the horizontal asymptote.

[4]


(This question continues on the following page)



(Question 3 continued)

(b) On the following grid, sketch the graph of  $f$ , for  $-4 \leq x \leq 4$ .

[3]




4. [Maximum mark: 8]

The height,  $h$  metres, of a seat on a Ferris wheel after  $t$  minutes is given by

$$h(t) = -15 \cos 1.2t + 17, \text{ for } t \geq 0.$$

- (a) Find the height of the seat when  $t = 0$ . [2]
- (b) The seat first reaches a height of 20 m after  $k$  minutes. Find  $k$ . [3]
- (c) Calculate the time needed for the seat to complete a full rotation, giving your answer correct to one decimal place. [3]






5. [Maximum mark: 6]

Consider the expansion of  $\left(x^2 + \frac{2}{x}\right)^{10}$ .

- (a) Write down the number of terms of this expansion. [1]
- (b) Find the coefficient of  $x^8$ . [5]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



6. [Maximum mark: 6]

A competition consists of two independent events, shooting at 100 targets and running for one hour.


The number of targets a contestant hits is the  $S$  score. The  $S$  scores are normally distributed with mean 65 and standard deviation 10.

(a) A contestant is chosen at random. Find the probability that their  $S$  score is less than 50. [2]

The distance in km that a contestant runs in one hour is the  $R$  score. The  $R$  scores are normally distributed with mean 12 and standard deviation 2.5. The  $R$  score is independent of the  $S$  score.

Contestants are disqualified if their  $S$  score is less than 50 **and** their  $R$  score is less than  $x$  km.

(b) Given that 1% of the contestants are disqualified, find the value of  $x$ . [4]



7. [Maximum mark: 7]

A particle moves in a straight line. Its velocity  $v \text{ m s}^{-1}$  after  $t$  seconds is given by

$$v = 6t - 6, \text{ for } 0 \leq t \leq 2.$$

After  $p$  seconds, the particle is 2 m from its initial position. Find the possible values of  $p$ .

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



12EP09

Turn over

Do **not** write solutions on this page.

### Section B

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 15]

The price of a used car depends partly on the distance it has travelled. The following table shows the distance and the price for seven cars on 1 January 2010.

<b>Distance, <math>x</math> km</b>	11 500	7500	13 600	10 800	9500	12 200	10 400
<b>Price, <math>y</math> dollars</b>	15 000	21 500	12 000	16 000	19 000	14 500	17 000

The relationship between  $x$  and  $y$  can be modelled by the regression equation  $y = ax + b$ .

- (a) (i) Find the correlation coefficient.
- (ii) Write down the value of  $a$  and of  $b$ . [4]

On 1 January 2010, Lina buys a car which has travelled 11 000 km.

- (b) Use the regression equation to estimate the price of Lina's car, giving your answer to the nearest 100 dollars. [3]

The price of a car decreases by 5% each year.

- (c) Calculate the price of Lina's car after 6 years. [4]

Lina will sell her car when its price reaches 10 000 dollars.

- (d) Find the year when Lina sells her car. [4]



Do **not** write solutions on this page.

9. [Maximum mark: 14]

Let  $f(x) = \frac{1}{x-1} + 2$ , for  $x > 1$ .

(a) Write down the equation of the horizontal asymptote of the graph of  $f$ . [2]


(b) Find  $f'(x)$ . [2]

Let  $g(x) = ae^{-x} + b$ , for  $x \geq 1$ . The graphs of  $f$  and  $g$  have the same horizontal asymptote.

(c) Write down the value of  $b$ . [2]

(d) Given that  $g'(1) = -e$ , find the value of  $a$ . [4]

(e) There is a value of  $x$ , for  $1 < x < 4$ , for which the graphs of  $f$  and  $g$  have the same gradient. Find this gradient. [4]



Do **not** write solutions on this page.

10. [Maximum mark: 15]

Consider the points  $A(1, 5, -7)$  and  $B(-9, 9, -6)$ .

(a) Find  $\vec{AB}$ . [2]

Let  $C$  be a point such that  $\vec{AC} = \begin{pmatrix} 6 \\ -4 \\ 0 \end{pmatrix}$ .


(b) Find the coordinates of  $C$ . [2]

The line  $L$  passes through  $B$  and is parallel to  $(AC)$ .

(c) Write down a vector equation for  $L$ . [2]

(d) Given that  $|\vec{AB}| = k|\vec{AC}|$ , find  $k$ . [3]

(e) The point  $D$  lies on  $L$  such that  $|\vec{AB}| = |\vec{BD}|$ . Find the possible coordinates of  $D$ . [6]



**Mathematics**  
**Standard level**  
**Paper 2**

Friday 11 November 2016 (morning)

Candidate session number

1 hour 30 minutes

--	--	--	--	--	--	--	--	--	--

**Instructions to candidates**

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[90 marks]**.





Please **do not** write on this page.  
Answers written on this page will not  
be marked.





Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example, if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

### Section A

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 7]

Let  $f(x) = x^2 + 2x + 1$  and  $g(x) = x - 5$ , for  $x \in \mathbb{R}$ .

- (a) Find  $f(8)$ . [2]
- (b) Find  $(g \circ f)(x)$ . [2]
- (c) Solve  $(g \circ f)(x) = 0$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



2. [Maximum mark: 7]


Let  $f(x) = 0.225x^3 - 2.7x$ , for  $-3 \leq x \leq 3$ . There is a local minimum point at A.

(a) Find the coordinates of A. [2]

(b) On the following grid,

(i) sketch the graph of  $f$ , clearly indicating the point A;

(ii) sketch the tangent to the graph of  $f$  at A. [5]



.....

.....

.....

.....

.....

.....

.....

.....

.....


.....



3. [Maximum mark: 7]

The following diagram shows a circle, centre  $O$  and radius  $r$  mm. The circle is divided into five equal sectors.

diagram not to scale



One sector is  $OAB$ , and  $\widehat{AOB} = \theta$ .

(a) Write down the **exact** value of  $\theta$  in radians. [1]

The area of sector  $AOB$  is  $20\pi$  mm<sup>2</sup>.

(b) Find the value of  $r$ . [3]

(c) Find  $AB$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



4. [Maximum mark: 6]

Let  $f(x) = xe^{-x}$  and  $g(x) = -3f(x) + 1$ .

The graphs of  $f$  and  $g$  intersect at  $x = p$  and  $x = q$ , where  $p < q$ .

- (a) Find the value of  $p$  and of  $q$ . [3]
- (b) Hence, find the area of the region enclosed by the graphs of  $f$  and  $g$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



5. [Maximum mark: 6]

The weights,  $W$ , of newborn babies in Australia are normally distributed with a mean 3.41 kg and standard deviation 0.57 kg. A newborn baby has a low birth weight if it weighs less than  $w$  kg.

(a) Given that 5.3% of newborn babies have a low birth weight, find  $w$ . [3]

(b) A newborn baby has a low birth weight.  
Find the probability that the baby weighs at least 2.15 kg. [3]


A large rectangular area containing horizontal dotted lines for writing answers. A large, faint watermark logo is centered in this area. The logo is circular with 'SAT PREP' at the top and 'AHMEDABAD' at the bottom. In the center of the logo are the letters 'SP' in a stylized font, with a green leaf-like shape behind the 'P'.



6. [Maximum mark: 6]

All lengths in this question are in metres.

Let  $f(x) = -0.8x^2 + 0.5$ , for  $-0.5 \leq x \leq 0.5$ . Mark uses  $f(x)$  as a model to create a barrel. The region enclosed by the graph of  $f$ , the  $x$ -axis, the line  $x = -0.5$  and the line  $x = 0.5$  is rotated  $360^\circ$  about the  $x$ -axis. This is shown in the following diagram.



- (a) Use the model to find the volume of the barrel. [3]
- (b) The empty barrel is being filled with water. The volume  $V \text{ m}^3$  of water in the barrel after  $t$  minutes is given by  $V = 0.8(1 - e^{-0.1t})$ . How long will it take for the barrel to be half-full? [3]

Area for student response with horizontal dotted lines.



7. [Maximum mark: 6]

A jar contains 5 red discs, 10 blue discs and  $m$  green discs. A disc is selected at random and replaced. This process is performed four times.

(a) Write down the probability that the first disc selected is red. [1]

(b) Let  $X$  be the number of red discs selected. Find the smallest value of  $m$  for which  $\text{Var}(X) < 0.6$ . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



16EP09

Do **not** write solutions on this page.

### Section B

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 16]

Ten students were surveyed about the number of hours,  $x$ , they spent browsing the Internet during week 1 of the school year. The results of the survey are given below.

$$\sum_{i=1}^{10} x_i = 252, \sigma = 5 \text{ and median} = 27.$$

- (a) Find the mean number of hours spent browsing the Internet. [2]
- (b) During week 2, the students worked on a major project and they each spent an additional five hours browsing the Internet. For week 2, write down
- (i) the mean; [2]
- (ii) the standard deviation.
- (c) During week 3 each student spent 5% less time browsing the Internet than during week 1. For week 3, find
- (i) the median; [6]
- (ii) the variance.
- (d) During week 4, the survey was extended to all 200 students in the school. The results are shown in the cumulative frequency graph on the following page.
- (i) Find the number of students who spent between 25 and 30 hours browsing the Internet.
- (ii) Given that 10% of the students spent more than  $k$  hours browsing the Internet, find the maximum value of  $k$ . [6]


(This question continues on the following page)





Do **not** write solutions on this page.

**(Question 8 continued)**



16EP11

Turn over


Do **not** write solutions on this page.

9. [Maximum mark: 14]

A particle P starts from a point A and moves along a horizontal straight line. Its velocity  $v \text{ cm s}^{-1}$  after  $t$  seconds is given by

$$v(t) = \begin{cases} -2t + 2, & \text{for } 0 \leq t \leq 1 \\ 3\sqrt{t} + \frac{4}{t^2} - 7, & \text{for } 1 \leq t \leq 12 \end{cases}$$

The following diagram shows the graph of  $v$ .



(a) Find the initial velocity of P. [2]

P is at rest when  $t = 1$  and  $t = p$ .

(b) Find the value of  $p$ . [2]

When  $t = q$ , the acceleration of P is zero.

(c) (i) Find the value of  $q$ . [4]

(ii) Hence, find the **speed** of P when  $t = q$ . [4]

(d) (i) Find the total distance travelled by P between  $t = 1$  and  $t = p$ .


(ii) Hence or otherwise, find the displacement of P from A when  $t = p$ . [6]



Do **not** write solutions on this page.

10. [Maximum mark: 15]

The following diagram shows the graph of  $f(x) = a \sin bx + c$ , for  $0 \leq x \leq 12$ .



The graph of  $f$  has a minimum point at  $(3, 5)$  and a maximum point at  $(9, 17)$ .

- (a) (i) Find the value of  $c$ .
- (ii) Show that  $b = \frac{\pi}{6}$ .
- (iii) Find the value of  $a$ . [6]

The graph of  $g$  is obtained from the graph of  $f$  by a translation of  $\begin{pmatrix} k \\ 0 \end{pmatrix}$ . The maximum point on the graph of  $g$  has coordinates  $(11.5, 17)$ .

- (b) (i) Write down the value of  $k$ .
- (ii) Find  $g(x)$ . [3]

The graph of  $g$  changes from concave-up to concave-down when  $x = w$ .

- (c) (i) Find  $w$ .
- (ii) Hence or otherwise, find the maximum positive rate of change of  $g$ . [6]





Please **do not** write on this page.  
Answers written on this page will not  
be marked.





Please **do not** write on this page.  
Answers written on this page will not  
be marked.





Please **do not** write on this page.  
Answers written on this page will not  
be marked.



**Mathematics**  
**Standard level**  
**Paper 2**

Friday 5 May 2017 (morning)

Candidate session number

1 hour 30 minutes

--	--	--	--	--	--	--	--	--	--

**Instructions to candidates**

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions. Answers must be written within the answer boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[90 marks]**.



Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

### Section A

Answer **all** questions. Answers must be written within the answer boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 7]

Consider the following frequency table.

$x$	Frequency
2	8
4	15
7	21
10	28
11	3

- (a) (i) Write down the mode.
- (ii) Find the value of the range. [3]
- (b) (i) Find the mean.
- (ii) Find the variance. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





2. [Maximum mark: 6]

Let  $\mathbf{v} = \begin{pmatrix} -10 \\ 2 \\ 1 \end{pmatrix}$  and  $\mathbf{w} = \begin{pmatrix} 3 \\ -4 \\ 0 \end{pmatrix}$ . Find the angle between  $\mathbf{v}$  and  $\mathbf{w}$ , giving your answer correct to one decimal place.

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



Turn over

3. [Maximum mark: 6]

Consider the graph of  $f(x) = \frac{e^x}{5x-10} + 3$ , for  $x \neq 2$ .

- (a) Find the  $y$ -intercept. [2]
- (b) Find the equation of the vertical asymptote. [2]
- (c) Find the minimum value of  $f(x)$  for  $x > 2$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



4. [Maximum mark: 6]

In a large university the probability that a student is left handed is 0.08. A sample of 150 students is randomly selected from the university. Let  $k$  be the expected number of left-handed students in this sample.

(a) Find  $k$ . [2]

(b) Hence, find the probability that

(i) exactly  $k$  students are left handed;


(ii) fewer than  $k$  students are left handed. [4]



5. [Maximum mark: 7]

The following diagram shows the chord  $[AB]$  in a circle of radius 8 cm, where  $AB = 12$  cm.

diagram not to scale



Find the area of the shaded segment.

Handwriting practice area with horizontal dotted lines. A large watermark for 'SP AHMEDABAD' is visible in the background.



6. [Maximum mark: 7]

Let  $f(x) = (x^2 + 3)^7$ . Find the term in  $x^5$  in the expansion of the derivative,  $f'(x)$ .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....



12EP07

7. [Maximum mark: 7]

A particle P moves along a straight line. Its velocity  $v_p \text{ m s}^{-1}$  after  $t$  seconds is given by  $v_p = \sqrt{t} \sin\left(\frac{\pi}{2}t\right)$ , for  $0 \leq t \leq 8$ . The following diagram shows the graph of  $v_p$ .



- (a) (i) Write down the first value of  $t$  at which P changes direction.
- (ii) Find the **total** distance travelled by P, for  $0 \leq t \leq 8$ . [3]
- (b) A second particle Q also moves along a straight line. Its velocity,  $v_Q \text{ m s}^{-1}$  after  $t$  seconds is given by  $v_Q = \sqrt{t}$  for  $0 \leq t \leq 8$ . After  $k$  seconds Q has travelled the same total distance as P.
- Find  $k$ . [4]

(This question continues on the following page)



(Question 7 continued)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....




Do **not** write solutions on this page.

### Section B

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 14]

At Grande Anse Beach the height of the water in metres is modelled by the function  $h(t) = p \cos(q \times t) + r$ , where  $t$  is the number of hours after 21:00 hours on 10 December 2017. The following diagram shows the graph of  $h$ , for  $0 \leq t \leq 72$ .



The point  $A(6.25, 0.6)$  represents the first low tide and  $B(12.5, 1.5)$  represents the next high tide.


- (a) (i) How much time is there between the first low tide and the next high tide?
- (ii) Find the difference in height between low tide and high tide. [4]
- (b) Find the value of
  - (i)  $p$ ;
  - (ii)  $q$ ;
  - (iii)  $r$ . [7]
- (c) There are two high tides on 12 December 2017. At what time does the second high tide occur? [3]





9. [Maximum mark: 15]

A random variable  $X$  is normally distributed with mean,  $\mu$ . In the following diagram, the shaded region between 9 and  $\mu$  represents 30% of the distribution.



(a) Find  $P(X < 9)$ . [2]


The standard deviation of  $X$  is 2.1.

(b) Find the value of  $\mu$ . [3]

The random variable  $Y$  is normally distributed with mean  $\lambda$  and standard deviation 3.5. The events  $X > 9$  and  $Y > 9$  are independent, and  $P((X > 9) \cap (Y > 9)) = 0.4$ .

(c) Find  $\lambda$ . [5]

(d) Given that  $Y > 9$ , find  $P(Y < 13)$ . [5]



10. [Maximum mark: 15]

Let  $f(x) = \ln x$  and  $g(x) = 3 + \ln\left(\frac{x}{2}\right)$ , for  $x > 0$ .

The graph of  $g$  can be obtained from the graph of  $f$  by two transformations:

a horizontal stretch of scale factor  $q$  followed by  
a translation of  $\begin{pmatrix} h \\ k \end{pmatrix}$ .

(a) Write down the value of


(i)  $q$ ;

(ii)  $h$ ;

(iii)  $k$ .

[3]

Let  $h(x) = g(x) \times \cos(0.1x)$ , for  $0 < x < 4$ . The following diagram shows the graph of  $h$  and the line  $y = x$ .



The graph of  $h$  intersects the graph of  $h^{-1}$  at two points. These points have  $x$  coordinates 0.111 and 3.31, correct to three significant figures.

(b) (i) Find  $\int_{0.111}^{3.31} (h(x) - x) dx$ .

(ii) Hence, find the area of the region enclosed by the graphs of  $h$  and  $h^{-1}$ .

[5]

(c) Let  $d$  be the vertical distance from a point on the graph of  $h$  to the line  $y = x$ . There is a point  $P(a, b)$  on the graph of  $h$  where  $d$  is a maximum. Find the coordinates of  $P$ , where  $0.111 < a < 3.31$ .

[7]



**Mathematics**  
**Standard level**  
**Paper 2**

Friday 5 May 2017 (morning)

Candidate session number

1 hour 30 minutes

--	--	--	--	--	--	--	--	--	--

**Instructions to candidates**

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions. Answers must be written within the answer boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[90 marks]**.



Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.


### Section A

Answer **all** questions. Answers must be written within the answer boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 6]

The following diagram shows a circle with centre  $O$  and radius 40 cm.

diagram not to scale



The points  $A$ ,  $B$  and  $C$  are on the circumference of the circle and  $\hat{AOC} = 1.9$  radians.

- (a) Find the length of arc  $ABC$ . [2]
- (b) Find the perimeter of sector  $OABC$ . [2]
- (c) Find the area of sector  $OABC$ . [2]

(This question continues on the following page)



(Question 1 continued)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



2. [Maximum mark: 7]

The maximum temperature  $T$ , in degrees Celsius, in a park on six randomly selected days is shown in the following table. The table also shows the number of visitors,  $N$ , to the park on each of those six days.

Maximum temperature ( $T$ )	4	5	17	31	29	11
Number of visitors ( $N$ )	24	26	36	38	46	28

The relationship between the variables can be modelled by the regression equation  $N = aT + b$ .

- (a) (i) Find the value of  $a$  and of  $b$ .
- (ii) Write down the value of  $r$ . [4]
- (b) Use the regression equation to estimate the number of visitors on a day when the maximum temperature is  $15^\circ\text{C}$ . [3]

ST PREP

SP

HYDRABAD

---



---



---



---



---



---



---



---



---



---



---



---



---




---



3. [Maximum mark: 6]

The following diagram shows the graph of a function  $y = f(x)$ , for  $-6 \leq x \leq -2$ .  
The points  $(-6, 6)$  and  $(-2, 6)$  lie on the graph of  $f$ . There is a minimum point at  $(-4, 0)$ .

(a) Write down the range of  $f$ . [2]



Let  $g(x) = f(x - 5)$ .

(b) On the grid above, sketch the graph of  $g$ . [2]

(c) Write down the domain of  $g$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



4. [Maximum mark: 6]

The depth of water in a port is modelled by the function  $d(t) = p \cos qt + 7.5$ , for  $0 \leq t \leq 12$ , where  $t$  is the number of hours after high tide.

At high tide, the depth is 9.7 metres.

At low tide, which is 7 hours later, the depth is 5.3 metres.

- (a) Find the value of  $p$ . [2]
- (b) Find the value of  $q$ . [2]
- (c) Use the model to find the depth of the water 10 hours after high tide. [2]

A large rectangular area with horizontal dotted lines for writing. In the center, there is a faint circular watermark logo. The logo contains the text 'SAT PREP' at the top and 'AHMEDABAD' at the bottom. In the center of the logo are the letters 'SP' in a stylized font, with a green leaf-like shape below the 'P'.






5. [Maximum mark: 6]

Consider a geometric sequence where the first term is 768 and the second term is 576.

Find the least value of  $n$  such that the  $n$ th term of the sequence is less than 7.




6. [Maximum mark: 8]

Let  $f(x) = x^2 - 1$  and  $g(x) = x^2 - 2$ , for  $x \in \mathbb{R}$ .

(a) Show that  $(f \circ g)(x) = x^4 - 4x^2 + 3$ . [2]

(b) On the following grid, sketch the graph of  $(f \circ g)(x)$ , for  $0 \leq x \leq 2.25$ . [3]



(c) The equation  $(f \circ g)(x) = k$  has exactly two solutions, for  $0 \leq x \leq 2.25$ . Find the possible values of  $k$ . [3]

(This question continues on the following page)



(Question 6 continued)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....



7. [Maximum mark: 6]

**Note: In this question, distance is in metres and time is in seconds.**

A particle moves along a horizontal line starting at a fixed point A. The velocity  $v$  of the particle, at time  $t$ , is given by  $v(t) = \frac{2t^2 - 4t}{t^2 - 2t + 2}$ , for  $0 \leq t \leq 5$ . The following diagram shows the graph of  $v$



There are  $t$ -intercepts at  $(0, 0)$  and  $(2, 0)$ .

Find the maximum distance of the particle from A during the time  $0 \leq t \leq 5$  and justify your answer.

**(This question continues on the following page)**



(Question 7 continued)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



16EP11

Turn over


Do **not** write solutions on this page.

**Section B**

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 15]

Let  $f(x) = -0.5x^4 + 3x^2 + 2x$ . The following diagram shows part of the graph of  $f$ .



There are  $x$ -intercepts at  $x = 0$  and at  $x = p$ . There is a maximum at A where  $x = a$ , and a point of inflexion at B where  $x = b$ .

- (a) Find the value of  $p$ . [2]
- (b) (i) Write down the coordinates of A. [3]
- (ii) Write down the rate of change of  $f$  at A. [3]
- (c) (i) Find the coordinates of B. [7]
- (ii) Find the the rate of change of  $f$  at B. [7]
- (d) Let  $R$  be the region enclosed by the graph of  $f$ , the  $x$ -axis, the line  $x = b$  and the line  $x = a$ . The region  $R$  is rotated  $360^\circ$  about the  $x$ -axis. Find the volume of the solid formed. [3]




Do **not** write solutions on this page.

9. [Maximum mark: 15]

A ship is sailing north from a point A towards point D. Point C is 175 km north of A. Point D is 60 km north of C. There is an island at E. The bearing of E from A is  $055^\circ$ . The bearing of E from C is  $134^\circ$ . This is shown in the following diagram.

**diagram not to scale**



- (a) Find the bearing of A from E. [2]
- (b) Find CE. [5]
- (c) Find DE. [3]
- (d) When the ship reaches D, it changes direction and travels directly to the island at 50 km per hour. At the same time as the ship changes direction, a boat starts travelling to the island from a point B. This point B lies on (AC), between A and C, and is the closest point to the island. The ship and the boat arrive at the island at the same time. Find the speed of the boat. [5]



Do **not** write solutions on this page.

10. [Maximum mark: 15]

The following table shows a probability distribution for the random variable  $X$ , where  $E(X) = 1.2$ .

$x$	0	1	2	3
$P(X=x)$	$p$	$\frac{1}{2}$	$\frac{3}{10}$	$q$

(a) (i) Find  $q$ .

(ii) Find  $p$ .

[4]

A bag contains white and blue marbles, with at least three of each colour. Three marbles are drawn from the bag, without replacement. The number of blue marbles drawn is given by the random variable  $X$ .

(b) (i) Write down the probability of drawing three blue marbles.

(ii) Explain why the probability of drawing three white marbles is  $\frac{1}{6}$ .

(iii) The bag contains a total of ten marbles of which  $w$  are white. Find  $w$ .

[5]

A game is played in which three marbles are drawn from the bag of ten marbles, without replacement. A player wins a prize if three white marbles are drawn.

(c) Jill plays the game nine times. Find the probability that she wins exactly two prizes.

[2]

(d) Grant plays the game until he wins two prizes. Find the probability that he wins his second prize on his eighth attempt.

[4]







Please **do not** write on this page.

Answers written on this page  
will not be marked.



16EP15



Please **do not** write on this page.

Answers written on this page  
will not be marked.



16EP16

**Mathematics**  
**Standard level**  
**Paper 2**

Tuesday 14 November 2017 (morning)

Candidate session number

1 hour 30 minutes

--	--	--	--	--	--	--	--	--	--

**Instructions to candidates**

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions. Answers must be written within the answer boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[90 marks]**.





Please **do not** write on this page.

Answers written on this page  
will not be marked.



Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, for example if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.


### Section A

Answer **all** questions. Answers must be written within the answer boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 6]

The following diagram shows a triangle ABC.

diagram not to scale



$AB = 5 \text{ cm}$ ,  $\hat{C}AB = 50^\circ$  and  $\hat{A}CB = 112^\circ$

- (a) Find BC. [3]
- (b) Find the area of triangle ABC. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....




Turn over

2. [Maximum mark: 7]

Let  $f(x) = \frac{6x^2 - 4}{e^x}$ , for  $0 \leq x \leq 7$ .

- (a) Find the  $x$ -intercept of the graph of  $f$ . [2]
- (b) The graph of  $f$  has a maximum at the point A. Write down the coordinates of A. [2]
- (c) On the following grid, sketch the graph of  $f$ . [3]



.....

.....

.....

.....

.....

.....

.....

.....



3. [Maximum mark: 6]

Let  $\vec{AB} = \begin{pmatrix} 4 \\ 1 \\ 2 \end{pmatrix}$ .

(a) Find  $|\vec{AB}|$ . [2]

(b) Let  $\vec{AC} = \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix}$ . Find  $\hat{BAC}$ . [4]

Area for student response with a large watermark logo in the center. The logo is circular and contains the text "SAT PREP" at the top, "AHMEDABAD" at the bottom, and a large "SP" in the middle with a green leaf-like shape behind it. The response area is filled with horizontal dotted lines.



4. [Maximum mark: 8]

A discrete random variable  $X$  has the following probability distribution.

$X$	0	1	2	3
$P(X = x)$	0.475	$2k^2$	$\frac{k}{10}$	$6k^2$

- (a) Find the value of  $k$ . [4]
- (b) Write down  $P(X = 2)$ . [1]
- (c) Find  $P(X = 2 | X > 0)$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....




5. [Maximum mark: 5]

Let  $f(x) = 6 - \ln(x^2 + 2)$ , for  $x \in \mathbb{R}$ . The graph of  $f$  passes through the point  $(p, 4)$ , where  $p > 0$ .

(a) Find the value of  $p$ . [2]

(b) The following diagram shows part of the graph of  $f$ .



The region enclosed by the graph of  $f$ , the  $x$ -axis and the lines  $x = -p$  and  $x = p$  is rotated  $360^\circ$  about the  $x$ -axis. Find the volume of the solid formed.

[3]

Area for student response with horizontal dotted lines.



6. [Maximum mark: 6]

In the expansion of  $ax^3(2+ax)^{11}$ , the coefficient of the term in  $x^5$  is 11880.  
Find the value of  $a$ .

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



7. [Maximum mark: 7]

The heights of adult males in a country are normally distributed with a mean of 180 cm and a standard deviation of  $\sigma$  cm. 17% of these men are shorter than 168 cm. 80% of them have heights between  $(192 - h)$  cm and 192 cm.

Find the value of  $h$ .

A large rectangular area for writing, containing horizontal dotted lines for guidance. A watermark logo for 'SAT PREP AHMEDABAD' with the letters 'SP' in the center is visible over the writing area.



Do **not** write solutions on this page.

### Section B

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

8. [Maximum mark: 14]


Adam is a beekeeper who collected data about monthly honey production in his bee hives. The data for six of his hives is shown in the following table.

<b>Number of bees (<math>N</math>)</b>	190	220	250	285	305	320
<b>Monthly honey production in grams (<math>P</math>)</b>	900	1100	1200	1500	1700	1800

The relationship between the variables is modelled by the regression line with equation  $P = aN + b$ .

- (a) Write down the value of  $a$  and of  $b$ . [3]
- (b) Use this regression line to estimate the monthly honey production from a hive that has 270 bees. [2]

Adam has 200 hives in total. He collects data on the monthly honey production of all the hives. This data is shown in the following cumulative frequency graph.



(This question continues on the following page)



Do **not** write solutions on this page.

**(Question 8 continued)**

Adam's hives are labelled as low, regular or high production, as defined in the following table.


Type of hive	low	regular	high
Monthly honey production in grams ( $P$ )	$P \leq 1080$	$1080 < P \leq k$	$P > k$

- (c) Write down the number of low production hives. [1]

Adam knows that 128 of his hives have a regular production.

- (d) Find
- (i) the value of  $k$ ;
  - (ii) the number of hives that have a high production. [5]

- (e) Adam decides to increase the number of bees in each low production hive. Research suggests that there is a probability of 0.75 that a low production hive becomes a regular production hive. Calculate the probability that 30 low production hives become regular production hives. [3]





Please **do not** write on this page.

Answers written on this page  
will not be marked.



Do **not** write solutions on this page.

9. [Maximum mark: 14]

**Note: In this question, distance is in metres and time is in seconds.**

A particle P moves in a straight line for five seconds. Its acceleration at time  $t$  is given by  $a = 3t^2 - 14t + 8$ , for  $0 \leq t \leq 5$ .


(a) Write down the values of  $t$  when  $a = 0$ . [2]

(b) Hence or otherwise, find all possible values of  $t$  for which the velocity of P is decreasing. [2]

When  $t = 0$ , the velocity of P is  $3 \text{ m s}^{-1}$ .

(c) Find an expression for the velocity of P at time  $t$ . [6]

(d) Find the total distance travelled by P when its velocity is increasing. [4]



Do **not** write solutions on this page.

10. [Maximum mark: 17]

**Note: In this question, distance is in millimetres.**

Let  $f(x) = x + a \sin\left(x - \frac{\pi}{2}\right) + a$ , for  $x \geq 0$ .

(a) Show that  $f(2\pi) = 2\pi$ . [3]


The graph of  $f$  passes through the origin. Let  $P_k$  be any point on the graph of  $f$  with  $x$ -coordinate  $2k\pi$ , where  $k \in \mathbb{N}$ . A straight line  $L$  passes through all the points  $P_k$ .

(b) (i) Find the coordinates of  $P_0$  and of  $P_1$ . [6]

(ii) Find the equation of  $L$ . [2]

(c) Show that the distance between the  $x$ -coordinates of  $P_k$  and  $P_{k+1}$  is  $2\pi$ . [2]

(This question continues on the following page)






Do **not** write solutions on this page.

**(Question 10 continued)**

Diagram 1 shows a saw. The length of the toothed edge is the distance AB.

Diagram 1


**diagram not to scale**



The toothed edge of the saw can be modelled using the graph of  $f$  and the line  $L$ . Diagram 2 represents this model.

Diagram 2

**diagram not to scale**



The shaded part on the graph is called a tooth. A tooth is represented by the region enclosed by the graph of  $f$  and the line  $L$ , between  $P_k$  and  $P_{k+1}$ .

- (d) A saw has a toothed edge which is 300 mm long. Find the number of complete teeth on this saw.

[6]





Please **do not** write on this page.  
Answers written on this page  
will not be marked.



16EP16