

A-level
Topic :Differential Equation
May 2013-May 2023
Answers

Question 1

- (i) State $\frac{dV}{dt} = 80 - kV$ B1
 Correctly separate variables and attempt integration of one side M1
 Obtain $a \ln(80 - kV) = t$ or equivalent M1*
 Obtain $-\frac{1}{k} \ln(80 - kV) = t$ or equivalent A1
 Use $t = 0$ and $V = 0$ to find constant of integration or as limits M1 (dep*)
 Obtain $-\frac{1}{k} \ln(80 - kV) = t - \frac{1}{k} \ln 80$ or equivalent A1
 Obtain given answer $V = \frac{1}{k}(80 - 80e^{-kt})$ correctly A1 [7]
- (ii) Use iterative formula correctly at least once M1
 Obtain final answer 0.14 A1
 Show sufficient iterations to 4 s.f. to justify answer to 2 s.f. or show a sign change in the interval (0.135, 0.145) A1 [3]
- (iii) State a value between 530 and 540 cm³ inclusive B1
 State or imply that volume approaches 569 cm³ (allowing any value between 567 and 571 inclusive) B1 [2]

Question 2

- (i) Use any relevant method to determine a constant M1
 Obtain one of the values $A = 1, B = -2, C = 4$ A1
 Obtain a second value A1
 Obtain the third value A1 [4]
 [If A and C are found by the cover up rule, give B1 + B1 then M1A1 for finding B . If only one is found by the rule, give B1M1A1A1.]
- (ii) Separate variables and obtain one term by integrating $\frac{1}{y}$ or a partial fraction M1
 Obtain $\ln y = -\frac{1}{2} - 2 \ln(2x + 1) + c$, or equivalent A3✓

Question 3

- (i) Separate variables correctly and integrate at least one side M1
 Obtain term $\ln t$, or equivalent B1
 Obtain term of the form $a \ln(k - x^3)$ M1
 Obtain term $-\frac{2}{3} \ln(k - x^3)$, or equivalent A1
- EITHER:* Evaluate a constant or use limits $t = 1, x = 1$ in a solution containing $a \ln t$ and $b \ln(k - x^3)$ M1*
- Obtain correct answer in any form e.g. $\ln t = -\frac{2}{3} \ln(k - x^3) + \frac{2}{3} \ln(k - 1)$ A1
- Use limits $t = 4, x = 2$, and solve for k M1(dep*)
 Obtain $k = 9$ A1
- OR:* Using limits $t = 1, x = 1$ and $t = 4, x = 2$ in a solution containing $a \ln t$ and $b \ln(k - x^3)$ obtain an equation in k M1*
- Obtain a correct equation in any form, e.g. $\ln 4 = -\frac{2}{3} \ln(k - 8) + \frac{2}{3} \ln(k - 1)$ A1
- Solve for k M1(dep*)
 Obtain $k = 9$ A1

Substitute $k = 9$ and obtain $x = (9 - 8t^{\frac{3}{2}})^{\frac{1}{3}}$ A1 [9]

- (ii) State that x approaches $9^{\frac{1}{3}}$, or equivalent B1✓ [1]

Question 4

- (i) State or imply $V = \pi h^3$ B1
 State or imply $\frac{dV}{dt} = -k\sqrt{h}$ B1
 Use $\frac{dV}{dt} = \frac{dV}{dh} \cdot \frac{dh}{dt}$, or equivalent M1
 Obtain the given equation A1 [4]

[The M1 is only available if $\frac{dV}{dh}$ is in terms of h and has been obtained by a correct method.]

[Allow B1 for $\frac{dV}{dt} = k\sqrt{h}$ but withhold the final A1 until the polarity of the constant

$\frac{k}{3\pi}$ has been justified.]

- (ii) Separate variables and integrate at least one side M1
 Obtain terms $\frac{2}{5}h^{\frac{5}{2}}$ and $-At$, or equivalent A1
 Use $t = 0, h = H$ in a solution containing terms of the form $ah^{\frac{5}{2}}$ and $bt + c$ M1
 Use $t = 60, h = 0$ in a solution containing terms of the form $ah^{\frac{5}{2}}$ and $bt + c$ M1
 Obtain a correct solution in any form, e.g. $\frac{2}{5}h^{\frac{5}{2}} = \frac{1}{150}H^{\frac{5}{2}}t + \frac{2}{5}H^{\frac{5}{2}}$ A1

- (ii) Obtain final answer $t = 60 \left(1 - \left(\frac{h}{H} \right)^{\frac{5}{2}} \right)$, or equivalent A1 [6]

- (iii) Substitute $h = \frac{1}{2}H$ and obtain answer $t = 49.4$ B1 [1]

Question 5

- Use $2\cos^2 x = 1 + \cos 2x$ or equivalent B1
 Separate variables and integrate at least one side M1
 Obtain $\ln(y^3 + 1) = \dots$ or equivalent A1
 Obtain $\dots = 2x + \sin 2x$ or equivalent A1
 Use $x = 0, y = 2$ to find constant of integration (or as limits) in an expression containing at least two terms of the form $a \ln(y^3 + 1), bx$ or $c \sin 2x$ M1*
 Obtain $\ln(y^3 + 1) = 2x + \sin 2x + \ln 9$ or equivalent e.g. implied by correct constant A1
 Identify at least one of $\frac{1}{2}\pi$ and $\frac{3}{2}\pi$ as x -coordinate at stationary point B1
 Use correct process to find y -coordinate for at least one x -coordinate M1(d*M)
 Obtain 5.9 A1
 Obtain 48.1 A1 [10]

Question 6

- Separate variables correctly and recognisable attempt at integration of at least one side M1
 Obtain $\ln y$, or equivalent B1
 Obtain $k \ln(2 + e^{3x})$ B1
 Use $y(0) = 36$ to find constant in $y = A(2 + e^{3x})^k$ or $\ln y = k \ln(2 + e^{3x}) + c$ or equivalent M1*
 Obtain equation correctly without logarithms from $\ln y = \ln \left(A(2 + e^{3x})^k \right)$ *M1
 Obtain $y = 4(2 + e^{3x})^2$ A1 [6]

Question 7

- (i) State or imply $\frac{dN}{dt} = kN(1 - 0.01N)$ and obtain the given answer $k = 0.02$ B1 **1**
- (ii) Separate variables and attempt integration of at least one side M1
 Integrate and obtain term $0.02t$, or equivalent A1
- Carry out a relevant method to obtain A or B such that $\frac{1}{N(1 - 0.01N)} \equiv \frac{A}{N} + \frac{B}{1 - 0.01N}$, or
 equivalent M1*
 Obtain $A = 1$ and $B = 0.01$, or equivalent A1
 Integrate and obtain terms $\ln N - \ln(1 - 0.01N)$, or equivalent A1✓
 Evaluate a constant or use limits $t = 0$, $N = 20$ in a solution with terms $a \ln N$ and
 $b \ln(1 - 0.01N)$, $ab \neq 0$ M1(dep*)
 Obtain correct answer in any form, e.g. $\ln N - \ln(1 - 0.01N) = 0.02t + \ln 25$ A1
 Rearrange and obtain $t = 50 \ln(4N / (100 - N))$, or equivalent A1 **8**
- (iii) Substitute $N = 40$ and obtain $t = 49.0$ B1 **1**

Question 8

- Separate variables correctly and attempt integration of at least one side B1
 Obtain term in the form $a\sqrt{2x+1}$ M1
 Express $1/(\cos^2 \theta)$ as $\sec^2 \theta$ B1
 Obtain term of the form $k \tan \theta$ M1
 Evaluate a constant, or use limits $x = 0$, $\theta = \frac{1}{4}\pi$ in a solution with terms $a\sqrt{2x+1}$ and $k \tan \theta$,
 $ak \neq 0$ M1
 Obtain correct solution in any form, e.g. $\sqrt{2x+1} = \frac{1}{2} \tan \theta + \frac{1}{2}$ A1
 Rearrange and obtain $x = \frac{1}{8}(\tan \theta + 1)^2 - \frac{1}{2}$, or equivalent A1 **7**

Question 9

- (i) Separate variables correctly and attempt to integrate at least one side B1
 Obtain term $\ln R$ B1
 Obtain $\ln x - 0.57x$ B1
 Evaluate a constant or use limits $x = 0.5$, $R = 16.8$, in a solution containing terms of the form $a \ln R$ and $b \ln x$ M1
 Obtain correct solution in any form A1
 Obtain a correct expression for R , e.g. $R = xe^{(3.80 - 0.57x)}$, $R = 44.7xe^{-0.57x}$ or
 $R = 33.6xe^{(0.285 - 0.57x)}$ A1 [6]
- (ii) Equate $\frac{dR}{dx}$ to zero and solve for x M1
 State or imply $x = 0.57^{-1}$, or equivalent, e.g. 1.75 A1
 Obtain $R = 28.8$ (allow 28.9) A1 [3]

Question 10

- (i) Sensibly separate variables and attempt integration of at least one side M1
 Obtain $2y^{\frac{1}{2}} = \dots$ or equivalent A1
 Correct integration by parts of $x \sin \frac{1}{3}x$ as far as $ax \cos \frac{1}{3}x \pm \int b \cos \frac{1}{3}x dx$ M1
 Obtain $-3x \cos \frac{1}{3}x + \int 3 \cos \frac{1}{3}x dx$ or equivalent A1
 Obtain $-3x \cos \frac{1}{3}x + 9 \sin \frac{1}{3}x$ or equivalent A1
 Obtain $y = \left(-\frac{3}{10}x \cos \frac{1}{3}x + \frac{9}{10} \sin \frac{1}{3}x + c \right)^2$ or equivalent A1 [6]
- (ii) Use $x = 0$ and $y = 100$ to find constant M*1
 Substitute 25 and calculate value of y DM*1
 Obtain 203 A1 [3]

Question 11

Separate variables and factorise to obtain $\frac{dy}{(3y+1)(y+3)} = 4x \, dx$ or equivalent	B1	
State or imply the form $\frac{A}{3y+1} + \frac{B}{y+3}$ and use a relevant method to find A or B	M1	
Obtain $A = \frac{3}{8}$ and $B = -\frac{1}{8}$	A1	
Integrate to obtain form $k_1 \ln(3y+1) + k_2 \ln(y+3)$	M1	
Obtain correct $\frac{1}{8} \ln(3y+1) - \frac{1}{8} \ln(y+3) = 2x^2$ or equivalent	A1	
Substitute $x = 0$ and $y = 1$ in equation of form $k_1 \ln(3y+1) + k_2 \ln(y+3) = k_3 x^2 + c$ to find a value of c	M1	
Obtain $c = 0$	A1	
Use correct process to obtain equation without natural logarithm present	M1	
Obtain $y = \frac{3e^{16x^2} - 1}{3 - e^{16x^2}}$ or equivalent	A1	[9]

Question 12

(i) Separate variables correctly and attempt integration of one side	B1	
Obtain term $\ln x$	B1	
Obtain term of the form $a \ln(k + e^{-t})$	M1	
Obtain term $-\ln(k + e^{-t})$	A1	
Evaluate a constant or use limits $x = 10, t = 0$ in a solution containing terms $a \ln(k + e^{-t})$ and $b \ln x$	M1*	
Obtain correct solution in any form, e.g. $\ln x - \ln 10 = -\ln(k + e^{-t}) + \ln(k+1)$	A1	[6]
(ii) Substitute $x = 20, t = 1$ and solve for k	M1(dep*)	
Obtain the given answer	A1	[2]
(iii) Using $e^{-t} \rightarrow 0$ and the given value of k , find the limiting value of x	M1	
Justify the given answer	A1	[2]

Question 13

- (i) Separate variables correctly and integrate one side B1
 Obtain term $2\sqrt{M}$, or equivalent B1
 Obtain term $50k \sin(0.02t)$, or equivalent B1
 Evaluate a constant of integration, or use limits $M = 100, t = 0$ in a solution with terms of
 the form $a\sqrt{M}$ and $b \sin(0.02t)$ M1*
 Obtain correct solution in any form, e.g. $2\sqrt{M} = 50k \sin(0.02t) + 20$ A1 **5**
- (ii) Use values $M = 196, t = 50$ and calculate k M1(dep*)
 Obtain answer $k = 0.190$ A1 **2**
- (iii) State an expression for M in terms of t , e.g. $M = (4.75 \sin(0.02t) + 10)^2$ M1(dep*)
 State that the least possible number of micro-organisms is 28 or 27.5 or 27.6 (27.5625) A1 **2**

Question 14

- Separate variables and integrate one side B1
 Obtain term $\ln(x + 2)$ B1
 Use $\cos 2A$ formula to express $\sin^2 2\theta$ in the form $a + b \cos 4\theta$ M1
 Obtain correct form $(1 - \cos 4\theta) / 2$, or equivalent A1
 Integrate and obtain term $\frac{1}{2}\theta - \frac{1}{8}\sin 4\theta$, or equivalent A1[✓]
 Evaluate a constant, or use $\theta = 0, x = 0$ as limits in a solution containing terms
 $c \ln(x + 2), d \sin(4\theta), e\theta$ M1
 Obtain correct solution in any form, e.g. $\ln(x + 2) = \frac{1}{2}\theta - \frac{1}{8}\sin 4\theta + \ln 2$ A1
 Use correct method for solving an equation of the form $\ln(x + 2) = f$ M1
 Obtain answer $x = 0.962$ A1 [9]

Question 15

- (i) State $\frac{dN}{dt} = k(N - 150)$ B1 [1]
- (ii) Substitute $\frac{dN}{dt} = 60$ and $N = 900$ to find value of k M1
 Obtain $k = 0.08$ A1
 Separate variables and obtain general solution involving $\ln(N - 150)$ M1*
 Obtain $\ln(N - 150) = 0.08t + c$ (following their k) or $\ln(N - 150) = kt + c$ A1[✓]
 Substitute $t = 0$ and $N = 650$ to find c dep M1*
 Obtain $\ln(N - 150) = 0.08t + \ln 500$ or equivalent A1
 Obtain $N = 500e^{0.08t} + 150$ A1 [7]
- (iii) Either Substitute $t = 15$ to find N or solve for t with $N = 2000$ M1
Obtain Either $N = 1810$ or $t = 16.4$ and conclude target not met A1 [2]

Question 16

- (i) Separate variables and attempt integration of one side **M1**
 Obtain term $-e^{-y}$ **A1**
 Integrate xe^x by parts reaching $xe^x \pm \int e^x dx$ **M1**
 Obtain integral $xe^x - e^x$ **A1**
 Evaluate a constant, or use limits $x = 0, y = 0$ **M1**
 Obtain correct solution in any form **A1**
 Obtain final answer $y = -\ln(e^x(1-x))$, or equivalent **A1** [7]
- (ii) Justify the given statement **B1** [1]

Question 17

- Separate variables and attempt integration of at least one side **M1***
 Obtain term $\ln y$ **A1**
 Obtain terms $\ln x - x^2$ **A1**
 Use $x = 1$ and $y = 2$ to evaluate a constant, or as limits **DM1***
 Obtain correct solution in any form, e.g. $\ln y = \ln x - x^2 + \ln 2 + 1$ **A1**
 Obtain correct expression for y , free of logarithms, i.e. $y = 2x \exp(1 - x^2)$ **A1**
 [6]

Question 18

- (i) Separate variables correctly and attempt integration of at least one side **B1**
 Obtain term $\ln x$ **B1**
 Obtain term of the form $k \ln(3 + \cos 2\theta)$, or equivalent **M1**
 Obtain term $-\frac{1}{2} \ln(3 + \cos 2\theta)$, or equivalent **A1**
 Use $x = 3, \theta = \frac{1}{4}\pi$ to evaluate a constant or as limits in a solution with terms $a \ln x$ and $b \ln(3 + \cos 2\theta)$, where $ab \neq 0$ **M1**
 State correct solution in any form, e.g. $\ln x = -\frac{1}{2} \ln(3 + \cos 2\theta) + \frac{3}{2} \ln 3$ **A1**
 Rearrange in a correct form, e.g. $x = \sqrt{\left(\frac{27}{3 + \cos 2\theta}\right)}$ **A1** [7]
- (ii) State answer $x = 3\sqrt{3}/2$, or exact equivalent (accept decimal answer in [2.59, 2.60]) **B1** [1]

Question 19

Separate variables and make reasonable attempt at integration of either integral	M1
Obtain term $\frac{1}{2}e^{2y}$	B1
Use Pythagoras	M1
Obtain terms $\tan x - x$	A1
Evaluate a constant or use $x = 0, y = 0$ as limits in a solution containing terms $ae^{\pm 2y}$ and $b \tan x, (ab \neq 0)$	M1
Obtain correct solution in any form, e.g. $\frac{1}{2}e^{2y} = \tan x - x + \frac{1}{2}$	A1
Set $x = \frac{1}{4}\pi$ and use correct method to solve an equation of the form $e^{\pm 2y} = a$ or $e^{\pm y} = a$, where $a > 0$	M1
Obtain answer $y = 0.179$	A1
	[8]

Question 20

(i)	Separate variables correctly and integrate at least one side	M1
	Integrate and obtain term kt , or equivalent	A1
	Carry out a relevant method to obtain A and B such that $\frac{1}{x(4-x)} \equiv \frac{A}{x} + \frac{B}{4-x}$, or equivalent	M1*
	Obtain $A = B = \frac{1}{4}$, or equivalent	A1
	Integrate and obtain terms $\frac{1}{4} \ln x - \frac{1}{4} \ln(4-x)$, or equivalent	A1✓
	EITHER: Use a pair of limits in an expression containing $p \ln x, q \ln(4-x)$ and rt and evaluate a constant	DM1
	Obtain correct answer in any form, e.g. $\ln x - \ln(4-x) = 4kt - \ln 9$,	
	or $\ln\left(\frac{x}{4-x}\right) = 4kt - 8k$	A1
	Use a second pair of limits and determine k	DM1
	Obtain the given exact answer correctly	A1
	OR: Use both pairs of limits in a definite integral	M1*
	Obtain the given exact answer correctly	A1
	Substitute k and either pair of limits in an expression containing $p \ln x, q \ln(4-x)$ and rt and evaluate a constant	DM1
	Obtain $\ln \frac{x}{4-x} = t \ln 3 - \ln 9$ or equivalent	A1
		[9]
<hr style="border-top: 1px dashed black;"/>		
(ii)	Substitute $x = 3.6$ and solve for t	M1
	Obtain answer $t = 4$	A1
		[2]

Question 21

(i)	State equation $\frac{dy}{dx} = \frac{1}{2}xy$	B1	[1]
(ii)	Separate variables correctly and attempts to integrate one side of equation Obtain terms of the form $a \ln y$ and bx^2 Use $x = 0$ and $y = 2$ to evaluate a constant, or as limits, in expression containing $a \ln y$ or bx^2 Obtain correct solution in any form, e.g. $\ln y = \frac{1}{4}x^2 + \ln 2$ Obtain correct expression for y , e.g. $y = 2e^{\frac{1}{4}x^2}$	M1 A1 M1 A1 A1	[5]
(iii)	Show correct sketch for $x \geq 0$. Needs through (0, 2) and rapidly increasing positive gradient.	B1	[1]

Question 22

(i)	State or imply $\frac{dV}{dt} = 2\frac{dh}{dt}$		B1
	State or imply $\frac{dV}{dt} = 1 - 0.2\sqrt{h}$		B1
	Obtain the given answer correctly		B1
	Total:		3
(ii)	State or imply $du = -\frac{1}{2\sqrt{h}} dh$, or equivalent		B1
	Substitute for h and dh throughout		M1
	Obtain $T = \int_3^5 \frac{20(5-u)}{u} du$, or equivalent		A1
	Integrate and obtain terms $100 \ln u - 20u$, or equivalent		A1
	Substitute limits $u = 3$ and $u = 5$ correctly		M1
	Obtain answer 11.1, with no errors seen		A1
	Total:		6

Question 23

(i)	Carry out a relevant method to obtain A and B such that $\frac{1}{x(2x+3)} \equiv \frac{A}{x} + \frac{B}{2x+3}$, or equivalent	M1
	Obtain $A = \frac{1}{3}$ and $B = -\frac{2}{3}$, or equivalent	A1
	Total:	2
(ii)	Separate variables and integrate one side	B1
	Obtain term $\ln y$	B1
	Integrate and obtain terms $\frac{1}{3} \ln x - \frac{1}{3} \ln(2x+3)$, or equivalent	B2 FT
	Use $x = 1$ and $y = 1$ to evaluate a constant, or as limits, in a solution containing $a \ln y$, $b \ln x$, $c \ln(2x+3)$	M1
	Obtain correct solution in any form, e.g. $\ln y = \frac{1}{3} \ln x - \frac{1}{3} \ln(2x+3) + \frac{1}{3} \ln 5$	A1
	Obtain answer $y = 1.29$ (3s.f. only)	A1
	Total:	7

Question 24

(i)	State $\frac{dy}{dt} = -\frac{2y}{(1+t)^2}$, or equivalent	B1
	Separate variables correctly and attempt integration of one side	M1
	Obtain term $\ln y$, or equivalent	A1
	Obtain term $\frac{2}{(1+t)}$, or equivalent	A1
	Use $y = 100$ and $t = 0$ to evaluate a constant, or as limits in an expression containing terms of the form $a \ln y$ and $\frac{b}{1+t}$	M1
	Obtain correct solution in any form, e.g. $\ln y = \frac{2}{1+t} - 2 + \ln 100$	A1
	Total:	6

(ii)	State that the mass of B approaches $\frac{100}{e^2}$, or exact equivalent	B1
	State or imply that the mass of A tends to zero	B1
	Total:	2

Question 25

(i)	Justify the given differential equation	B1
	Total:	1
(ii)	Separate variables correctly and attempt to integrate one side	B1
	Obtain term kt , or equivalent	B1
	Obtain term $-\ln(50-x)$, or equivalent	B1
	Evaluate a constant, or use limits $x=0, t=0$ in a solution containing terms $a \ln(50-x)$ and bt	M1*
	Obtain solution $-\ln(50-x) = kt - \ln 50$, or equivalent	A1
	Use $x=25, t=10$ to determine k	DM1
	Obtain correct solution in any form, e.g. $\ln 50 - \ln(50-x) = \frac{1}{10}(\ln 2)t$	A1
	Obtain answer $x = 50(1 - \exp(-0.0693t))$, or equivalent	A1
	Total:	8

Question 26

Separate variables correctly and attempt integration of one side	B1
Obtain term $\tan y$, or equivalent	B1
Obtain term of the form $k \ln \cos x$, or equivalent	M1
Obtain term $-4 \ln \cos x$, or equivalent	A1
Use $x = 0$ and $y = \frac{1}{4}\pi$ in solution containing $a \tan y$ and $b \ln \cos x$ to evaluate a constant, or as limits	M1
Obtain correct solution in any form, e.g. $\tan y = 4 \ln \sec x + 1$	A1
Substitute $y = \frac{1}{3}\pi$ in solution containing terms $a \tan y$ and $b \ln \cos x$, and use correct method to find x	M1
Obtain answer $x = 0.587$	A1
	8

Question 27

Separate variables and obtain $\int \frac{1}{y} dy = \int \frac{x+2}{x+1} dx$	B1
Obtain term $\ln y$	B1
Use an appropriate method to integrate $(x+2)/(x+1)$	*M1
Obtain integral $x + \ln(x+1)$, or equivalent, e.g. $\ln(x+1) + x + 1$	A1
Use $x = 1$ and $y = 2$ to evaluate a constant, or as limits	DM1
Obtain correct solution in x and y in any form e.g. $\ln y = x + \ln(x+1) - 1$	A1
Obtain answer $y = (x+1)e^{x-1}$	A1
	7

Question 28

(i)	Show sufficient working to justify the given statement	AG	B1
			1
(ii)	Separate variables correctly and attempt integration of at least one side		B1
	Obtain term $\frac{1}{2}x^2$		B1
	Obtain terms $\tan^2 \theta + \tan \theta$, or $\sec^2 \theta + \tan \theta$		B1 + B1
	Evaluate a constant, or use limits $x = 1$, $\theta = \frac{1}{4}\pi$, in a solution with two terms of the form ax^2 and $b \tan \theta$, where $ab \neq 0$		M1
	State correct answer in any form, e.g. $\frac{1}{2}x^2 = \tan^2 \theta + \tan \theta - \frac{3}{2}$		A1
	Substitute $\theta = \frac{1}{3}\pi$ and obtain $x = 2.54$		A1
			7

Question 29

(i)	Separate variables correctly and integrate at least one side		B1
	Obtain term $\ln x$		B1
	Obtain term $-\frac{2}{3}kt\sqrt{t}$, or equivalent		B1
	Evaluate a constant, or use limits $x = 100$ and $t = 0$, in a solution containing terms $a \ln x$ and $b t\sqrt{t}$		M1
	Obtain correct solution in any form, e.g. $\ln x = -\frac{2}{3}kt\sqrt{t} + \ln 100$		A1
(ii)	Substitute $x = 80$ and $t = 25$ to form equation in k		M1
	Substitute $x = 40$ and eliminate k		M1
	Obtain answer $t = 64.1$		A1

Question 30

(i)	Fully justify the given statement	B1	Some indication of use of gradient of curve = gradient of tangent (PT) and no errors seen /no incorrect statements
		1	
(ii)	Separate variables and attempt integration of at least one side Obtain terms $\ln y$ and $\frac{1}{2}x$	B1 B1	Must be working from $\int \frac{1}{y} dy = \int k dx$ B marks are not available for fortuitously correct answers
	Use $x = 4, y = 3$ to evaluate a constant or as limits in a solution with terms $a \ln y$ and bx , where $ab \neq 0$	M1	
	Obtain correct solution in any form	A1	$\ln y = \frac{1}{2}x + \ln 3 - 2$
	Obtain answer $y = 3e^{\frac{1}{2}x-2}$, or equivalent	A1	Accept $y = e^{\frac{1}{2}x + \ln 3 - 2}$, $y = e^{\frac{x-1.80}{2}}$, $y = 3\sqrt{e^{x-4}}$ $ y = \dots$ scores A0
		5	

Question 31

(i)	Carry out relevant method to find A and B such that $\frac{1}{4-y^2} \equiv \frac{A}{2+y} + \frac{B}{2-y}$	M1
	Obtain $A = B = \frac{1}{4}$	A1
	Total:	2
(ii)	Separate variables correctly and integrate at least one side to obtain one of the terms $a \ln x, b \ln(2+y)$ or $c \ln(2-y)$	M1
	Obtain term $\ln x$	B1
	Integrate and obtain terms $\frac{1}{4} \ln(2+y) - \frac{1}{4} \ln(2-y)$	A1FT
	Use $x = 1$ and $y = 1$ to evaluate a constant, or as limits, in a solution containing at least two terms of the form $a \ln x, b \ln(2+y)$ and $c \ln(2-y)$	M1
	Obtain a correct solution in any form, e.g. $\ln x = \frac{1}{4} \ln(2+y) - \frac{1}{4} \ln(2-y) - \frac{1}{4} \ln 3$	A1
	Rearrange as $\frac{2(3x^4-1)}{(3x^4+1)}$, or equivalent	A1
	Total:	6

Question 32

Separate variables correctly and integrate at least one side	B1
Obtain term $\ln y$	B1
Obtain terms $2 \ln x - \frac{1}{2} x^2$	B1+B1
Use $x = 1, y = 1$ to evaluate a constant, or as limits	M1
Obtain correct solution in any form, e.g. $\ln y = 2 \ln x - \frac{1}{2} x^2 + \frac{1}{2}$	A1
Rearrange as $y = x^2 \exp\left(\frac{1}{2} - \frac{1}{2} x^2\right)$, or equivalent	A1
	7

Question 33

State equation $\frac{dy}{dx} = k \frac{y^2}{x}$, or equivalent	B1	SC: If $k = 1$ seen or implied give B0 and then allow B1B1B0M1, max 3/8.
Separate variables correctly and integrate at least one side	B1	$\int \frac{k}{x} dx = \int \frac{1}{y^2} dy$ Allow with incorrect value substituted for k
Obtain terms $-\frac{1}{y}$ and $k \ln x$	B1 + B1	Incorrect k used scores max. B1B0
Use given coordinates correctly to find k and/or a constant of integration C in an equation containing terms $\frac{a}{y}$, $b \ln x$ and C	M1	SC: If an incorrect method is used to find k , M1 is allowable for a correct method to find C
Obtain $k = \frac{1}{2}$ and $c = -1$, or equivalent	A1 + A1	$\frac{1}{2} \ln x = 1 - \frac{1}{y}$ A0 for fortuitous answers.
Obtain answer $y = \frac{2}{2 - \ln x}$, or equivalent, and ISW	A1	$y = \frac{-1}{-1 + \ln \sqrt{x}}$
		SC: MR of the fraction. $\frac{dy}{dx} = k \frac{y^2}{x^2}$ B1 Separate variables and integrate B1 $\frac{-1}{y} = \frac{-k}{x} (+C)$ B1+B1 Substitute to find k and/or c M1 $k = \frac{e}{2(e-1)}, c = \frac{2-e}{2(e-1)}$ A1+A1 Answer A0
	8	

Question 34

Separate variables correctly and attempt integration of at least one side	B1
Obtain term $-\frac{1}{2y^2}$, or equivalent	B1
Obtain term $-k e^{-x}$	B1
Use a pair of limits, e.g. $x = 0, y = 1$ to obtain an equation in k and an arbitrary constant c	M1
Use a second pair of limits, e.g. $x = 1, y = \sqrt{e}$, to obtain a second equation and solve for k or for c	M1
Obtain $k = \frac{1}{2}$ and $c = 0$	A1
Obtain final answer $y = e^{\frac{1}{2}x}$, or equivalent	A1
	7

Question 35

(i)	Use chain rule	M1	$k \cos \theta \sin^{-3} \theta (= -k \operatorname{cosec}^2 \theta \cot \theta)$ Allow M1 for $-2 \cos \theta \sin^{-1} \theta$
	Obtain correct answer in any form	A1	e.g. $-2 \operatorname{cosec}^2 \theta \cot \theta$, $\frac{-2 \cos \theta}{\sin^3 \theta}$ Accept $\frac{-2 \sin \theta \cos \theta}{\sin^4 \theta}$
		2	
(ii)	Separate variables correctly and integrate at least one side	B1	$\int x \, dx = \int -\operatorname{cosec}^2 \theta \cot \theta \, d\theta$
	Obtain term $\frac{1}{2}x^2$	B1	
	Obtain term of the form $\frac{k}{\sin^2 \theta}$	M1*	or equivalent
	Obtain term $\frac{1}{2 \sin^2 \theta}$	A1	or equivalent
	Use $x = 4, \theta = \frac{1}{6}\pi$ to evaluate a constant, or as limits, in a solution with terms ax^2 and $\frac{b}{\sin^2 \theta}$, where $ab \neq 0$	DM1	Dependent on the preceding M1
	Obtain solution $x = \sqrt{(\operatorname{cosec}^2 \theta + 12)}$	A1	or equivalent
		6	

Question 36

(i)	Separate variables correctly and attempt integration of at least one side	B1	$\int e^{-y} dy = \int xe^x dx$
	Obtain term $-e^{-y}$	B1	B0B1 is possible
	Commence integration by parts and reach $xe^x \pm \int e^x dx$	M1	B0B0M1A1 is possible
	Obtain $xe^x - e^x$	A1	or equivalent
			B1B1M1A1 is available if there is no constant of integration
	Use $x = 0, y = 0$ to evaluate a constant, or as limits in a definite integral, in a solution with terms ae^{-y}, bxe^x and ce^x , where $abc \neq 0$	M1	Must see this step
	Obtain correct solution in any form	A1	e.g. $e^{-y} = e^x - xe^x$
	Rearrange as $y = -\ln(1-x) - x$	A1	or equivalent e.g. $y = \ln \frac{1}{e^x(1-x)}$
			ISW
			7
(ii)	Justify the given statement	B1	e.g. require $1-x > 0$ for the \ln term to exist, hence $x < 1$ Must be considering the range of values of x , and must be relevant to <i>their</i> y involving $\ln(1-x)$
			1

Question 37

Separate variables correctly and integrate at least one side	B1
Obtain term $\ln(x+1)$	B1
Obtain term of the form $a \ln(y^2 + 5)$	M1
Obtain term $\frac{1}{2} \ln(y^2 + 5)$	A1
Use $y = 2, x = 0$ to determine a constant, or as limits, in a solution containing terms $a \ln(y^2 + 5)$ and $b \ln(x+1)$, where $ab \neq 0$	M1
Obtain correct solution in any form	A1
Obtain final answer $y^2 = 9(x+1)^2 - 5$	A1
	7

Question 38

(i)	State $\frac{dN}{dt} = ke^{-0.02t}N$ and show $k = -0.01$	B1	OE ($-10 = k \times 1 \times 1000$)
		1	
(ii)	Separate variables correctly and integrate at least one side	B1	$\int \frac{1}{N} dN = \int -0.01e^{-0.02t} dt$
	Obtain term $\ln N$	B1	OE
	Obtain term $0.5e^{-0.02t}$	B1	OE
	Use $N = 1000$, $t = 0$ to evaluate a constant, or as limits, in a solution with terms $a \ln N$ and $be^{-0.02t}$, where $ab \neq 0$	M1	
	Obtain correct solution in any form e.g. $\ln N - \ln 1000 = 0.5(e^{-0.02t} - 1)$	A1	$\ln 1000 - \frac{1}{2} = 6.41$
	Substitute $N = 800$ and obtain $t = 29.6$	A1	
		6	
(iii)	State that N approaches $\frac{1000}{\sqrt{e}}$	B1	Accept 606 or 607 or 606.5
		1	

Question 39

Separate variables correctly to obtain $\int \frac{1}{x+2} dx = \int \cot \frac{1}{2} \theta d\theta$	B1	Or equivalent integrands. Integral signs SOI
Obtain term $\ln(x+2)$	B1	Modulus signs not needed.
Obtain term of the form $k \ln \sin \frac{1}{2} \theta$	M1	
Obtain term $2 \ln \sin \frac{1}{2} \theta$	A1	
Use $x = 1$, $\theta = \frac{1}{3} \pi$ to evaluate a constant, or as limits, in an expression containing $p \ln(x+2)$ and $q \ln(\sin \frac{1}{2} \theta)$	M1	Reach $C =$ an expression or a decimal value
Obtain correct solution in any form e.g. $\ln(x+2) = 2 \ln \sin \frac{1}{2} \theta + \ln 12$	A1	$\ln 12 = 2.4849 \dots$ Accept constant to at least 3 s.f. Accept with $\ln 3 - 2 \ln \frac{1}{2}$
Remove logarithms and use correct double angle formula	M1	Need correct algebraic process. $\left(\frac{x+2}{12} = \frac{1-\cos \theta}{2} \right)$
Obtain answer $x = 4 - 6 \cos \theta$	A1	
	8	

Question 40

(i)	Separate variables correctly and integrate one side	B1
	Obtain term $0.2t$, or equivalent	B1
	Carry out a relevant method to obtain A and B such that $\frac{1}{(20-x)(40-x)} \equiv \frac{A}{20-x} + \frac{B}{40-x}$	*M1
	Obtain $A = \frac{1}{20}$ and $B = -\frac{1}{20}$	A1
	Integrate and obtain terms $-\frac{1}{20}\ln(20-x) + \frac{1}{20}\ln(40-x)$ OE	A1FT +A1FT
	Use $x = 10, t = 0$ to evaluate a constant, or as limits	DM1
	Obtain correct answer in any form	A1
	Obtain final answer $x = \frac{60e^{4t} - 40}{3e^{4t} - 1}$	A1
		9
(ii)	State that x approaches 20	B1
		1

Question 41

(a)	Separate variables correctly and attempt integration of at least one side	B1
	Obtain term of the form $a \tan^{-1}(2y)$	M1
	Obtain term $\frac{1}{2} \tan^{-1}(2y)$	A1
	Obtain term $-e^{-x}$	B1
	Use $x = 1, y = 0$ to evaluate a constant or as limits in a solution containing terms of the form $a \tan^{-1}(by)$ and $ce^{\pm x}$	M1
	Obtain correct answer in any form	A1
	Obtain final answer $y = \frac{1}{2} \tan(2e^{-1} - 2e^{-x})$, or equivalent	A1
		7
(b)	State that y approaches $\frac{1}{2} \tan(2e^{-1})$, or equivalent	B1FT
		1

Question 42

(a)	State $\frac{dy}{dx} = k \frac{y}{x\sqrt{x}}$, or equivalent	B1
	Separate variables correctly and attempt integration of at least one side	M1
	Obtain term $\ln y$, or equivalent	A1
	Obtain term $-2k \frac{1}{\sqrt{x}}$, or equivalent	A1
	Use given coordinates to find k or a constant of integration c in a solution containing terms of the form $a \ln y$ and $\frac{b}{\sqrt{x}}$, where $ab \neq 0$	M1
	Obtain $k = 1$ and $c = 2$	A1 + A1
	Obtain final answer $y = \exp\left(-\frac{2}{\sqrt{x}} + 2\right)$, or equivalent	A1
		8
(b)	State that y approaches e^2 (FT their c in part (a) of the correct form)	B1FT
		1

Question 43

Separate variables correctly and integrate at least one side	B1
Obtain term $\ln(y-1)$	B1
Carry out a relevant method to determine A and B such that $\frac{1}{(x+1)(x+3)} \equiv \frac{A}{x+1} + \frac{B}{x+3}$	M1
Obtain $A = \frac{1}{2}$ and $B = -\frac{1}{2}$	A1
Integrate and obtain terms $\frac{1}{2}\ln(x+1) - \frac{1}{2}\ln(x+3)$ $\frac{1}{2}\ln(x+1) - \frac{1}{2}\ln(x+3)$, or equivalent (FT is on A and B)	A1 FT + A1 FT
Use $x=0, y=2$ to evaluate a constant, or as limits in a solution containing terms of the form $a \ln(y-1), b \ln(x+1)$ and $c \ln(x+3)$, where $abc \neq 0$	M1
Obtain correct answer in any form	A1
Obtain final answer $y = 1 + \sqrt{\frac{3x+3}{x+3}}$, or equivalent	A1
	9

Question 44

(a)	State or imply $\frac{dV}{dt} = -k\sqrt{h}$	B1
	State or imply $\frac{dV}{dh} = 2\pi rh - \pi h^2$, or equivalent	B1
	Use $\frac{dV}{dt} = \frac{dV}{dh} \cdot \frac{dh}{dt}$	M1
	Obtain the given answer correctly	A1
		4
(b)	Separate variables and attempt integration of at least one side	M1
	Obtain terms $\frac{4}{3}rh^{\frac{3}{2}} - \frac{2}{5}h^{\frac{5}{2}}$ and $-Bt$	A3, 2, 1, 0
	Use $t=0, h=r$ to find a constant of integration c	M1
	Use $t=14, h=0$ to find B	M1
	Obtain correct c and B , e.g. $c = \frac{14}{15}r^{\frac{5}{2}}, B = \frac{1}{15}r^{\frac{5}{2}}$	A1
	Obtain final answer $t = 14 - 20\left(\frac{h}{r}\right)^{\frac{3}{2}} + 6\left(\frac{h}{r}\right)^{\frac{5}{2}}$, or equivalent	A1
		8

Question 45

Separate variables correctly and attempt integration of at least one side	B1	$\frac{1}{y} dy = \frac{1-2x^2}{x} dx$
Obtain term $\ln y$	B1	
Obtain terms $\ln x - x^2$	B1	
Use $x = 1, y = 1$ to evaluate a constant, or as limits, in a solution containing at least 2 terms of the form $a \ln y, b \ln x$ and cx^2	M1	The 2 terms of required form must be from correct working e.g. $\ln y = \ln x - x^2 + 1$
Obtain correct solution in any form	A1	
Rearrange and obtain $y = xe^{1-x^2}$	A1	OE
	6	

Question 46

(a)	Correct separation of variables	B1	$\int \sec^2 2x dx = \int e^{-3t} dt$ Needs correct structure
	Obtain term $-\frac{1}{3}e^{-3t}$	B1	
	Obtain term of the form $k \tan 2x$	M1	From correct working
	Obtain term $\frac{1}{2} \tan 2x$	A1	
	Use $x = 0, t = 0$ to evaluate a constant, or as limits in a solution containing terms of the form $a \tan 2x$ and be^{-3t} , where $ab \neq 0$	M1	
	Obtain correct solution in any form	A1	e.g. $\frac{1}{2} \tan 2x = -\frac{1}{3}e^{-3t} + \frac{1}{3}$
	Obtain final answer $x = \frac{1}{2} \tan^{-1} \left(\frac{2}{3} (1 - e^{-3t}) \right)$	A1	
		7	
(b)	State that x approaches $\frac{1}{2} \tan^{-1} \left(\frac{2}{3} \right)$	B1 FT	Correct value. Accept $x \rightarrow 0.294$ The FT is dependent on letting $e^{-3t} \rightarrow 0$ in a solution containing e^{-3t} .
		1	

Question 47

Separate variables correctly and attempt integration of at least one side	B1	$\frac{1}{y} dy = \frac{1-2x^2}{x} dx$
Obtain term $\ln y$	B1	
Obtain terms $\ln x - x^2$	B1	
Use $x = 1, y = 1$ to evaluate a constant, or as limits, in a solution containing at least 2 terms of the form $a \ln y, b \ln x$ and cx^2	M1	The 2 terms of required form must be from correct working e.g. $\ln y = \ln x - x^2 + 1$
Obtain correct solution in any form	A1	
Rearrange and obtain $y = xe^{1-x^2}$	A1	OE
	6	


Question 48

(a)	Separate variables correctly and attempt integration of at least one side	M1
	Obtain term $\ln y$	A1
	Obtain term of the form $\pm \ln(1 - \cos x)$	M1
	Obtain term $\ln(1 - \cos x)$	A1
	Use $x = \pi, y = 4$ to evaluate a constant, or as limits, in a solution containing terms of the form $a \ln y$ and $b \ln(1 - \cos x)$	M1
	Obtain final answer $y = 2(1 - \cos x)$	A1
		6
(b)	Show a correct graph for $0 < x < 2\pi$ with the maximum at $x = \pi$	B1 FT
		1

Question 49

(a)(i)	Justify the given statement $\frac{MN}{y} = \frac{dy}{dx}$	B1
		1
(a)(ii)	Express the area of PMN in terms of y and $\frac{dy}{dx}$ and equate to $\tan x$	M1
	Obtain the given equation correctly	A1
		2

(b)	Separate variables and integrate at least one side	M1
	Obtain term $\frac{1}{6}y^3$	A1
	Obtain term of the form $\pm \ln \cos x$	M1
	Evaluate a constant or use $x = 0$ and $y = 1$ in a solution containing terms ay^3 and $\pm \ln \cos x$, or equivalent	M1
	Obtain correct answer in any form, e.g. $\frac{1}{6}y^3 = -\ln \cos x + \frac{1}{6}$	A1
	Obtain final answer $y = \sqrt[3]{(1 - 6 \ln \cos x)}$	A1
		6



Question 50

State equation $\frac{dy}{dx} = k \frac{y}{\sqrt{x+1}}$	B1
Separate variables correctly for <i>their</i> differential equation and integrate at least one side	*M1
Obtain $\ln y$	A1
Obtain $2[k]\sqrt{x+1}$	A1
Use (0, 1) and (3, e) in an expression containing $\ln y$ and $\sqrt{x+1}$ and a constant of integration to determine k and/or a constant of integration c (or use (0, 1), (3, e) and (x, y) as limits on definite integrals)	DM1
Obtain $k = \frac{1}{2}$ and $c = -1$	A1
Obtain $y = \exp(\sqrt{x+1} - 1)$	A1
	7

Question 51

State a suitable form of partial fractions for $\frac{1}{x^2(1+2x)}$	B1
Use a relevant method to determine a constant	M1
Obtain one of $A = -2$, $B = 1$ and $C = 4$	A1
Obtain a second value	A1
Obtain the third value	A1
Separate variables correctly and integrate at least one term	M1
Obtain terms $-2 \ln x - \frac{1}{x} + 2 \ln(1+2x)$ and t	B3 FT
Evaluate a constant, or use limits $x = 1$, $t = 0$ in a solution containing terms t , $a \ln x$ and $b \ln(1+2x)$, where $ab \neq 0$	M1
Obtain a correct expression for t in any form, e.g. $t = -\frac{1}{x} + 2 \ln\left(\frac{1+2x}{3x}\right) + 1$	A1
	11

Question 52

(a)	State or imply equation of the form $\frac{dx}{dt} = k \frac{x}{20-x}$	M1	
	Obtain $k = 19$	A1	AG
			2
(b)	Separate variables and integrate at least one side	M1	
	Obtain terms $20 \ln x - x$ and $19t$, or equivalent	A1 A1	
	Evaluate a constant or use $t = 0$ and $x = 1$ as limits in a solution containing terms $a \ln x$ and bt	M1	
	Substitute $t = 1$ and rearrange the equation in the given form	A1	AG
			5
(c)	Use $x_{n+1} = e^{0.9+0.05x_n}$ correctly at least once	M1	
	Obtain final answer $x = 2.83$	A1	
	Show sufficient iterations to 4 decimal places to justify 2.83 to 2 d.p. or show there is a sign change in the interval (2.825, 2.835)	A1	
			3
(d)	Set $x = 20$ and obtain answer $t = 2.15$	B1	
			1

Question 53

Separate variables correctly	B1	$\int \frac{1}{y^2} dy = \int 4xe^{-2x} dx$
$\int \frac{1}{y^2} dy = -\frac{1}{y}$	B1	OE
Commence the other integration and reach $axe^{-2x} + b \int e^{-2x} dx$	M1	
Obtain $-2xe^{-2x} + 2 \int e^{-2x} dx$ or $-\frac{1}{2}xe^{-2x} + \frac{1}{2} \int e^{-2x} dx$	A1	SOI (might have taken out factor of 4)
Complete integration and obtain $-2xe^{-2x} - e^{-2x}$	A1	
Evaluate a constant or use $x = 0$ and $y = 1$ as limits in a solution containing terms of the form $\frac{p}{y}$, qxe^{-2x} , re^{-2x} , or equivalent.	M1	
Obtain $y = \frac{e^{2x}}{2x+1}$, or equivalent expression for y	A1	ISW
		7

Question 54

(a)	Show sufficient working to justify the given answer	B1	
		1	
(b)	Correct separation of variables	B1	e.g. $-\int \frac{1}{t} dt = \int \frac{1}{x \ln x} dx$
	Obtain term $\ln(\ln x)$	B1	
	Obtain term $-\ln t$	B1	
	Evaluate a constant or use $x = e$ and $t = 2$ as limits in an expression involving $\ln(\ln x)$	M1	
	Obtain correct solution in any form, e.g. $\ln(\ln x) = -\ln t + \ln 2$	A1	
	Use log laws to enable removal of logarithms	M1	
	Obtain answer $x = e^{\frac{2}{t}}$, or simplified equivalent	A1	
		7	
(c)	State that x tends to 1 coming from $x = e^{\frac{k}{t}}$	B1	
		1	

Question 55

Correctly separate variables and integrate at least one side	M1	To obtain $a \ln y$ or $b \ln(x+1) + c \ln(3x+1)$
Obtain term $\ln y$ from integral of $1/y$	B1	
State or imply the form $\frac{A}{x+1} + \frac{B}{3x+1}$ and use a correct method to find a constant	M1	
Obtain $A = -\frac{1}{2}$ and $B = \frac{3}{2}$	A1	
Obtain terms $-\frac{1}{2} \ln(x+1) + \frac{1}{2} \ln(3x+1)$ or $-\frac{1}{2} \ln(2x+2) + \frac{1}{2} \ln(6x+2)$ or combination of these terms	A1 FT + A1 FT	The FT is on the values of A and B .
Use $x = 1$ and $y = 1$ to evaluate a constant, or expression for a constant, (decimal equivalent of \ln terms allowed) or as limits, in a solution containing terms $a \ln y$, $b \ln(x+1)$ and $c \ln(3x+1)$, where $abc \neq 0$	*M1	e.g. $\ln y = -\frac{1}{2} \ln(x+1) + \frac{1}{2} \ln(3x+1) - \frac{1}{2} \ln 2$
Obtain an expression for y or y^2 and substitute $x = 3$	DM1	Do not accept decimal equivalent of \ln terms
Obtain answer $y = \frac{1}{2}\sqrt{5}$ or $\sqrt{\frac{5}{4}}$ or $\sqrt{\frac{10}{8}}$	A1	ISW. Must be simplified and exact, do not allow 1.118 or $e^{\frac{1}{2} \ln \frac{5}{4}}$.
	9	

Question 56

(a)	Separate variables correctly	B1	$\frac{dN}{N^{\frac{3}{2}}} = (k \cos 0.02t) dt$ Allow without integral signs.
	Obtain term $-\frac{2}{\sqrt{N}}$	B1	OE Ignore position of k .
	Obtain term $50 \sin 0.02t$	B1	OE Ignore position of k .
	Use $t = 0, N = 100$ to evaluate a constant, or as limits, in a solution containing terms $\frac{a}{\sqrt{N}}$ and $b \sin 0.02t$, where $ab \neq 0$	M1	$\left[\text{e.g. } c = -0.2 \text{ or } c = \frac{-0.2}{k} \right]$
	Obtain correct solution in any form, e.g. $-\frac{2}{\sqrt{N}} = 50k \sin 0.02t - 0.2$	A1	OE ISW $\text{e.g. } N = \frac{1}{(25k \sin 0.02t - 0.1)^2} \quad -2N^{\frac{1}{2}} = \frac{k}{0.02} \sin 0.02t - \frac{1}{5}$ $50k \sin 0.02t = -\frac{2}{\sqrt{N}} + \frac{1}{5} \quad \frac{1}{\sqrt{N}} = -\frac{1}{2}k(50 \sin 0.02t) + \frac{1}{10}$ $50 \sin\left(\frac{1}{50}t\right) = -\frac{2\sqrt{N}}{kN} + \frac{20}{100k}$
		5	
(b)	Use the substitution $N = 625$ and $t = 50$ in expression of appropriate form to evaluate k	M1	Expression must contain $a + b k \sin 0.02t \cdot (\sqrt{N})^{\pm n}$, where $n = -1, 1, 3$ or 5 and a and b are constants $ab \neq 0$ or $(a + b k \sin 0.02t)^{\pm 2}$ and $(N)^{\pm n}$. Allow with k replaced by $\frac{1}{k}$, error due to $k(N^{-3/2})$ when separating variables in 8(a) . If invert term by term when 3 terms shown then M0.
	Obtain $k = 0.00285[2148]$	A1	Must evaluate $\sin 1$. Degrees $k = 0.138$ M1 A0.
		2	
(c)	Rearrange and obtain $N = 4(0.2 - 0.142(607) \sin 0.02t)^{-2}$ Substitution for k required	M1	Anything of the form $N = c(d - ek \sin 0.02t)^{-2}$, where c, d and e are constants $cde \neq 0$ and value of k substituted. Allow with k replaced by $1/k$, error due to $k(N^{-3/2})$ when separating variables in 8(a) . OE ISW e.g. $N = \left(\frac{10}{0.7125 \sin 0.02t - 1} \right)^2 \quad N = \frac{1}{(-0.0713 \sin 0.02t + 0.1)^2}$ $N = \frac{100}{\left(\left(\frac{0.6}{\sin 1} \right) \sin 0.02t - 1 \right)^2} \quad N = \frac{1}{\left(\frac{3}{-50 \sin 1} \times \sin 0.02t + \frac{1}{10} \right)^2}$ $N = \left(-\frac{0.06}{\sin 1} \sin 0.02t + 0.1 \right)^{-2} \quad N = \left(\frac{800}{80 - 57 \sin 0.02t} \right)^2$ Do not need to substitute for $\sin(0.02t) = 1$, but must substitute for k .
	Accept answers between 1209 and 1215	A1	ISW Substitute $\sin 0.02t = 1$ or $t = 50 \sin^{-1} 1$ or 78.5 or 25π . Answer with no working (rubric) 0/2. SC $N = \dots$ not seen but correct numerical answer B1 1/2.
		2	

Question 57

(a)	Correct separation of variables	B1	$\int e^{-y} dy = \int xe^{-x} dx$ Condone missing integral signs.
	Obtain term $-e^{-y}$	B1	
	Commence integration by parts and reach $\pm xe^{-x} \pm \int e^{-x} dx$	*M1	M0 if clearly using differentiation of a product.
	Complete integration and obtain $-xe^{-x} - e^{-x}$	A1	
	Use $x = 0$ and $y = 0$ to evaluate a constant or as limits in a solution containing or derived from terms ae^{-y} , bxe^{-x} and ce^{-x} , where $abc \neq 0$	DM1	Must see working for this. In a correct solution they should have $-e^{-y} + C = -xe^{-x} - e^{-x}$ or equivalent. If they take logarithms before finding the constant, the constant must be of the right form.
	Correct solution in any form Must follow from correct working	A1	e.g. $-e^{-y} = -xe^{-x} - e^{-x}$ A0 if constant of integration ignored or assumed to be zero.
	Obtain final answer $y = -\ln((x+1)e^{-x})$ from correct working	A1	OE e.g. $y = x - \ln(x+1)$, $y = \ln\left(\frac{e^x}{x+1}\right)$. A0 if constant of integration ignored or assumed to be zero.
			7
(b)	Obtain answer $(y=)1 - \ln 2$	B1	Must follow from at least 6 or 7 obtained in part 6(a).
		1	

Question 58

	Separate variables correctly	B1	$\int \frac{x}{1+x^2} dx = \int \frac{1}{y} dy$ Accept without integral signs.
	Obtain term $\ln y$	B1	
	State term of the form $k \ln(1+x^2)$	M1	
	State correct term $\frac{1}{2} \ln(1+x^2)$	A1	OE
	Evaluate a constant, or use limits $x = 0$, $y = 2$ in a solution containing terms $a \ln y$ and $b \ln(1+x^2)$ where $ab \neq 0$	M1	If they remove logs first the constant must be of the correct form.
	Obtain correct solution in any form	A1	e.g. $\ln y + \ln \frac{1}{2} = \frac{1}{2} \ln(1+x^2)$
	Simplify and obtain $y = 2\sqrt{1+x^2}$	A1	OE The question asks for simplification, so need to deal with $\exp(\ln(\dots))$.
			7

Question 59

(a)	$a = 30$ and $b = 0.01$	B1	
		1	
(b)	Separate variables and integrate one side	M1	
	Obtain terms $-100\ln(30-0.01V)$ and t , or equivalent	A1 FT + A1 FT	FT <i>their a and b.</i>
	Evaluate a constant, or use $t = 0, V = 0$ as limits, in a solution containing terms $c \ln(30-0.01V)$ and dt where $cd \neq 0$	M1	
	Obtain solution $100\ln 30 - 100\ln(30-0.01V) = t$, or equivalent	A1	
	Substitute $V = 1000$ and obtain answer $t = 40.5$	A1	
		6	
(c)	Obtain $V = 3000(1 - e^{-0.01t})$	B1	OE
	State that V approaches 3000	B1	
		2	

Question 60

(a)	Show sufficient working to justify the given statement	B1	e.g. see $2 \cot \theta \times -\cos \text{csc}^2 \theta$ in the working or express in terms of $\sin \theta$ and $\cos \theta$ and use quotient rule to obtain the given result. Solution must have θ present throughout and must reach the given answer.
		1	
(b)	Separate variables correctly	B1	$\int x dx = \int \frac{\tan^2 \theta}{\sin^2 \theta} - \frac{2 \cot \theta}{\sin^2 \theta} d\theta$ Condone incorrect notation e.g. missing dx. Need either the integral sign or the dx, dθ.
	Check for relevant working in (a)		
	Obtain term $\frac{1}{2}x^2$	B1	
	Obtain terms $\tan \theta + \cot^2 \theta$	B1 + B1	Alternative: $\int \frac{2 \cot \theta}{\sin^2 \theta} d\theta = \int \frac{2 \cos \theta}{\sin^3 \theta} d\theta = -\frac{1}{\sin^2 \theta} (+C)$
	Form an equation for the constant of integration, or use limits $x = 2, \theta = \frac{1}{4}\pi$, in a solution with at least two correctly obtained terms of the form $ax^2, b \tan \theta$ and $c \cot^2 \theta$, where $abc \neq 0$	M1	Need to have 3 terms. Constant of correct form.
	State correct solution in any form, e.g. $\frac{1}{2}x^2 = \tan \theta + \cot^2 \theta$	A1	or $\frac{1}{2}x^2 = \tan \theta + \text{cosec}^2 \theta - 1$ If everything else is correct, allow a correct final answer to imply this A1.
	Substitute $\theta = \frac{1}{6}\pi$ and obtain answer $x = 2.67$	A1	2.6748... $\sqrt{\frac{18+2\sqrt{3}}{3}}$ If see a correctly rounded value ISW.
		7	

Question 61

(a)	Separate variables correctly	B1	$\int \frac{1}{x} dx = \int k e^{-0.1t} dt$
	Obtain term $\ln x$	B1	
	Obtain term $-10k e^{-0.1t}$	B1	Not from $\int x e^{-0.1t} dt$
	Use $x = 20, t = 0$ to evaluate a constant or as limits in a solution containing terms $a \ln x, b e^{-0.1t}$ where $ab \neq 0$	M1	
	Obtain $\ln x = 10k(1 - e^{-0.1t}) + \ln 20$	A1	or equivalent ISW
		5	
(b)	Use $x = 40, t = 10$ to find k or $10k$	M1	Available for their function of the correct structure even if they found no constant in (a).
	Obtain $10k = 1.09654$	A1	or equivalent e.g. $10k = \frac{\ln 2}{1 - e^{-1}}$
	State that x tends to 59.9	A1	Need a number, not an expression for that value 3 sf or better 59.87595.....
		3	

Question 62

	Separate variables correctly and obtain e^{-3y} and $\sin^2 2x$ on the opposite sides	B1	
	Obtain term $-\frac{1}{3} e^{-3y}$	B1	
	Use correct double angle formula for $\sin^2 2x = (1/2)[1 - \cos 4x]$	M1	
	Obtain terms $\frac{1}{2} \left[x - \frac{1}{4} \sin 4x \right]$ oe	A1	
	Use $x = 0, y = 0$ to evaluate a constant or as limits in a solution containing terms of the form ax and $b \sin 4x$ and $ce^{\pm 3y}$	M1	
	Obtain correct answer in any form e.g. $-\frac{1}{3} e^{-3y} = \frac{1}{2} \left[x - \frac{1}{4} \sin 4x \right] - \frac{1}{3}$	A1	
	Substitute $x = \frac{1}{2}$ and obtain $y = 0.175$ or $-\frac{1}{3} \ln \left(\frac{1}{4} + \frac{3}{8} \sin 2 \right)$	A1	OE ISW
		7	

Question 63

Separate the variables correctly	B1	$\frac{y+4}{y^2+4} dy = \frac{1}{x} dx.$
Obtain $\ln x$	B1	
Split the fraction and integrate to obtain $p \ln(y^2+4)$ or $q \tan^{-1} \frac{y}{2}$ correctly	*M1	Only following subdivision into $\frac{y}{y^2+4} + \frac{4}{y^2+4}$. If no subdivision seen then both terms $p \ln(y^2+4)$ and $q \tan^{-1} \frac{y}{2}$ must be present.
Obtain $\frac{1}{2} \ln(y^2+4)$	A1	
Obtain $2 \tan^{-1} \frac{y}{2}$	A1	
Use $(4, 2\sqrt{3})$ in an expression containing at least 2 of $a \ln x$, $b \ln(y^2+4)$ and $c \tan^{-1} \frac{y}{2}$ to obtain constant of integration	DM1	Allow one sign or arithmetic error e.g. $\frac{2\pi}{3}$. May use $(4, 2\sqrt{3})$ and $(x, 2)$ as limits to find x for the final 3 marks.
Correct solution (any form) e.g. $\frac{1}{2} \ln(y^2+4) + 2 \tan^{-1} \frac{y}{2} = \ln x + \frac{2\pi}{3}$ or $\frac{1}{2} \ln(y^2+4) + 2 \tan^{-1} \frac{y}{2} = \ln x + 2 \tan^{-1} \sqrt{3} + \frac{1}{2} \ln 16 - \ln 4$	A1	However solution not asked for so allow $\frac{1}{2} \ln 8 + 2 \tan^{-1} 1 = \ln x + 2 \tan^{-1} \sqrt{3} + \frac{1}{2} \ln 16 - \ln 4.$
Obtain $\sqrt{8} e^{-\frac{1}{6}\pi}$ or 1.68 or more accurate or $2\sqrt{2} e^{-\frac{1}{6}\pi}$ or $\frac{\sqrt{8}}{e^{\frac{1}{6}\pi}}$ or $e^{0.516}$	A1	ISW Must remove \ln so $x = e^{(\ln 2\sqrt{2} - \pi/6)}$ A0.
Alternative method for first *M1 A1 A1		
$p \left((y+4) \tan^{-1} \frac{y}{2} - \int \tan^{-1} \frac{y}{2} dy \right)$	*M1	Allow sign error.
$(y+4) \frac{1}{2} \tan^{-1} \frac{y}{2} - \frac{y}{2} \tan^{-1} \frac{y}{2} + \int \frac{y}{y^2+4} dy$	A1	
Obtain $2 \tan^{-1} \frac{y}{2} + \frac{1}{2} \ln(y^2+4)$	A1	
	8	

Question 64

(a)	Separate variables correctly	B1	$\int \frac{1}{4+9y^2} dy = \int e^{-(2x+1)} dx$. Condone missing integral signs or dx and dy missing.
	Obtain term $-\frac{1}{2}e^{-2x-1}$	B1	OE e.g. $-\frac{1}{2e}e^{-2x}$.
	Obtain term of the form $a \tan^{-1}\left(\frac{3y}{2}\right)$	M1	
	Obtain term $\frac{1}{6} \tan^{-1}\left(\frac{3y}{2}\right)$	A1	OE e.g. $\frac{1}{9} \times \frac{3}{2} \tan^{-1}\frac{3y}{2}$.
	Use $x = 1, y = 0$ to evaluate a constant or as limits in a solution containing or derived from terms of the form $a \tan^{-1}(by)$ and $ce^{\pm(2x+1)}$	M1	If they rearrange before evaluating the constant, the constant must be of the correct form.
	Obtain correct answer in any form	A1	e.g. $\frac{1}{6} \tan^{-1}\frac{3y}{2} = \frac{1}{2}e^{-3} - \frac{1}{2}e^{-(2x+1)}$.
	Obtain final answer $y = \frac{2}{3} \tan(3e^{-3} - 3e^{-2x-1})$	A1	OE Allow with $3e^{-3} = 0.149\dots$
		7	
(b)	State that y approaches $\frac{2}{3} \tan(3e^{-3})$	B1 FT	Or exact equivalent. The FT is on correct work on a solution containing e^{-2x-1} . Condone $y = \dots$. Accept correct answer stated with minimal wording. $0.10032\dots$ is not exact so B0.
		1	

Question 65

Correct separation of variables	B1	$\int \sin^2 3y dy = \int 4 \sec 2x \tan 2x dx$ or equivalent. Condone missing integral signs or dx and dy.
Integrate to obtain $k \sec 2x$	M1	
Obtain $2 \sec 2x$	A1	
Use double angle formula and integrate to obtain $py + q \sin 6y$	M1	Or two cycles of integration by parts.
Obtain $\frac{1}{2}y - \frac{1}{12} \sin 6y$	A1	
Use $y = 0, x = \frac{\pi}{6}$ in a solution containing terms $\lambda \sec 2x$ and $\mu \sin 6y$ to find the constant of integration	M1	
Obtain $\frac{1}{2}y - \frac{1}{12} \sin 6y = 2 \sec 2x - 4$	A1	Or equivalent seen or implied by $\frac{\pi}{2} \left(-\frac{1}{12} \sin \pi \right) = 2 \sec 2x - 4$.
Obtain $x = 0.541$	A1	From correct working (not by using the calculator to integrate).
	8	