

A-Level

Topic : Continuous Random Variable

May 2013-May 2023

Questions

Question 1

A random variable X has probability density function given by

$$f(x) = \begin{cases} \frac{k}{x^3} & x \geq 1, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (i) Show that $k = 2$. [2]
- (ii) Find $P(1 \leq X \leq 2)$. [2]
- (iii) Find $E(X)$. [3]

Question 2

A random variable X has probability density function given by

$$f(x) = \begin{cases} \frac{2}{3}x & 1 \leq x \leq 2, \\ 0 & \text{otherwise.} \end{cases}$$

- (i) Find $E(X)$. [3]
- (ii) Find $P(X < E(X))$. [2]
- (iii) Hence explain whether the mean of X is less than, equal to or greater than the median of X . [2]

Question 3

The time in minutes taken by people to read a certain booklet is modelled by the random variable T with probability density function given by

$$f(t) = \begin{cases} \frac{1}{2\sqrt{t}} & 4 \leq t \leq 9, \\ 0 & \text{otherwise.} \end{cases}$$

- (i) Find the time within which 90% of people finish reading the booklet. [3]
- (ii) Find $E(T)$ and $\text{Var}(T)$. [6]

Question 4

The waiting time, T weeks, for a particular operation at a hospital has probability density function given by

$$f(t) = \begin{cases} \frac{1}{2500}(100t - t^3) & 0 \leq t \leq 10, \\ 0 & \text{otherwise.} \end{cases}$$

- (i) Given that $E(T) = \frac{16}{3}$, find $\text{Var}(T)$. [3]
- (ii) 10% of patients have to wait more than n weeks for their operation. Find the value of n , giving your answer correct to the nearest integer. [5]

Question 5

The volume, in cm^3 , of liquid left in a glass by people when they have finished drinking all they want is modelled by the random variable X with probability density function given by

$$f(x) = \begin{cases} k(x-2)^2 & 0 \leq x \leq 2, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (i) Show that $k = \frac{3}{8}$. [2]
- (ii) 20% of people leave at least $d \text{ cm}^3$ of liquid in a glass. Find d . [3]
- (iii) Find $E(X)$. [3]

Question 6


The lifetime, X years, of a certain type of battery has probability density function given by

$$f(x) = \begin{cases} \frac{k}{x^2} & 1 \leq x \leq a, \\ 0 & \text{otherwise,} \end{cases}$$

where k and a are positive constants.

- (i) State what the value of a represents in this context. [1]
- (ii) Show that $k = \frac{a}{a-1}$. [3]
- (iii) Experience has shown that the longest that any battery of this type lasts is 2.5 years. Find the mean lifetime of batteries of this type. [3]

Question 7



A random variable X takes values between 0 and 4 only and has probability density function as shown in the diagram. Calculate the median of X . [3]

Question 8

The time, T hours, spent by people on a visit to a museum has probability density function

$$f(t) = \begin{cases} kt(16 - t^2) & 0 \leq t \leq 4, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (i) Show that $k = \frac{1}{64}$. [3]
- (ii) Calculate the probability that two randomly chosen people each spend less than 1 hour on a visit to the museum. [4]
- (iii) Find the mean time spent on a visit to the museum. [3]

Question 9


A random variable X has probability density function given by

$$f(x) = \begin{cases} \frac{k}{x} & 1 \leq x \leq a, \\ 0 & \text{otherwise,} \end{cases}$$

where k and a are positive constants.

- (i) Show that $k = \frac{1}{\ln a}$. [3]
- (ii) Find $E(X)$ in terms of a . [3]
- (iii) Find the median of X in terms of a . [4]

Question 10



The diagram shows the graph of the probability density function, f , of a random variable X .

- (i) Find the value of the constant c . [2]
- (ii) Find the value of a such that $P(a < X < 1) = 0.1$. [4]
- (iii) Find $E(X)$. [2]


Question 11

- (a) The time for which Lucy has to wait at a certain traffic light each day is T minutes, where T has probability density function given by

$$f(t) = \begin{cases} \frac{3}{2}t - \frac{3}{4}t^2 & 0 \leq t \leq 2, \\ 0 & \text{otherwise.} \end{cases}$$

Find the probability that, on a randomly chosen day, Lucy has to wait for less than half a minute at the traffic light. [3]

- (b)



The diagram shows the graph of the probability density function, g , of a random variable X . The graph of g is a semicircle with centre $(0, 0)$ and radius a . Elsewhere $g(x) = 0$.

- (i) Find the value of a . [2]
 (ii) State the value of $E(X)$. [1]
 (iii) Given that $P(X < -c) = 0.2$, find $P(X < c)$. [2]

Question 12

The probability density function of the random variable X is given by

$$f(x) = \begin{cases} \frac{3}{4}x(c-x) & 0 \leq x \leq c, \\ 0 & \text{otherwise,} \end{cases}$$

where c is a constant.

- (i) Show that $c = 2$. [3]
 (ii) Sketch the graph of $y = f(x)$ and state the median of X . [3]
 (iii) Find $P(X < 1.5)$. [4]
 (iv) Hence write down the value of $P(0.5 < X < 1)$. [1]

Question 13


The waiting time, T minutes, for patients at a doctor's surgery has probability density function given by

$$f(t) = \begin{cases} k(225 - t^2) & 0 \leq t \leq 15, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (i) Show that $k = \frac{1}{2250}$. [3]
 (ii) Find the probability that a patient has to wait for more than 10 minutes. [3]
 (iii) Find the mean waiting time. [4]

Question 14



The random variable X has probability density function, f , as shown in the diagram, where a is a constant. Find the value of a and hence show that $E(X) = 0.943$ correct to 3 significant figures. [5]

Question 15

A random variable X has probability density function given by

$$f(x) = \begin{cases} k(4 - x^2) & -2 \leq x \leq 2, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (i) Show that $k = \frac{3}{32}$. [3]
- (ii) Sketch the graph of $y = f(x)$ and hence write down the value of $E(X)$. [2]
- (iii) Find $P(X < 1)$. [3]

Question 16

A random variable X has probability density function given by


$$f(x) = \begin{cases} k(3 - x) & 1 \leq x \leq 2, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (i) Show that $k = \frac{2}{3}$. [3]
- (ii) Find the median of X . [4]

Question 17

(a)



The diagram shows the graph of the probability density function, f , of a random variable X , where

$$f(x) = \begin{cases} \frac{2}{9}(3x - x^2) & 0 \leq x \leq 3, \\ 0 & \text{otherwise.} \end{cases}$$

- (i) State the value of $E(X)$ and find $\text{Var}(X)$. [4]
- (ii) State the value of $P(1.5 \leq X \leq 4)$. [1]
- (iii) Given that $P(1 \leq X \leq 2) = \frac{13}{27}$, find $P(X > 2)$. [2]

(b) A random variable, W , has probability density function given by

$$g(w) = \begin{cases} aw & 0 \leq w \leq b, \\ 0 & \text{otherwise,} \end{cases}$$

where a and b are constants. Given that the median of W is 2, find a and b . [4]

Question 18

The time, T minutes, taken by people to complete a test has probability density function given by


$$f(t) = \begin{cases} k(10t - t^2) & 5 \leq t \leq 10, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (i) Show that $k = \frac{3}{250}$. [3]
- (ii) Find $E(T)$. [3]
- (iii) Find the probability that a randomly chosen value of T lies between $E(T)$ and the median of T . [3]
- (iv) State the greatest possible length of time taken to complete the test. [1]

Question 19

(a)



The diagram shows the graph of the probability density function of a variable X . Given that the graph is symmetrical about the line $x = 1$ and that $P(0 < X < 2) = 0.6$, find $P(X > 0)$. [2]

(b) A flower seller wishes to model the length of time that tulips last when placed in a jug of water. She proposes a model using the random variable X (in hundreds of hours) with probability density function given by

$$f(x) = \begin{cases} k(2.25 - x^2) & 0 \leq x \leq 1.5, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

(i) Show that $k = \frac{4}{9}$. [3]

(ii) Use this model to find the mean number of hours that a tulip lasts in a jug of water. [4]

The flower seller wishes to create a similar model for daffodils. She places a large number of daffodils in jugs of water and the longest time that any daffodil lasts is found to be 290 hours.

(iii) Give a reason why $f(x)$ would not be a suitable model for daffodils. [1]

(iv) The flower seller considers a model for daffodils of the form

$$g(x) = \begin{cases} c(a^2 - x^2) & 0 \leq x \leq a, \\ 0 & \text{otherwise,} \end{cases}$$

where a and c are constants. State a suitable value for a . (There is no need to evaluate c .)

[1]

Question 20

In each turn of a game, a coin is pushed and slides across a table. The distance, X metres, travelled by the coin has probability density function given by

$$f(x) = \begin{cases} kx^2(2 - x) & 0 \leq x \leq 2, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.





(i) State the greatest possible distance travelled by the coin in one turn. [1]

(ii) Show that $k = \frac{3}{4}$. [3]

(iii) Find the mean distance travelled by the coin in one turn. [3]

(iv) Out of 400 turns, find the expected number of turns in which the distance travelled by the coin is less than 1 metre. [3]

Question 21



The diagrams show the probability density functions of four random variables W , X , Y and Z . Each of the four variables takes values between -3 and 3 only, and their standard deviations are σ_W , σ_X , σ_Y and σ_Z respectively.

(i) List σ_W , σ_X , σ_Y and σ_Z in order of size, starting with the largest. [2]

(ii) The probability density function of X is given by




$$f(x) = \begin{cases} \frac{1}{18}x^2 & -3 \leq x \leq 3, \\ 0 & \text{otherwise.} \end{cases}$$

(a) Show that $\sigma_X = 2.32$ correct to 3 significant figures. [3]

(b) Calculate $P(X > \sigma_X)$. [3]

(c) Write down the value of $P(X > 2\sigma_X)$. [1]

Question 22



The diagrams show the probability density functions of four random variables W , X , Y and Z . Each of the four variables takes values between 0 and 3 only, and their medians are m_W , m_X , m_Y and m_Z respectively.

- (i) List m_W , m_X , m_Y and m_Z in order of size, starting with the largest. [2]

- (ii) The probability density function of X is given by

$$f(x) = \begin{cases} \frac{4}{81}x^3 & 0 \leq x \leq 3, \\ 0 & \text{otherwise.} \end{cases}$$


- (a) Show that $E(X) = \frac{12}{5}$. [3]

- (b) Calculate $P(X > E(X))$. [3]

- (c) Write down the value of $P(X < 2E(X))$. [1]

Question 23

(a)



The diagram shows the graphs of three functions, f_1 , f_2 and f_3 . The function f_1 is a probability density function.

- (i) State the value of k . [1]
- (ii) For each of the functions f_2 and f_3 , state why it cannot be a probability density function. [2]
- (b) The probability density function g is defined by

$$g(x) = \begin{cases} 6(a^2 - x^2) & -a \leq x \leq a, \\ 0 & \text{otherwise,} \end{cases}$$

where a is a constant.

- (i) Show that $a = \frac{1}{2}$. [3]
- (ii) State the value of $E(X)$. [1]
- (iii) Find $\text{Var}(X)$. [2]

Question 24




The diagram shows the graph of the probability density function, f , of a continuous random variable X , where f is defined by

$$f(x) = \begin{cases} k(x - x^2) & 0 \leq x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

- (i) Show that the value of the constant k is 6. [3]
- (ii) State the value of $E(X)$ and find $\text{Var}(X)$. [4]
- (iii) Find $P(0.4 < X < 2)$. [3]

Question 25



The diagram shows the graph of the probability density function, f , of a random variable X which takes values between 0 and a only. It is given that $P(X < 1) = 0.25$.

(i) Find, in any order,


- (a) $P(X < 2)$,
- (b) the value of a ,
- (c) $f(x)$.

[5]

(ii) Find the median of X .

[3]

Question 26



The time, X minutes, taken by a large number of runners to complete a certain race has probability density function f given by

$$f(x) = \begin{cases} \frac{k}{x^2} & 5 \leq x \leq 10, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant, as shown in the diagram.

(i) Without calculation, explain how you can tell that there were more runners whose times were below 7.5 minutes than above 7.5 minutes. [1]

(ii) Show that $k = 10$. [3]

(iii) Find $E(X)$. [3]

(iv) Find $\text{Var}(X)$. [2]

Question 27

A continuous random variable, X , has probability density function given by

$$f(x) = \begin{cases} \frac{1}{4}(x+1) & 0 \leq x \leq 2, \\ 0 & \text{otherwise.} \end{cases}$$

- (i) Find $E(X)$. [3]
(ii) Find the median of X . [3]

Question 28

The random variable X has probability density function given by

$$f(x) = \begin{cases} \frac{k}{\sqrt{x}} & 0 < x \leq a, \\ 0 & \text{otherwise,} \end{cases}$$

where k and a are constants. It is given that $E(X) = 3$.

- (i) Find the value of a and show that $k = \frac{1}{6}$. [7]
(ii) Find the median of X . [3]

Question 29

A random variable X has probability density function given by

$$f(x) = \begin{cases} 6x(1-x) & 0 \leq x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

- (i) Find the probability that X does not lie between 0.3 and 0.7. [4]
(ii) Sketch the graph of the probability density function and hence state the value of $E(X)$. [2]
(iii) Find $\text{Var}(X)$. [3]

Question 30


A random variable X has probability density function defined by

$$f(x) = \begin{cases} k \left(\frac{1}{x^2} + \frac{1}{x^3} \right) & 1 \leq x \leq 2, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (i) Show that $k = \frac{8}{7}$. [3]
(ii) Find $E(X)$. [3]
(iii) Three values of X are chosen at random. Find the probability that one of these values is less than 1.5 and the other two are greater than 1.5. [5]

Question 31



Question 32

The time, in minutes, taken by people to complete a test is modelled by the continuous random variable X with probability density function given by

$$f(x) = \begin{cases} \frac{k}{x^2} & 5 \leq x \leq 10, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (i) Show that $k = 10$. [3]
- (ii) Show that $E(X) = 10 \ln 2$. [2]
- (iii) Find $P(X > 9)$. [3]
- (iv) Given that $P(X < a) = 0.6$, find a . [3]

Question 33

The random variable X has probability density function given by

$$f(x) = \begin{cases} kx^{-1} & 2 \leq x \leq 6, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (i) Show that $k = \frac{1}{\ln 3}$. [2]
- (ii) Show that $E(X) = 3.64$, correct to 3 significant figures. [3]
- (iii) Given that the median of X is m , find $P(m < X < E(X))$. [4]

Question 34


The time, X hours, taken by a large number of runners to complete a race is modelled by the probability density function given by

$$f(x) = \begin{cases} \frac{k}{(x+1)^2} & 0 \leq x \leq a, \\ 0 & \text{otherwise,} \end{cases}$$

where k and a are constants.

- (i) Show that $k = \frac{a+1}{a}$. [3]
 - (ii) State what the constant a represents in this context. [1]
- Three quarters of the runners take half an hour or less to complete the race.
- (iii) Find the value of a . [3]

Question 35



A random variable X has probability density function given by

$$f(x) = \begin{cases} (\sqrt{2}) \cos x & 0 \leq x \leq \frac{1}{4}\pi, \\ 0 & \text{otherwise,} \end{cases}$$

as shown in the diagram.

- (i) Find $P(X > \frac{1}{6}\pi)$. [2]
- (ii) Find the median of X . [4]
- (iii) Find $E(X)$. [4]

Question 36

A function f is defined by

$$f(x) = \begin{cases} \frac{3x^2}{a^3} & 0 \leq x \leq a, \\ 0 & \text{otherwise,} \end{cases}$$

where a is a constant.

- (i) Show that f is a probability density function for all positive values of a . [3]
- The random variable X has probability density function f and the median of X is 2.
- (ii) Show that $a = 2.52$, correct to 3 significant figures. [3]
 - (iii) Find $E(X)$. [3]

Question 37

X is a random variable with probability density function given by


$$f(x) = \begin{cases} \frac{a}{x^2} & 1 \leq x \leq b, \\ 0 & \text{otherwise,} \end{cases}$$

where a and b are constants.

- (i) Show that $b = \frac{a}{a-1}$. [3]
- (ii) Given that the median of X is $\frac{3}{2}$, find the values of a and b . [3]
- (iii) Use your values of a and b from part (ii) to find $E(X)$. [3]

Question 38

(a)



The diagram shows the graph of the probability density function, f , of a random variable X , where a is a constant greater than 0.5. The graph between $x = 0$ and $x = a$ is a straight line parallel to the x -axis.

(i) Find $P(X < 0.5)$ in terms of a . [2]

(ii) Find $E(X)$ in terms of a . [1]

(iii) Show that $\text{Var}(X) = \frac{1}{12}a^2$. [2]

(b) A random variable T has probability density function given by

$$g(t) = \begin{cases} \frac{3}{2(t-1)^2} & 2 \leq t \leq 4, \\ 0 & \text{otherwise.} \end{cases}$$

Find the value of b such that $P(T \leq b) = \frac{3}{4}$. [4]

Question 39

A random variable X has probability density function given by

$$f(x) = \begin{cases} k(3x - x^2) & 0 \leq x \leq 3, \\ 0 & \text{otherwise.} \end{cases}$$

(i) Show that $k = \frac{2}{9}$. [3]

(ii) Find $P(1 \leq X \leq 2)$. [2]

(iii) Find $\text{Var}(X)$. [5]

Question 40

A random variable X has probability density function given by

$$f(x) = \begin{cases} \frac{1}{2}x & 0 \leq x \leq a, \\ 0 & \text{otherwise,} \end{cases}$$

where a is a constant.

(i) Find a . [2]

(ii) Show that $E(X) = \frac{4}{3}$. [3]

The median of X is denoted by m .

(iii) Find $P(E(X) < X < m)$. [4]

Question 41

Bottles of Lanta contain approximately 300 ml of juice. The volume of juice, in millilitres, in a bottle is $300 + X$, where X is a random variable with probability density function given by

$$f(x) = \begin{cases} \frac{3}{4000}(100 - x^2) & -10 \leq x \leq 10, \\ 0 & \text{otherwise.} \end{cases}$$

- (a) Find the probability that a randomly chosen bottle of Lanta contains more than 305 ml of juice. [3]
- (b) Given that 25% of bottles of Lanta contain more than $(300 + p)$ ml of juice, show that

$$p^3 - 300p + 1000 = 0. \quad [4]$$

Question 42

The length, X centimetres, of worms of a certain type is modelled by the probability density function

$$f(x) = \begin{cases} \frac{6}{125}(10 - x)(x - 5) & 5 \leq x \leq 10, \\ 0 & \text{otherwise.} \end{cases}$$

- (a) State the value of $E(X)$. [1]
- (b) Find $\text{Var}(X)$. [3]
- (c) Two worms of this type are chosen at random.

Find the probability that exactly one of them has length less than 6 cm. [5]

Question 43

A random variable X has probability density function given by

$$f(x) = \begin{cases} \frac{k}{x^2} & 1 \leq x \leq a, \\ 0 & \text{otherwise,} \end{cases}$$

where k and a are positive constants.

- (a) Show that $k = \frac{a}{a-1}$. [3]
- (b) Find $E(X)$ in terms of a . [3]
- (c) Find the 60th percentile of X in terms of a . [4]

Question 44

The length of time, T minutes, that a passenger has to wait for a bus at a certain bus stop is modelled by the probability density function given by


$$f(t) = \begin{cases} \frac{3}{4000}(20t - t^2) & 0 \leq t \leq 20, \\ 0 & \text{otherwise.} \end{cases}$$

- (a) Sketch the graph of $y = f(t)$. [1]
- (b) Hence explain, without calculation, why $E(T) = 10$. [1]
- (c) Find $\text{Var}(T)$. [3]
- (d) It is given that $P(T < 10 + a) = p$, where $0 < a < 10$.

Find $P(10 - a < T < 10 + a)$ in terms of p . [2]

- (e) Find $P(8 < T < 12)$. [3]


Question 45



The diagram shows the probability density function, $f(x)$, of a random variable X . For $0 \leq x \leq a$, $f(x) = k$; elsewhere $f(x) = 0$.

- (a) Express k in terms of a . [1]
 (b) Given that $\text{Var}(X) = 3$, find a . [4]


Question 46



A random variable X takes values between 0 and 3 only and has probability density function as shown in the diagram, where c is a constant.

- (a) Show that $c = \frac{2}{3}$. [1]
 (b) Find $P(X > 2)$. [2]
 (c) Calculate $E(X)$. [4]

Question 47



The diagram shows the graph of the probability density function, f , of a random variable X .

- (a) Find the value of the constant k . [2]
 (b) Using this value of k , find $f(x)$ for $0 \leq x \leq k$ and hence find $E(X)$. [3]
 (c) Find the value of p such that $P(p < X < 1) = 0.25$. [4]

Question 48

Alethia models the length of time, in minutes, by which her train is late on any day by the random variable X with probability density function given by

$$f(x) = \begin{cases} \frac{3}{8000}(x-20)^2 & 0 \leq x \leq 20, \\ 0 & \text{otherwise.} \end{cases}$$

(a) Find the probability that the train is more than 10 minutes late on each of two randomly chosen days. [4]


(b) Find $E(X)$. [4]

(c) The median of X is denoted by m .

Show that m satisfies the equation $(m-20)^3 = -4000$, and hence find m correct to 3 significant figures. [4]

(d) State one way in which Alethia's model may be unrealistic. [1]

Question 49



The random variable X takes values in the range $1 \leq x \leq p$, where p is a constant. The graph of the probability density function of X is shown in the diagram.

(a) Show that $p = 2$. [2]

(b) Find $E(X)$. [5]

Question 50

The probability density function, f , of a random variable X is given by

$$f(x) = \begin{cases} k(6x - x^2) & 0 \leq x \leq 6, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

State the value of $E(X)$ and show that $\text{Var}(X) = \frac{9}{5}$. [6]

Question 51

The graph of the probability density function of a random variable X is symmetrical about the line $x = 4$.

Given that $P(X < 5) = \frac{20}{27}$, find $P(3 < X < 5)$. [2]

Question 52

A random variable X has probability density function given by

$$f(x) = \begin{cases} \frac{1}{18}(9 - x^2) & 0 \leq x \leq 3, \\ 0 & \text{otherwise.} \end{cases}$$

(a) Find $P(X < 1.2)$. [3]

(b) Find $E(X)$. [3]

The median of X is m .

(c) Show that $m^3 - 27m + 27 = 0$. [3]

Question 53

(a) The probability density function of the random variable X is given by

$$f(x) = \begin{cases} kx(4 - x) & 0 \leq x \leq 2, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

(i) Show that $k = \frac{3}{16}$. [3]

(ii) Find $E(X)$. [3]

(b) The random variable Y has the following properties.

- Y takes values between 0 and 5 only.
- The probability density function of Y is symmetrical.

Given that $P(Y < a) = 0.2$, find $P(2.5 < Y < 5 - a)$ illustrating your method with a sketch on the axes provided. [3]



Question 54

A random variable X has probability density function given by

$$f(x) = \begin{cases} \frac{1}{18}(9 - x^2) & 0 \leq x \leq 3, \\ 0 & \text{otherwise.} \end{cases}$$

(a) Find $P(X < 1.2)$. [3]

(b) Find $E(X)$. [3]

The median of X is m .

(c) Show that $m^3 - 27m + 27 = 0$. [3]

Question 55

In a game a ball is rolled down a slope and along a track until it stops. The distance, in metres, travelled by the ball is modelled by the random variable X with probability density function

$$f(x) = \begin{cases} -k(x-1)(x-3) & 1 \leq x \leq 3, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

(a) Without calculation, explain why $E(X) = 2$. [1]

(b) Show that $k = \frac{3}{4}$. [3]

(c) Find $\text{Var}(X)$. [3]

One turn consists of rolling the ball 3 times and noting the largest value of X obtained. If this largest value is greater than 2.5, the player scores a point.

(d) Find the probability that on a particular turn the player scores a point. [4]

Question 56

The random variables X and W have probability density functions f and g defined as follows:

$$f(x) = \begin{cases} p(a^2 - x^2) & 0 \leq x \leq a, \\ 0 & \text{otherwise,} \end{cases}$$
$$g(w) = \begin{cases} q(a^2 - w^2) & -a \leq w \leq a, \\ 0 & \text{otherwise,} \end{cases}$$

where a , p and q are constants.

(a) (i) Write down the value of $P(X \geq 0)$. [1]

(ii) Write down the value of $P(W \geq 0)$. [1]

(iii) Write down an expression for q in terms of p only. [1]

(b) Given that $E(X) = 3$, find the value of a . [6]

Question 57

A random variable X has probability density function f . The graph of $f(x)$ is a straight line segment parallel to the x -axis from $x = 0$ to $x = a$, where a is a positive constant.


(a) State, in terms of a , the median of X . [1]

(b) Find $P(X > \frac{3}{4}a)$. [1]

(c) Show that $\text{Var}(X) = \frac{1}{12}a^2$. [5]

(d) Given that $P(X < b) = p$, where $0 < b < \frac{1}{2}a$, find $P(\frac{1}{3}b < X < a - \frac{1}{3}b)$ in terms of p . [2]


Question 58



The diagram shows the graph of the probability density function of a random variable X that takes values between -1 and 3 only. It is given that the graph is symmetrical about the line $x = 1$. Between $x = -1$ and $x = 3$ the graph is a quadratic curve.


The random variable S is such that $E(S) = 2 \times E(X)$ and $\text{Var}(S) = \text{Var}(X)$.

- (a) On the grid below, sketch a quadratic graph for the probability density function of S . [1]



The random variable T is such that $E(T) = E(X)$ and $\text{Var}(T) = \frac{1}{4} \text{Var}(X)$.

- (b) On the grid below, sketch a quadratic graph for the probability density function of T . [2]




It is now given that

$$f(x) = \begin{cases} \frac{3}{32}(3 + 2x - x^2) & -1 \leq x \leq 3, \\ 0 & \text{otherwise.} \end{cases}$$

- (c) Given that $P(1 - a < X < 1 + a) = 0.5$, show that $a^3 - 12a + 8 = 0$. [3]

- (d) Hence verify that $0.69 < a < 0.70$. [1]

Question 59



The diagram shows the graph of the probability density function, f , of a random variable X which takes values between -3 and 2 only.

- (a) Given that the graph is symmetrical about the line $x = -0.5$ and that $P(X < 0) = p$, find $P(-1 < X < 0)$ in terms of p . [2]


- (b) It is now given that the probability density function shown in the diagram is given by

$$f(x) = \begin{cases} a - b(x^2 + x) & -3 \leq x \leq 2, \\ 0 & \text{otherwise,} \end{cases}$$

where a and b are positive constants.

- (i) Show that $30a - 55b = 6$. [3]
- (ii) By substituting a suitable value of x into $f(x)$, find another equation relating a and b and hence determine the values of a and b . [3]


Question 60



The diagram shows the graph of the probability density function of a random variable X that takes values between -1 and 3 only. It is given that the graph is symmetrical about the line $x = 1$. Between $x = -1$ and $x = 3$ the graph is a quadratic curve.


The random variable S is such that $E(S) = 2 \times E(X)$ and $\text{Var}(S) = \text{Var}(X)$.

- (a) On the grid below, sketch a quadratic graph for the probability density function of S . [1]



The random variable T is such that $E(T) = E(X)$ and $\text{Var}(T) = \frac{1}{4} \text{Var}(X)$.

- (b) On the grid below, sketch a quadratic graph for the probability density function of T . [2]




It is now given that

$$f(x) = \begin{cases} \frac{3}{32}(3 + 2x - x^2) & -1 \leq x \leq 3, \\ 0 & \text{otherwise.} \end{cases}$$

- (c) Given that $P(1 - a < X < 1 + a) = 0.5$, show that $a^3 - 12a + 8 = 0$. [3]

- (d) Hence verify that $0.69 < a < 0.70$. [1]

Question 61



The diagram shows the graph of the probability density function, f , of a random variable X that takes values between $x = 0$ and $x = 3$ only. The graph is symmetrical about the line $x = 1.5$.

- (a) It is given that $P(X < 0.6) = a$ and $P(0.6 < X < 1.2) = b$.

Find $P(0.6 < X < 1.8)$ in terms of a and b . [2]

- (b) It is now given that the equation of the probability density function of X is


$$f(x) = \begin{cases} kx^2(3-x)^2 & 0 \leq x \leq 3, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

- (i) Show that $k = \frac{10}{81}$. [3]

- (ii) Find $\text{Var}(X)$. [3]

Question 62



A random variable X has probability density function f , where the graph of $y = f(x)$ is a semicircle with centre $(0, 0)$ and radius $\sqrt{\frac{2}{\pi}}$, entirely above the x -axis. Elsewhere $f(x) = 0$ (see diagram).

(a) Verify that f can be a probability density function. [2]

A and B are the points where the line $x = \sqrt{\frac{1}{\pi}}$ meets the x -axis and the semicircle respectively.

(b) Show that angle AOB is $\frac{1}{4}\pi$ radians and hence find $P\left(X > \sqrt{\frac{1}{\pi}}\right)$. [6]

Question 63

A random variable X has probability density function f , where


$$f(x) = \begin{cases} \frac{3}{2}(1 - x^2) & 0 \leq x \leq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Find $E(X)$.

[3]

Question 64

(a)




The diagram shows the graph of the probability density function, f , of a random variable X which takes values between 0 and 4 only. Between these two values the graph is a straight line.

(i) Show that $f(x) = kx$ for $0 \leq x \leq 4$, where k is a constant to be determined. [2]

(ii) Hence, or otherwise, find $E(X)$. [3]

(b)




The diagram shows the graph of the probability density function, g , of a random variable W which takes values between 0 and a only, where $a > 0$. Between these two values the graph is a straight line.

Given that the median of W is 1, find the value of a . [3]

Question 65

(a)




The graph of the function f is a straight line segment from $(0, 0)$ to $(2, 1)$.

Show that f could be a probability density function.

[2]

(b)




The graph of the function g is a semicircle, centre $(0, 0)$, entirely above the x -axis.

Given that g is a probability density function, find the radius of the semicircle.

[2]

(c)



The time, X minutes, taken by a large number of students to complete a test has probability density function h , as shown in the diagram.

- (i) Without calculation, use the diagram to explain how you can tell that the median time is less than 15 minutes. [1]

It is now given that

$$h(x) = \begin{cases} \frac{40}{x^2} - \frac{1}{10} & 10 \leq x \leq 20, \\ 0 & \text{otherwise.} \end{cases}$$

- (ii) Find the mean time. [3]