

A-Level

Topic : Discrete , Continuous Hypothesis and Types of Error

May 2013–May 2023

Answers

Question 1

Assume shots independent OR prob of scoring constant	B1	In context
$H_0: P(\text{score}) = 0.82$ $H_1: P(\text{score}) > 0.82$	B1	Both. Allow 'p'
$20 \times 0.82^{19} \times 0.18 + 0.82^{20}$ $= 0.102$ (3 sf)	M1	For use of Bin(20,0.82) and either P(19)
No evidence that improved	A1	and/or P(20) attempted
	B1f	Valid comparison seen (with 0.05 if H_1 $p \neq$
	5	0.82) and correct conclusion ft numerical errors in 0.102 only
		Normal approx'n: B1 B1 ($\mu = 16.4$ acceptable here) if earned, then:
		$CR = 1.222$ (from $\frac{18.5 - 20 \times 0.82}{\sqrt{20 \times 0.82 \times (1 - 0.82)}}$,
		need cc)
		comp $z = 1.282$
		No evidence that improved SC 1
		Same scheme for proportions
[Total: 5]		

Question 2

<p>(i) Assume sd unchanged or 4500</p> <p>H_0: Pop mean = 34600 H_1: Pop mean > 34600</p> $\frac{35400 - 34600}{\frac{4500}{\sqrt{90}}}$ <p>= 1.687/1.686 (1.69) cf 1.645 < 1.686 Evidence that mean wkly profit has increased</p>	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1 M1</p> <p>A1 f 6</p>	<p>Both. Allow just μ, but not just "mean"</p> <p>Allow without $\sqrt{90}$</p> <p>Valid comparison (or 0.0458/0.0459 < 0.05 or 35380 < 35400 or 34600 < 34620) If $H_1: \neq$, and 1.96 used, max B1B0M1A1M1A1f No contradictions</p> <p>Allow not Normal</p> <p>B1* dep 2</p> <p>B1 1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>M1 A1 6</p> <p>Attempt to find cv must see (+) 1.645 allow without $\sqrt{90}$. If found in (i) award when used</p> <p>Standardising with their "CV" must use $\sqrt{90}$</p> <p>Correct tail</p>
<p>(ii) Distr'n of X unknown.</p> <p>Yes</p>	<p>B1*</p> <p>B1* dep 2</p>	<p>Allow not Normal</p>
<p>(iii) 0.05 or 5%</p>	<p>B1 1</p>	
<p>(iv) $\frac{a - 34600}{\frac{4500}{\sqrt{90}}} = 1.645$ $a = 35380$ $\frac{35380 - 36500}{\frac{4500}{\sqrt{90}}} (= -2.361)$ $1 - \Phi(2.361)$ $= 0.0091$</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>M1 A1 6</p>	<p>Attempt to find cv must see (+) 1.645 allow without $\sqrt{90}$. If found in (i) award when used</p> <p>Standardising with their "CV" must use $\sqrt{90}$</p> <p>Correct tail</p>
<p>[Total: 14]</p>		

Question 3

<p>(i) $\frac{73.1 - 75.2}{\frac{5.7}{\sqrt{n}}} = -1.563$ $n = \{-1.563 \times 5.7 \div (-2.1)\}^2$ $n = 18$ Assume s.d. for the region is 5.7</p>	<p>M1</p> <p>A1</p> <p>A1 B1 [4]</p>	<p>For standardising (with \sqrt{n})</p> <p>Any correct expression for n or \sqrt{n}. May be implied by ans.</p>
<p>(ii) H_0: pop mean (or μ) = 75.2 H_0: pop mean (or μ) < 75.2 1.563 comp 1.555 Evidence that plants shorter</p>	<p>B1</p> <p>M1</p> <p>A1 [3]</p>	<p>Both (could be stated in (i))</p> <p>For comparison of z values / areas / x values CWO. No contradictions</p>
<p>[Total: 7]</p>		

Question 4

(i)	H_0 : Pop mean (or λ or μ) is 5.3 H_1 : Pop mean (or λ or μ) is less than 5.3 $P(X \leq 1) = e^{-5.3}(1 + 5.3)$ $P(X \leq 2) = e^{-5.3}(1 + 5.3 + \frac{5.3^2}{2})/P(X=2)$ $P(X \leq 1) = 0.0314$ or 0.0315 & $P(X \leq 2) = 0.102/P(X=2)=0.7071$ CR is 0 or 1 cases No evidence mean has decreased	B1 M1 A1 A1 B1f [5]	Both Both attempted Both correct Dep. M1 and $P(X \leq 1) < 0.05 < P(X \leq 2)$ ft their CR
(ii)	Concluding mean has decreased when it hasn't '0.0314 or 0.0315'	B1 B1f[2]	In context ft their $P(X \leq 1)$, dep. < 0.05
(iii)	$(Po(18.4))$ $N(18.4, 18.4)$ $\frac{20.5-18.4}{\sqrt{18.4}} \quad (= 0.490)$ $1 - \Phi(0.490)$ $= 0.312$ (3 s.f.)	B1 B1f M1 M1 A1 [5]	Stated or implied B1 for $N(18.4, ..)$; B1f for var. = 18.4 For standardising with or without cc.Allow without $\sqrt{\quad}$ Use of tables and attempt to find area consistent with their working

[Total: 12]

Question 5

(i)	Conclude die is biased when it isn't oe ${}^5C_3 \left(\frac{1}{6}\right)^3 \left(\frac{5}{6}\right)^2 + 5 \left(\frac{1}{6}\right)^4 \left(\frac{5}{6}\right) + \left(\frac{1}{6}\right)^5 + 5$ $= \frac{23}{648}$ or 0.0355 (3 sf)	B1 M1 A1 [3]	In context or $1 - \left({}^5C_2 \left(\frac{1}{6}\right)^2 \left(\frac{5}{6}\right) + 5 \left(\frac{1}{6}\right) \left(\frac{5}{6}\right)^4 + \left(\frac{5}{6}\right)^5 \right)$ allow 1 end error
(ii)	State or attempt $P(0, 1, 2)$ with $p = \frac{2}{3}$ ${}^5C_2 \left(\frac{2}{3}\right)^2 \left(\frac{1}{3}\right)^3 + 5 \left(\frac{2}{3}\right) \left(\frac{1}{3}\right)^4 + \left(\frac{1}{3}\right)^5$ $= \frac{17}{81}$ or 0.210 (3 sf)	M1 M1 A1 [3]	Or $1 - P(3,4,5)$ Attempt at correct expression Allow 0.21
(iii)	Est $\text{Var}(P_s) = \frac{0.625 \times (1 - 0.625)}{80}$ $(= \frac{3}{1024})$ $z = 2.054$ (or 2.055) $0.625 \pm z \times \sqrt{\frac{3}{1024}}$ $= 0.514$ to 0.736 (3 sf)	M1 B1 M1 A1 [4]	Any z

Question 6

H_0 : Pop mean = 17 H_1 : Pop mean \neq 17	B1	Both correct. Allow μ , but not just "mean"	
$\frac{18.2 - 17}{\frac{2.4}{\sqrt{5}}}$ = 1.12 (3 sf)	M1	Allow incorrect 18.2. Must have $\sqrt{5}$	$17 \pm 1.96 \frac{2.4}{\sqrt{5}}$ M1
'1.12' < 1.96 oe	A1		= (14.9, 19.1) A1
Claim can be accepted	M1	Comp '1.12' with 1.96 or area '0.132' with 0.025	'14.9' < 18.2 < '19.1' M1
	A1ft	ft their '1.12'	
		If $H_1: \mu > 17$ and cf 1.645: can score max B0M1A1M1A1ft	
	[5]		

Question 7

(i) $E(X) = 3.5$ $(1^2 + 2^2 + 3^2 + 4^2 + 5^2 + 6^2) \div 6 - "3.5"^2$ (= $\frac{35}{12}$ AG)	B1 B1	21/6 2	oe, must see correct expression and no incorrect working
(ii) Attempt $P(X < 3)$ or $1 - P(X \geq 3)$ $N(3.5, \frac{35}{12}/50)$ $\frac{3 - "3.5"}{\sqrt{\frac{35}{12}/50}}$ (= -2.070) $\Phi(" -2.070") = 1 - \Phi("2.070")$ = 0.0192 as final answer	M1 M1 M1 M1		seen or implied seen or implied or $\frac{2.99 - "3.5"}{\sqrt{\frac{35}{12}/50}}$ (= -2.111) $\Phi(" -2.111") = 1 - \Phi("2.111")$ = 0.0174 or 0.0173 Consistent area As final answer or valid total method 5 Allow with incorrect cc (e.g. 2.5) OR no $\sqrt{\quad}$. Must have $\div 50$
(iii) Die is biased (towards lower numbers) Mean of 50 throws ≥ 3 (Allow > 3) or Equal nos of high and low scores or More high scores	B1 indep B1 indep	2	Comment implying die is biased Comment implying results of exp't do not indicate bias (or indicate bias towards higher numbers) Both must be in context

Question 8

(i)	Assume sd unchanged or sd = 10.4	B1		Oe e.g. var unchanged
	H ₀ : Pop mean speed (or μ) = 62.3 H ₁ : Pop mean speed (or μ) < 62.3	B1		Both. Not just "Mean . . "
	$\frac{59.9 - 62.3}{\frac{10.4}{\sqrt{75}}}$	M1		Accept sd/var mixes, but must have $\sqrt{75}$
	= -1.999 or -2.00 (allow + or -) Compare -2.054 or -2.055 No evidence that mean speed decreased	A1 M1 A1 ft	6	Correct z value (or correct critical value) Valid comparison of z's/areas/critical values No contradictions. Do not ft 2-tail test.
(ii)	Pop distribution unknown Yes	B1 B1	2	

Question 9

(i)	P(Type I) = 1 - P(≥ 4 assuming $p = 0.7$) $1 - ({}^6C_4 \times 0.7^4 \times 0.3^2 + {}^6C_5 \times 0.7^5 \times 0.3 + 0.7^6)$ (= 1 - 0.744) = 0.256 (3 s.f.)	M1		or P(≤ 3 assuming $p = 0.7$) May be implied ${}^6C_3 \times 0.7^3 \times 0.3^3 + {}^6C_2 \times 0.7^2 \times 0.3^4 + {}^6C_1 \times 0.7 \times 0.3^5 + 0.3^6$ Allow one end error = 0.256 (3 s.f.) SR if zero scored allow B1 for use of B(6, 0.7) in any two or more terms
		M1		
		A1 [3]		
(ii)	P(Type II) = P(≥ 4 assuming $p = 0.35$) $= {}^6C_4 \times 0.35^4 \times 0.65^2 + {}^6C_5 \times 0.35^5 \times 0.65 + 0.35^6$ = 0.117	M1		May be implied Allow one end error SR if zero scored allow B1 for use of B(6, 0.35) in any two or more terms
		M1		
		A1 [3]		
(iii)	Type 1 They will reject Luigi's belief, although it might be true.	B1 B1	[2]	In context

Question 10

<p>(i)</p> <p>$H_0: p = 0.2$ $H_1: p < 0.2$</p> <p>$P(0 \text{ or } 1 \text{ 5s in } 25 H_0)$</p> <p>$= 0.0274$ (3 s.f.)</p> <p>Comp with 0.025</p> <p>No evidence (at 2.5% level) to support claim</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1✓ [5]</p>	<p>(Allow π)</p> <p>$0.8^{25} + 25 \times 0.8^{24} \times 0.2$ Use of B(25,1/5) and P(0) or P(1) or both – may be implied by “0.0274”</p> <p>Valid comparison</p> <p>No contradictions SR Use of Normal N(5,4) leading to $z = 1.75$ or 0.0401 B1* $H_0 \mu = 5$ $H_1 \mu < 5$ B1. Comparison $1.75 < 1.96$ or $0.0401 > 0.025$ B1* dep</p>
<p>(ii)</p> <p>Normal</p> <p>$\mu = 200, \sigma^2 = 160$ or $\sigma = \sqrt{160}$</p>	<p>B1</p> <p>B1 [2]</p>	
<p>(iii)</p> <p>Concluding that the machine produces the right proportion of 5s, although it doesn't.</p>	<p>B1 [1]</p>	<p>Not concluding that the machine produces too few 5s although it does. Must be in context o.e. No contradictions</p>

Question 11

<p>$\bar{x} = 1.65$</p> <p>$\text{est}(\sigma^2) = \frac{100}{99} \left(\frac{276.25}{100} - 1.65^2 \right)$</p> <p>$= 0.040404\dots = 4/99$</p> <p>$(\pm) \frac{1.65 - 1.6}{\sqrt{\frac{0.040404}{100}}}$</p> <p>$= (\pm) 2.487/2.488$ accept 2.49 Or 0.0065/0.0064 if area comparison done</p> <p>comp with 1.96</p> <p>There is evidence that μ is not 1.6</p>	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1✓ [6]</p>	<p>Without $\frac{100}{99}$: $\frac{1.65 - 1.6}{\sqrt{\frac{0.04}{100}}}$ B1 B0 M1</p> <p>$= 2.50$ A1</p> <p>CV Method M1 must use 1.96 A1 for 1.639 or 1.6106</p> <p>For valid comparison (z/z Signs consistent or area/area cv)</p> <p>Accept Reject H_0 No contradictions</p>
--	--	---

Question 12

<p>(i) 2nd</p> <p>More representative of all appointments or Lengths may vary during the day or 1st does not include later appts so not representative</p>	<p>B1</p> <p>B1 [2]</p>	<p>Any implication that times or conditions vary throughout day, e.g. doctors get tired</p>
<p>(ii) 0.01 o.e.</p> <p>Concluding that times spent are too long when they are not.</p>	<p>B1</p> <p>B1 [2]</p>	<p>Concluding that the mean time spent is more than 10 mins when it is not. Must be in context.</p>
<p>(iii) H₀: Pop mean appt time (or μ) = 10 H₁: Pop mean appt time (or μ) > 10</p> $\frac{147-10}{\frac{12}{\frac{3.4}{\sqrt{12}}}} (\pm)$ <p>= (±)2.292 or (0.0109 if area comparison done)</p> <p>“2.292” < 2.326 o.e.</p> <p>(No evidence to reject H₀.) No reason to believe appts are too long</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1✓ [5]</p>	<p>Both correct. Allow μ, but not just “mean”</p> <p>Allow incorrect $\frac{147}{12}$ Must have $\sqrt{12}$ (accept totals method)</p> <p>For valid comparison Comp “2.292” with 2.326 Or 0.0109 with 0.01 Or 147/12 with 12.28</p> <p>Dep 2.326, ft their “2.292” No contradictions</p> <p>$10 + 2.326 \times \frac{3.4}{\sqrt{12}}$ M1 = 12.28 A1</p> <p>$\frac{147}{12} < 12.28$ M1</p>
<p>(iv) Normal population</p>	<p>B1 [1]</p>	<p>Must have “population” or equiv</p>

Question 13

<p>H₀: Pop mean (or μ or λ) = 50 (or 5)</p> <p>H₁: Pop mean (or μ or λ) \neq 50 (or 5)</p>	<p>B1</p>	<p>Not just “mean”</p>
$\frac{60.5-50}{\sqrt{50}} (\pm)$	<p>M1</p>	<p>For standardising with N(50,50) or N(5,5/$\sqrt{10}$)</p>
<p>= (±)1.485 OR 0.0687 OR C.V</p>	<p>A1</p>	<p>Allow M1 with wrong or no continuity correction OR no $\sqrt{\quad}$ (accept c.v method M1, A1 for 61.63 or 48.868)</p>
<p>1.485 < 1.645 or 0.0687 > 0.05 No evidence that mean changed</p>	<p>M1 A1✓ [5]</p>	<p>For valid comparison (zs or areas or cv) (S.R For cv comparison 61.63 only award final A1 if cc used)</p> <p>or if H₁: $\lambda > 50$, 1.485 < 1.96 M1 No evid mean changed A0 (i.e. if one-tail test, max B0 M1 A1 M1 A0)</p>

Question 14

<p>(i) H_0: Rate = 0.9 H_1: Rate < 0.9 $1 - P(17, 18, 19, 20)$</p> $1 - ({}^{20}C_{17} \times 0.1^3 \times 0.9^{17} + {}^{20}C_{18} \times 0.1^2 \times 0.9^{18} + 20 \times 0.1 \times 0.9^{19} + 0.9^{20})$ $= 0.133 \text{ (3 sf)}$	<p>B1 M1 M1 A1 [4]</p>	<p>$p = 0.9$ $p < 0.9$ Use of B(20,0.1)</p> <p>Allow $1 - P(18,19,20)$ or $1 - P(16,17,18,19,20)$</p>
<p>(ii) Type II H_0 will not be rejected</p>	<p>B1 B1 [2]</p>	<p>or Stephan will conclude standard not fallen No contradictions</p>
<p>[Total: 6]</p>		

Question 15

<p>$H_0: \mu = 250$ $H_1: \mu > 250$</p> $\frac{250.06 - 250}{0.2 \div \sqrt{40}}$ <p>= 1.90 comp with $z = 1.645$ Claim is justified or There is evidence that claim is true</p>	<p>B1 M1 A1 M1 A1 ✓ [5]</p>	<p>Both hypotheses</p> <p>M1 for standardising, must have $\sqrt{40}$. Accept cv method</p> <p>For valid comparison "1.90" with 1.645 or area comparison or CVs Correct conclusion. No contradictions NB 2-tail test scores B0 M1 A1 M1 (use 1.96) A0</p>
<p>[Total: 5]</p>		

Question 16

(i)	<p>H_0: population proportion = 0.1 oe H_1: population proportion > 0.1 oe</p> $P(X \geq 4) = 1 - P(X \leq 3) =$ $1 - \left(0.9^{18} + 18 \times 0.9^{17} \times 0.1 + \right.$ $\left. {}^{18}C_2 \times 0.9^{16} \times 0.1^2 + {}^{18}C_3 \times 0.9^{15} \times 0.1^3 \right)$ <p>= 0.0982 (3 sf)</p> <p>Comp 0.08</p> <p>No evidence that more reach 1m</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1[✓]</p>	<p></p> <p></p> <p></p> <p></p> <p>[5]</p> <p></p> <p>[2]</p> <p></p> <p>[3]</p>	<p>Allow “$p = 0.1$” and “$p > 0.1$”</p> <p>Allow 1 – (one term omitted or extra or wrong) (note CR method 0.0982 and $CR \geq 5$ for A1) Valid comparison ($0.9018 < 0.92$ also recovered previous A1). Or 4 is not in CR Dep M1M1 no contradictions “Accept H_0” provided H_0 defined</p> <p>Ft their (i) If (i) “reject H_0” then ft gives Type I error</p> <p>Attempt $P(X \geq 5)$ e.g. ‘0.0982’ – ${}^{18}C_4 \times 0.9^{14} \times 0.1^4$ oe. Valid comp of their ≥ 5 (if CR method used, could be awarded in (i))</p>
(ii)	<p>Not rejected H_0 Type II</p>	<p>B1[✓] B1dep[✓]</p>	<p></p>	<p></p>
(iii)	<p>$P(X \geq 5)$ (= 0.0282) 0.0282 < 0.08</p> <p>$P(\text{Type I error}) = 0.0282$ (3 sf)</p>	<p>M1 B1[✓]</p> <p>A1</p>	<p></p> <p></p>	<p></p>

Question 17

(i)	<p>“Different” being investigated</p>	<p>B1</p>	<p>[1]</p>	<p>Oe (“changed”, “not equal to”)</p>
(ii)	<p>H_0: Pop mean (or μ) in region same as elsewhere</p> <p>H_1: Pop mean (or μ) in region diff from elsewhere</p> <p>1.91 < 2.054 (or 2.055) or -1.91 > -2.054</p> <p>No evidence that mean is different</p>	<p>B1</p> <p>M1</p> <p>A1</p>	<p></p> <p></p> <p>[3]</p>	<p>Must be “pop mean”, not just “mean” Can be awarded in (i)</p> <p>oe</p> <p>or $P(z > 1.91) = 0.0281 > 0.02$ or $0.0562 > 0.04$ or $0.972 < 0.98$ Accept 2.05 if nothing better seen. inequality sign incorrect M1A0 no contradictions “accept H_0” provided H_0 reasonably well defined</p>

Question 18

(i)	$\lambda = 4.65$ $e^{-4.65} \times \frac{4.65^4}{4!}$ $= 0.186$ (3 sf)	B1 M1 A1	3	Poisson $P(X=4)$ with any λ
(ii)	$\lambda = 3.875$ $= e^{-3.875} \left(1 + 3.875 + \frac{3.875^2}{2!} \right) = 0.257$ (3 sf)	B1 M1 A1	3	$P(X=0, 1, 2)$ Attempted, any λ As final answer
(iii)	$\lambda = 1.5$ $1 - e^{-1.5} \left(1 + 1.5 + \frac{1.5^2}{2!} \right)$ $= 0.191$ (3 sf)	B1 M1 A1	3	$1 - P(X=0, 1, 2)$ Attempted, any λ As final answer
(iv)	He will reject H_0 .	B1	1	
		Total: 10		

Question 19

(i)	$\frac{4.8}{\sqrt{40}}$ $\frac{50.3 - 49.5}{\frac{4.8}{\sqrt{40}}}$ (= 1.054) $1 - \Phi(1.054)$ $= 0.146$ (3 sf)	B1 M1 M1 A1	4	or $\frac{4.8^2}{40}$. Accept $4.8\sqrt{40}$ or $4.8^2 \times 40$ for totals method For standardising with their SD Accept \pm Accept totals method. No mixed methods For use of tables and finding area consistent with their working
(ii) (a)	Looking for decrease	B1	1	
(b)	H_0 : Pop mean time spent (or μ) = 49.5 H_1 : Pop mean time spent (or μ) < 49.5 $\frac{\frac{1920}{40} - 49.5}{\frac{4.8}{\sqrt{40}}}$ (= -1.976) $'1.976' > 1.555$ (or $'-1.976' < -1.555$) There is evidence that mean time has decreased.	B1 M1 M1 A1	4	Not just "mean time spent" For standardising. Allow $\div \frac{4.8}{40}$ Accept totals method; CV method. No mixed methods For valid comparison (area comparison $0.024 < 0.06$) CWO. No contradictions in conclusions
(c)	Population normally distr so No	B1	1	Both needed

Question 20

(i)	Cables broken or not all cables can be accessed oe or Too many cables oe or too time consuming oe	B1 [1]	e.g. previous days' stocks may have gone
(ii)	$H_0: \text{Pop mean brk str (or } \mu) = 5$ $H_1: \text{Pop mean brk str (or } \mu) < 5$ $(\pm) \frac{4.95 - 5}{\frac{0.15}{\sqrt{60}}}$ $(= \pm 2.582)$ comp ± 2.326 There is evidence that mean breaking strength is less than it should be Or reject H_0 (H_0 correctly defined)	B1 M1 A1 B1 ft [4]	Not just "mean" Allow 60 instead of $\sqrt{60}$ Ft their -2.582 (No ft 2 tailed test) Correct comparison shown, no errors seen. Accept area comparison 0.0049 with 0.01 [CR method $(x - 5)/(0.15/\sqrt{60})$ $= -2.326$ M1 A1 leading to $x = 4.955$ compared to 4.95 and correct conclusion B1ft OR $((x - 4.95)/0.15/\sqrt{60})$ leading to 4.995 M1 A1 compared to 5 and correct conclusion B1ft]
(iii)	Population not necessarily normal so yes	B1 B1 dep [2]	SR B1 For "it" is not necc normal (no mention of population) AND Yes

Question 21

(i)	$H_0: P(\text{correct}) = \frac{1}{8}$ $H_1: P(\text{correct}) > \frac{1}{8}$	B1 [1]	Or $H_0 p = 1/8$ $H_1 p > 1/8$
(ii)	$1 - \left(\left(\frac{1}{8} \right)^{10} + 10 \left(\frac{1}{8} \right)^9 \left(\frac{7}{8} \right) + {}^{10}C_2 \left(\frac{1}{8} \right)^8 \left(\frac{7}{8} \right)^2 \right)$ $= 0.120$ (3 sf) or 0.119	M1 A1 A1 [3]	M1 for attempt at correct expression accept 1 error only, e.g. 1 term extra, omitted or wrong, or omit "1-" or incorrect p/q Correct expression Note Use of Poisson in (ii) could score M1 only for expression $1 - P(0,1,2) \lambda = 1.25$
(iii)	12%	B1f [1]	ft their (ii) Must be a probability

Question 21

(i)	Conclude flight times affected when in fact they have not been.	B1 B1	2	Or accept pop mean changed from 6.2 although pop mean has not changed from 6.2
(ii)	H_0 : Pop mean (or μ) = 6.2 H_0 : Pop mean (or μ) \neq 6.2 $\frac{5.98 - 6.2}{\frac{0.8}{\sqrt{40}}}$ = -1.739 (\pm) Accept (\pm)1.74 comp $z = 1.96$ No evidence that flight times affected	B1 M1 A1 B1 \checkmark	4	Allow with 40 instead of $\sqrt{40}$ Allow SD/Var mix (CV method 5.952 or 6.2279 M1 A1) For valid comparison or $P(z < -1.739) = 0.041 > 0.025$ or $5.98 > 5.952$ or $6.2 < 6.228$ and correct conclusion
(iii)	H_0 was not rejected oe Type II	B1* B1*dep	2	If in (ii) H_0 was rejected, then: H_0 rejected B1; Type I B1dep
Total			8	

Question 23

(i)	$H_0: \lambda = 0.5$ $H_1: \lambda > 0.5$	B1	1	or Pop mean = 0.5, not just Mean = 0.5 or Pop mean (per m^2) = 0.1 Accept μ instead of λ
(ii)	$1 - e^{-0.5(1 + 0.5)}$ = 0.0902 (3 sf) comp 0.1 Claim justified or there is evidence to support claim	M1 A1 M1 A1 \checkmark	4	$1 - P(X = 0,1)$ attempted, any λ . Allow 1 end error Allow 0.09 Valid comparison NB $0.9098 > 0.9$ recovers M1A1 M1 oe Accept 'Reject H_0 ' if correctly defined No contradictions.

Question 24

(i)	H_0 : pop mean (or μ) = 12.4 H_1 : pop mean (or μ) > 12.4 $\frac{12.9 - 12.4}{2.1 + \sqrt{50}}$ 1.684 comp cv $z = 1.96$ No evidence that pop mean time has increased	B1 M1 A1 B1f	[4]	not just "mean" Allow with 50 instead of $\sqrt{50}$ or $P(z > 1.684) = 0.0461 > 0.025$ Allow accept H_0 if correctly defined. Ft their test statistic. No contradictions
(ii)	Not reject (or accept) that mean time is unchanged (or is 12.4) oe although mean time has increased (or is more than 12.4) oe	B1 B1	[2]	
(iii)	True (or new) mean	B1	[1]	
			[Total: 7]	

Question 25

<p>(i) $H_0: p = 0.2$ or $\mu = 10$ $H_1: p > 0.2$ or $\mu > 10$</p>	<p>B1 [1]</p>	
<p>(ii) $N(10, 8)$ seen or implied</p> $\frac{125 - 10}{\sqrt{8}} \text{ or } \frac{\frac{125}{50} - 0.2}{\sqrt{\frac{0.2 \times 0.8}{50}}}$ <p>= 0.884</p> <p>comp 1.282</p>	<p>B1 M1 A1 M1f</p>	<p>or $N\left(0.2, \frac{0.2 \times 0.8}{50}\right)$</p> <p>For standardising allow with no or wrong cc</p> <p>Allow area comparison with 0.188 or comp 1.645 if $H_1 p \neq 0.2$</p>
<p>Claim not justified or No evidence to support claim</p>	<p>A1f [5]</p>	<p>Allow accept H_0 provided correctly defined. Follow through their test statistic ;dep 1-tail test No Contradictions</p> <p>SR; Use of $B(50,0.2)$ scores B1 provided at least two probabilities calculated. M1 For finding $P(X \geq 13)$ allow one end error. A1 for 0.186</p>
<p>[Total: 6]</p>		

Question 26

<p>(i) $H_0: \mu = 2.60$ $H_1: \mu > 2.60$</p> $\pm \frac{2.64 - 2.6}{0.2 \div \sqrt{75}}$ <p>= ± 1.732</p> <p>'1.732' > 1.645 Reject H_0. There is evidence that μ has increased</p>	<p>B1 M1 A1 B1 ✓ [4]</p>	<p>allow pop mean, not just 'mean'</p> <p>accept ± 1.73 (3 sf)</p> <p>valid comparison with 1.645 (or $0.0416 < 0.05$) and correct conclusion ✓ their 1.732 no contradictions (or CV method $x_{crit} = 2.638$ M1A1 comp $2.64 > 2.638$ and concln B1 ✓)</p> <p>SR two tail test, using 1.96 (or using 0.025) can score B0M1A1B1ft max 3/4</p>
<p>(ii) $\frac{x - 2.6}{0.2 \div \sqrt{75}} = 1.645$ ($x = 2.638$)</p> $\pm \frac{2.638 - 2.6}{0.2 \div \sqrt{75}}$ <p>= ± 1.819</p> <p>$\Phi(-1.819) = 1 - \Phi(1.819)$</p> <p>$\approx 0.0345$ or 0.0344</p>	<p>M1 M1 A1 M1 A1 [5]</p>	<p>for standardising with their " 2.638 " using 2.68 accept 1.82 (3 sf)</p> <p>indep M mark, calculate correct area/prob consistent with their working</p>

Question 27

(a)	$\lambda = 4.5$ $e^{-4.5}$ (= 0.011109) $\left(\frac{99}{100}\right)^{450}$ (= 0.010860) $\left(\frac{0.011109 - 0.010860}{0.010860} \times 100\right)$ = 2.29% (3 sf)	B1 M1 M1 A1	[4]	alone allow any λ
(b)	$H_0: P(6) = \frac{1}{6}$ or $p = \frac{1}{6}$ $H_1: P(6) < \frac{1}{6}$ or $p < \frac{1}{6}$ $\left(\frac{5}{6}\right)^{25} + 25\left(\frac{5}{6}\right)^{24} \times \frac{1}{6} + {}^{25}C_2 \left(\frac{5}{6}\right)^{23} \times \left(\frac{1}{6}\right)^2$ = 0.189 (3 sf) comp 0.1 No reason to believe die biased	B1 M1 A1 M1 A1	[5]	Both needed allow one error (extra term / missing term / incorrect term) CR method: attempt at least P(0) and P(0 and 1) (0.010... and 0.06... < 0.1) CR is 0,1 and must see 0.189 for A1 valid comp '0.189' with 0.1 oe valid comparison of 2 with CR correct conclusion, ✓ their 0.189 no contradictions

Question 28

(i)	Prob could be different later in day or on a different day oe	B1	[1]	or any explanation why not random or "Not random" or "Not representative"
(ii)	Looking for decrease (or improvement) $H_0: P(\text{not arrive}) = 0.2$ $H_1: P(\text{not arrive}) < 0.2$	B1 B1	[2]	oe Allow " $p = 0.2$ "
(iii)	Concluding that prob has decreased (or publicity has worked) when it hasn't oe	B1	[1]	In context
(iv)	$P(X = 0)$ and $P(X = 1)$ attempted $P(X \leq 2) = 0.8^{30} + 30 \times 0.8^{29} \times 0.2 + {}^{30}C_2 \times 0.8^{28} \times 0.2^2$ (= 0.0442) $P(X \leq 3) = 0.8^{30} + 30 \times 0.8^{29} \times 0.2 + {}^{30}C_2 \times 0.8^{28} \times 0.2^2 + {}^{30}C_3 \times 0.8^{27} \times 0.2^3$ = 0.123 cr is $X \leq 2$ $P(\text{Type I}) = 0.0442$ (3 sf)	M1 M1 B1 A1 A1	[5]	B(30, 0.2) Not nec'y added May be implied by calc $P(X \leq 2)$ or $P(X \leq 3)$ Attempt $P(X \leq 2)$ Or '0.0442' + ${}^{30}C_3 \times 0.8^{27} \times 0.2^3 = 0.123$
(v)	3 is outside cr No evidence that p has decreased (or that publicity has worked)	M1 A1 ✓	[2]	Comparison of 3 with their cr or $P(X \leq 3) = 0.123$ which is > 0.05 Correct conclusion. No contradictions

Question 29

(i)	H_0 : pop mean journey time = 35.2 mins H_1 : pop mean journey time < 35.2 mins	B1	Allow " μ ". Not "mean journey time"
	$\frac{34.7-35.2}{5.6/\sqrt{25}} \quad (= -0.446)$	M1	For standardising ($\sqrt{25}$ needed)
	$\Phi(< "-0.446") = 1 - \Phi("0.446")$ $= 0.328$ (3 sf)	M1 A1	For correct area consistent with their working As final answer
<hr/>			
(ii)	H_0 is rejected but Type II error can only be made if H_0 is <i>not</i> rejected	B1	[1] Allow just " H_0 is rejected." oe

Question 30

H_0 : P(hit target) = 0.65 H_1 : P(hit target) > 0.65	B1	Allow $p = 0.65$ Allow $p > 0.65$
${}^{20}C_2 \times 0.35^2 \times 0.65^{18} + 19 \times 0.35 \times 0.65^{19} + 0.65^{20}$ $= 0.0121$ (3 sf)	M1 A1	Allow one end error. Allow p/q mix. Allow (1-) for M mark A mark recovered following valid comparison
Comp 0.01 There is no evidence (at the 1% level) that she has improved	M1 A1 [✓]	For valid comparison She has probably not improved. No contradictions. (SR Use of Normal M0 , but M1A1 for valid comparison could be awarded)

Question 31

(i)	H_0 : Pop mean = 2.5 (or 7.5) H_0 : Pop mean < 2.5 (or 7.5)	B1	or $\lambda = 2.5$ (Not just "mean") Allow μ or $\lambda < 2.5$
	$\lambda = 7.5$ $P(X \leq 2) = e^{-7.5} (1 + 7.5 + \frac{7.5^2}{2}) = 0.0203$ $P(X \leq 3) = 0.0203 + e^{-7.5} \times \frac{7.5^3}{3!} = 0.0591$	M1 A1	Either $P(X \leq 2)$ or $P(X \leq 3)$, allow any λ <i>Both Correct</i>
	CR is $X \leq 2$	A1	<i>Clear statement</i>
	Reject H_0 Evidence that no of sightings fewer	A1 [✓] [5]	<i>Follow through their CR/their $P(X \leq 2)$</i>
(ii)	$P(\text{Type I}) = 0.0203$ (3 sf)	B1 [✓] [1]	fit their $P(X \leq 2)$
(iii)	H_0 was rejected oe	B1 [1]	or Type II is $P(\text{not reject } H_0)$ oe

Question 32

H_0 : Pop mean yield = 8.2 H_1 : Pop mean yield > 8.2	B1	or $\mu = 8.2$ (not just "mean") $\mu > 8.2$
$(\pm) \frac{8.7-8.2}{1.2/\sqrt{16}}$ $= (\pm) 1.667$ Comp $z = 1.645$ Or Area comparison 0.0475-0.0478	M1 A1 M1	Allow without $\sqrt{\quad}$ sign (Allow cc) Or comp $1 - \Phi('1.667')$ with 0.05 Valid Comparison z-values (same sign) or areas No Contradictions
Reject H_0 Evidence that mean yield has increased	A1 [✓] [5]	No follow through for 2 tail test

Question 33

$H_0: \lambda \text{ (or } \mu) = 42$ $H_1: \lambda \text{ (or } \mu) \neq 42$ $Po(42) \sim N(42, 42)$ stated or implied $\frac{53.5-42}{\sqrt{42}}$ $= 1.77(4)$ (or 0.038 for area comparison) comp 1.96 No evidence that mean has changed	B1 B1 [✓] M1 A1 M1 A1 [✓] [6]	Or pop weekly mean = 2.1 etc. allow 'population mean' not just 'mean' ft their '42' (Accept alt method $N(2.1, 2.1/20)$ allow with wrong or no cc. Accept alt method using $N(2.1, 2.1/20)$ with or without cc Valid comp z or $1 -$ ('1.774') with 0.025 seen allow comp 1.645 if $H_1: \lambda \text{ (or } \mu) > 42$ No contradictions. No ft for $H_1: \lambda \text{ (or } \mu) > 42$ Note – accept other valid methods(e.g. cv method)
---	--	--

Question 34

(i) Conclude less than 90% satisfied when this is not true oe	B11	In context
(ii) $1 - (0.9^{15} + 15 \times 0.9^{14} \times 0.1 + {}^{15}C_2 \times 0.9^{13} \times 0.1^2 + {}^{15}C_3 \times 0.9^{12} \times 0.1^3)$ $= 0.0556$ (3 sf) or 0.0555	M1 M1 A1 [3]	Attempt $(1 -)P(X = 15, 14, 13, 12)$ allow 1 end error Attempt fully correct expression

Question 35

(i) $H_0: \mu = 12.5$ $H_1: \mu \neq 12.5$ $\frac{13.5-12.5}{4.2/\sqrt{50}}$ $= 1.68(4)$ '1.684' < 1.96 No evidence that mean time has changed	B1 M1 A1 M1 A1 ft [5]	allow $4.2 \div 50$ comp 1.96 allow comp 1.645 if $H_1: \mu > 12.5$ or comp $1 -$ ('1.684') with 0.025 No contradictions ft their 1.684, but not comp 1.645
(ii) 0.05	B1 [1]	

Question 36

$B(200, \frac{1}{6}) \rightarrow N(\frac{100}{3}, \frac{250}{9})$ $\frac{25.5 - \frac{100}{3}}{\sqrt{\frac{250}{9}}}$ $= -1.486$ <p>comp '1.486' with 1.282</p> <p>Evidence to reject H_0 There is some evidence that $p < \frac{1}{6}$ or, e.g. It is likely that $p < \frac{1}{6}$ oe</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1 ft [5]</p>	<p>seen or implied</p> <p>allow with wrong or no cc</p> <p>(Accept alternative correct methods)</p> <p>or comp ('1.486') with 0.1</p> <p>No contradictions</p>
--	---	--

Question 37

<p>(i)</p> <p>$H_0: P(\text{free gift}) = 0.3$ or $p = 0.3$ $H_1: P(\text{free gift}) < 0.3$ or $p < 0.3$</p>	<p>B1</p>	<p>[1]</p>
<p>(ii)</p> <p>$P(X \leq 2) =$ $0.7^{20} + 20 \times 0.7^{19} \times 0.3 + {}^{20}C_2 \times 0.7^{18} \times 0.3^2$ $= 0.03548$ or 0.0355</p> <p>$P(X \leq 3) =$ $'0.03548' + {}^{20}C_3 \times 0.7^{17} \times 0.3^3 (=$ $0.107)$</p> <p>One comparison with 0.05 seen</p> <p>$P(\text{Type I error}) = 0.0355$ (3 sf)</p>	<p>M1*</p> <p>A1</p> <p>M1*</p> <p>M1*</p> <p>DA1 ✓</p>	<p>$P(X \leq 2)$ attempted</p> <p>$P(X \leq 3)$ attempted</p> <p>or implied by fully correct methods for $P(X \leq 2)$ and $P(X \leq 3)$</p> <p>dep on all 3 Ms</p>
<p>(iii)</p> <p>$P(X \leq 3) = '0.107'$ $'0.107' > 0.05$ or $cv = 2$ and compare $3 > 2$</p> <p>No evidence to reject claim oe</p>	<p>M1</p> <p>A1 ✓</p>	<p>Compare their $P(X \leq 3)$ with 0.05</p> <p>No evidence that 30% is not correct oe ft their 0.107</p>

Question 38

(i)	H_0 : Pop mean time (or μ) = 20.5 H_1 : Pop mean time (or μ) < 20.5 $\frac{20.3-20.5}{1.2\sqrt{100}}$ = -1.667 or 0.0478/0.952 if areas compared '1.667' < 1.751 (or '-1.667' > -1.751) No evidence that (pop) mean time has decreased	B1 M1 A1 M1 A1ft	[5]	Not just "mean" Allow without $\sqrt{\quad}$ sign (accept $\pm 1.667/1.67$) Correct comparison of their z_{calc} with 1.751/1.75 or valid comparison of areas (0.0478 > 0.04) No contradictions (ft their z)
(ii)	$\frac{cv-20.5}{1.2\sqrt{100}} = -1.751$ $cv = 20.29$ or 20.3 $\frac{20.29-20.1}{1.2\sqrt{100}} (= 1.583 \text{ or } 1.582)$ $1 - \Phi('1.583')$ = 0.0567 – 0.0569 (3 sf)	M1* A1 DM1 M1 A1	[5]	Allow $\frac{20.3-20.1}{1.2\sqrt{100}} (= 1.667)$ M1 $1 - \Phi('1.667')$ M1 = 0.0478 (3 sf) A1
(iii)	Concluding (mean) time not decreased when in fact it has.	B1	[1]	Must be in context oe

Question 39

(i)	H_0 : $P(6) = \frac{1}{6}$ H_1 : $P(6) < \frac{1}{6}$	B1	[1]	Allow $H_0: p = \frac{1}{6}$ $H_1: p < \frac{1}{6}$
(ii)	$(\frac{5}{6})^{15}$ = 0.065 > 0.05	M1 A1	[2]	Correct result and comparison needed for A1 SR if 2 tail test followed allow A1 for 0.065 > 0.025
(iii)	$(\frac{5}{6})^{16} = 0.054$ and $(\frac{5}{6})^{17} = 0.045$ Smallest n is 17 OR $(\frac{5}{6})^n < 0.05$ and attempt to solve $n \ln(\frac{5}{6}) < \ln 0.05$ smallest n is 17	M1 A1 M1 A1	[2]	both No errors seen

Question 40

(i)	$H_0: \text{Pop mean time (or } \mu) = 20.5$ $H_1: \text{Pop mean time (or } \mu) < 20.5$ $\frac{20.3-20.5}{1.2+\sqrt{100}}$ $= -1.667$ or 0.0478/0.952 if areas compared '1.667' < 1.751 (or '-1.667' > -1.751) No evidence that (pop) mean time has decreased	B1 M1 A1 M1 A1ft	[5]	Not just "mean" Allow without $\sqrt{\quad}$ sign (accept $\pm 1.667/1.67$) Correct comparison of their z_{calc} with 1.751/1.75 oe valid comparison of areas (0.0478 > 0.04) No contradictions (ft their z)
(ii)	$\frac{cv-20.5}{1.2+\sqrt{100}} = -1.751$ $cv = 20.29 \text{ or } 20.3$ $\frac{20.29-20.1}{1.2+\sqrt{100}} \quad (= 1.583 \text{ or } 1.582)$ $1 - \Phi('1.583')$ $= 0.0567 - 0.0569 \text{ (3 sf)}$	M1* A1 DM1 M1 A1	[5]	Allow $\frac{20.3-20.1}{1.2+\sqrt{100}}$ (= 1.667) M1 $1 - \Phi('1.667')$ M1 $= 0.0478 \text{ (3 sf)}$ A1
(iii)	Concluding (mean) time not decreased when in fact it has.	B1	[1]	Must be in context oe

Question 41

(i)	$(\lambda =) 4.5$	B1	
	$e^{-4.5}(1 + 4.5 + \frac{4.5^2}{2!})$	M1	Allow any λ . Allow one end error
	$= 0.174$	A1	
	Total:	3	
(ii)	Accept reduction in mean no. of missed appts although untrue	B1	or Mean is 0.9 (or 4.5) but < 3 missed appts. In context
	Total:	1	
(iii)	$P(X \geq 3)$	M1	Attempted
	$= 1 - e^{-1}(1 + 1 + \frac{1^2}{2!})$	M1	Allow any λ except 4.5 or 0.9, Allow one end error
	$= 0.0803 \text{ (3 sfs)}$	A1	

Question 42

(i)	$(H_1): \mu \neq 6.4$	B1	
	Total:	1	
(ii)	comp 2.43 with a z-value $z = 2.576$ AND	M1	oe valid comparison
	No evidence that μ is not 6.4 or do not reject $\mu = 6.4$	A1	Allow "Accept $\mu = 6.4$ " Must mention μ , not just " H_0 " or " H_1 "
	Total:	2	
iii)	Testing for an increase in μ , or for a decrease in μ , rather than a change	B1	Any equiv statement

Question 43

7(i)	H_0 : Pop mean no. accidents = 5.64 H_1 : Pop mean no. accidents < 5.64	B1	or "= 0.47 (per month)" not just "mean", but allow just " λ " or " μ "
	Use of $\lambda = 5.64$	B1	used in a Poisson calculation
	$= e^{-5.64} (1 + 5.64 + \frac{5.64^2}{2})$	M1	Allow incorrect λ in otherwise correct
	= 0.08(0)	A1	
	Comp with 0.05	M1	Valid comparison (Poisson only), no contradictions.
	No evidence to believe mean no. of accidents has decreased; accept H_0 (if correctly defined)	A1FT	Normal distribution: M0M0
	Total:	6	
(ii)	Mean < 0.47 but conclude that this is not so	B1	(Mean) no. of accidents reduced , but conclude not reduced. Must be in context.
	Total:	1	
(iii)	(Need greatest x such that $P(X \leq x) < 0.05$) $P(X \leq 1) = e^{-5.64} (1 + 5.64) = 0.024$ $P(X \leq 2) = 0.08$	B1	Both. could be seen in (i)
	Hence rejection region is $X \leq 1$	B1	Can be implied
	With $\lambda = 12 \times 0.05 = 0.6$, $1 - P(X \leq 1) = 1 - e^{-0.6}(1 + 0.6)$	M1	$\lambda = 0.6$ and $1 - P(X \leq 1)$
	= 0.122 (3 sf)	A1	Normal scores 0

Question 44

(i)	$0.75^{20} + 20 \times 0.75^{19} \times 0.25 + {}^{20}C_2 \times 0.75^{18} \times 0.25^2$	M1	No end errors
	= 0.0913	A1	As final answer
	Total:	2	
(ii)	H_0 : Pop proportion=0.25 H_1 : Pop proportion<0.25	B1	Allow p or π , not "proportion" (Accept anywhere in the question)
	$0.75^{25} + 25 \times 0.75^{24} \times 0.25$	M1	Must be B(25,0.25) No end errors
	= 0.00702	A1	
	comp 0.01	M1	Valid comparison
	There is evidence that the claim is not justified	A1 FT	OE. No contradictions

Question 45

5(i)	mean = 6.6	B1	B1 for 6.6 (could be scored in iii)
	$P(X \leq 1) = e^{-6.6} (1 + 6.6) = 0.0103$	M1	Allow incorrect λ in both probs
	$P(X \leq 2) = e^{-6.6} (1 + 6.6 + \frac{6.6^2}{2}) = 0.0400$	M1A1	A1 for both values
	CR is $X \leq 1$	DA1	Dep on at least one M
	$P(\text{Type I error}) = P(X \leq 1) = 0.0103$	B1FT	FT their $P(X \leq 1)$
	Total:	6	
(ii)	Wrongly concluding that (mean) no of (sports) injuries has decreased	B1	Must be in context
i(iii)	$H_0: \lambda = 6.6$ $H_1: \lambda < 6.6$	B1	Can be scored in (i). Allow μ or $\lambda / 1.1$ or 6.6 or $P(X \leq 2) = 0.0400 > 0.02$
	2 not in CR	M1	
	No evidence mean no. of injuries has decreased	A1FT	
	Total:	3	
i(iv)	$N(39.6, 39.6)$	B1	May be implied
	$\frac{29.5 - 39.6}{\sqrt{39.6}} \quad (= -1.605)$	M1	Allow with wrong or no cc
	$\Phi(-1.605) = 1 - \Phi(1.605)$	M1	For area consistent with their mean
	= 0.0543 (3 sfs)	A1	

Question 46


Assume sd still = 3.8	B1	or sd unchanged
$H_0: \mu = 64.0 \quad H_1: \mu < 64.0$	B1	
$\frac{63.3 - 64.0}{\frac{3.8}{\sqrt{100}}}$	M1	Standardising with their values (no sd / var mixes) Must have $\sqrt{100}$
= -1.842	A1	
comp "1.842" with z-value "1.842" < 1.96	M1	comp +ve with +ve or -ve with -ve or comp Φ ("1.842") with 0.975 0.9672 < 0.975 OE
No evidence that heights are shorter	A1FT	OE FT their z_{calc}

Question 47

3(i)	$\bar{x} = 27/150 (= 0.18)$	B1	
	$s = \sqrt{\frac{150}{149}} \times \sqrt{\frac{5.01}{150} - 0.18^2}$ or variance (= 0.031729) (var = 3/2980 = 0.0010067)	M1	or var = 1/149(5.01 – 27.0 ² /150)
	H ₀ : Pop mean = 0.185 H ₁ : Pop mean < 0.185	B1	allow just ‘μ’
	$\frac{0.18 - 0.185}{\frac{0.031729}{\sqrt{150}}}$	M1	standardising, need $\sqrt{150}$
	= (–) 1.930 (3 sfs) or 1.93	A1	
	Comp with z = (–) 2.326	M1	consistent signs or using probs 0.0268 > 0.01 or 0.9732 < 0.99 or using x _{crit} 0.18 > 0.17897
	There is no evidence (at 1% level) that concentration with drug is less than without drug	A1 FT	conclusion FT no contradictions
(ii)	$\frac{cv - 0.185}{\frac{0.031729}{\sqrt{150}}}$ (= – 2.326)	M1	must use 0.185 and $\sqrt{150}$
	= 0.17897 or 0.179	A1	acceptance region (for H ₀) is > 0.179
	"0.17897" – 0.175 (=1.534) $\frac{0.031729}{\sqrt{150}}$	M1	must use 0.175 and $\sqrt{150}$
	1 – φ("1.534")	M1	indep mark
	= 0.0625 (3 sf)	A1	Accept 0.0610 to 0.0628

Question 48

(i)	H_0 : mean no. sales = 3.5	B1	or “ ... = 0.7 (per day)”
	H_1 : mean no. sales > 3.5	M1	allow ‘ λ ’ or ‘ μ ’ but not just ‘mean’
	$P(X \geq 5) = 1 - e^{-3.5}(1 + 3.5 + \frac{3.5^2}{2!} + \frac{3.5^3}{3!} + \frac{3.5^4}{4!})$	M1	
	= 0.275	A1	allow 0.274
	Comp with 0.10	M1	valid comparison using Poisson
	No evidence (at 10%) to believe that sales per day have increased	A1 FT	correct conclusion FT no contradictions
(ii)	$\lambda = 3.9$	B1	-
	$e^{-3.9} \times \frac{3.9^2}{2!}$	M1	any λ ($\neq 0.7$ or 0.6), single term
	= 0.154 (3 sf)	A1	



Question 50

(i)	$\bar{m} = \frac{98.2}{100} = 0.982$	B1	Accept either
	$s = \sqrt{\frac{100}{99} \times \sqrt{\frac{104.52}{100} - 0.982^2}} (= 0.28582)$ or var = 0.08169	M1	
	H ₀ : Pop mean mass = 1.01 H ₁ : Pop mean mass < 1.01	B1	not just 'mean', but allow just ' μ '
	$\pm \frac{0.982 - 1.01}{\frac{0.28582}{\sqrt{100}}}$	M1	$\pm \frac{0.982 - 1.01}{\frac{0.284387}{\sqrt{100}}}$ M1
	= -0.980 (3 sf) accept \pm	A1	= -0.985 (3 sfs) accept \pm A1
	Comp with $z = -1.645$ (or areas 0.1635 > 0.05)	M1	Valid comparison of z 's or area's
	No evidence that (mean) mass is less than 1.01	A1 FT	Correct conclusion FT their z
(ii)	Distr of X normal (so distr of \bar{X} normal) Must state or imply No	B1	X /parent population

Question 51

(i)	$\bar{x} = 27/150 (= 0.18)$	B1	
	$s = \sqrt{\frac{150}{149} \times \sqrt{\frac{5.01}{150} - 0.18^2}}$ or variance (= 0.031729) (var = 3/2980 = 0.0010067)	M1	or var = 1/149(5.01 - 27.0 ² /150)
	H ₀ : Pop mean = 0.185 H ₁ : Pop mean < 0.185	B1	allow just ' μ '
	$\frac{0.18 - 0.185}{\frac{0.031729}{\sqrt{150}}}$	M1	standardising, need $\sqrt{150}$
	= (-) 1.930 (3 sfs) or 1.93	A1	
	Comp with $z = (-) 2.326$	M1	consistent signs or using probs 0.0268 > 0.01 or 0.9732 < 0.99 or using x_{crit} 0.18 > 0.17897
	There is no evidence (at 1% level) that concentration with drug is less than without drug	A1 FT	conclusion FT no contradictions


(ii)	$\frac{cv - 0.185}{\frac{0.031729}{\sqrt{150}}} (= -2.326)$	M1	must use 0.185 and $\sqrt{150}$
	= 0.17897 or 0.179	A1	acceptance region (for H_0) is > 0.179
	$\frac{0.17897 - 0.175}{\frac{0.031729}{\sqrt{150}}} (=1.534)$	M1	must use 0.175 and $\sqrt{150}$
	$1 - \phi(1.534)$	M1	indep mark
	= 0.0625 (3 sf)	A1	Accept 0.0610 to 0.0628

Question 52

(i)	H_0 : mean no. sales = 3.5	B1	or “ ... = 0.7 (per day)”
	H_1 : mean no. sales > 3.5	M1	allow ‘ λ ’ or ‘ μ ’ but not just ‘mean’
	$P(X \geq 5) = 1 - e^{-3.5}(1 + 3.5 + \frac{3.5^2}{2!} + \frac{3.5^3}{3!} + \frac{3.5^4}{4!})$	M1	
	= 0.275	A1	allow 0.274
	Comp with 0.10	M1	valid comparison using Poisson
	No evidence (at 10%) to believe that sales per day have increased	A1 FT	correct conclusion FT no contradictions
(ii)	$\lambda = 3.9$	B1	
	$e^{-3.9} \times \frac{3.9^2}{2!}$	M1	any λ ($\neq 0.7$ or 0.6), single term
	= 0.154 (3 sf)	A1	

Question 53

(i)	(Po)(2.4)	B1	seen or implied
	$e^{-2.4} \left(1 + 2.4 + \frac{2.4^2}{2} + \frac{2.4^3}{3!} \right)$	M1	allow + P(4)/one end error. Allow wrong λ
	= 0.779 (3 sfs)	A1	Final answer (Note: accept combination method)
		3	
(ii)	$H_0: \lambda$ (or mean) = 3.6 (or 0.9) $H_1: \lambda$ (or mean) < 3.6 (or 0.9)	B1	Accept μ for both
	$e^{-3.6} (1 + 3.6)$	M1	Allow any λ
	= 0.126	A1	
	0.126 > 0.1	M1	Valid comparison. (Comparison with 0.9 could recover previous M1A1)
	No evidence that fewer than usual sold	A1FT	Correct conclusion. No contradictions



Question 54

5(i)	$H_0: P(\text{Orange}) = 0.17$ $H_1: P(\text{Orange}) < 0.17$	B1	or $H_0: p = 0.17$ $H_1: p < 0.17$
i(ii)	Wrongly concluding that % age is less than 17%	B1	OE in context allow "fewer than 3 orange in packet even though average 17% is correct"
		1	
iii)	$B(30, 0.17)$ stated or implied	M1	eg by $0.17^p \times 0.83^q$ ($p + q = 30$) or ${}^{30}C_r$ ($r < 30$)
	$(1 - 0.17)^{30} + 30(1 - 0.17)^{29} \times 0.17 + {}^{30}C_2(1 - 0.17)^{28} \times 0.17^2$	M1	correct, but allow $+ {}^{30}C_3(1 - 0.17)^{27} \times 0.17^3$
	$= 0.0949$ (3 sf)	A1	(SR: use of $N(5.1, 4.233)$ M1 standardising (with or without cc) M1 max 2/3)
iv)	$P(\geq 3 \text{ orange} \mid p = 0.05)$	M1	stated or attempted; can be implied
	$= 1 - [(0.95)^{30} + 30(0.95)^{29} \times 0.05 + {}^{30}C_2(0.95)^{28} \times 0.05^2]$	M1	allow $+ {}^{30}C_3(0.95)^{27} \times 0.05^3$ in bracket, or ans 0.0608
	$= 0.188$ (3 sfs)	A1	

Question 55

5(i)	Assume (pop) sd same (0.3) $H_0: \text{Pop mean} = 2.4$	B1	
	$H_1: \text{Pop mean} \neq 2.4$	B1	Allow ' μ ' but not just 'mean'
	$\pm \frac{2.3 - 2.4}{\frac{0.3}{\sqrt{30}}}$	M1	Must have $\sqrt{30}$, Critical region approach (2.293, 2.507) or (2.193, 2.407)
	$= \pm 1.826$	A1	
	comp $z = \pm 1.96$	M1	Valid comparison (e.g. compare 0.034 with 0.025)
	No evidence that mean time changed	A1f	In context, allow accept H_0 if correctly defined, no contradictions. One-tail test can score B1, B0, M1, A1, M1, A0 Max 4/6
		6	
ii)(a)	0.05	B1	
		1	
ii)(b)	Concluding mean time has not changed when it has.	B1	OE, must have e.g. conclude/accept SR Allow mean has decreased if a one tailed test in Part (i)
		1	

Question 56

(i)	$H_0: P(10) = 0.1$ $H_1: P(10) > 0.1$	B1	Both. Allow 'p' for P(10)
	B(9,0.1) $P(X \geq 3) =$ $1 - (0.9^9 + 9 \times 0.9^8 \times 0.1 + {}^9C_2 \times 0.9^7 \times 0.1^2)$	M1	Allow one extra term in bracket
	$= 0.05297\dots$ or $0.053(0)$	A1	
	comp 0.01	M1	Valid comparison. (comparison with 0.99 can recover previous M1 A1 for 0.9470)
	No evidence (at 1% level) to reject H_0 Claim not justified	A1ft	No contradictions
		5	
(ii)	H_0 not rejected oe	B1	
		1	
(iii)	$P(X \geq 4)$ $= "0.05297" - {}^9C_3 \times 0.9^6 \times 0.1^3$	M1	or $1 - (0.9^9 + 9 \times 0.9^8 \times 0.1 + {}^9C_2 \times 0.9^7 \times 0.1^2 + {}^9C_3 \times 0.9^6 \times 0.1^3)$
	$= 0.00833$	A1	Note: 0.05297 and 0.00833 both needed in (i) or (iii) to justify CV
	Hence crit value is 4	B1	Allow without working. Or in (i) May be implied by attempt at $P(X < 4)$ below
	B(9,0.5) $P(X < 4)$	M1	stated or implied
	$= 0.5^9 + 9 \times 0.5^8 \times 0.5 + {}^9C_2 \times 0.5^7 \times 0.5^2 + {}^9C_3 \times 0.5^6 \times 0.5^3$	M1	Attempt $P(X < 4)$ with $p = 0.5$
	$P(\text{Type II}) = 0.254$ (3 sf)	A1	
		6	

Question 57

7(i)	Po(1.0)	B1	Seen or implied
	$e^{-1} (1 + 1 + \frac{1^2}{2})$	M1	Allow any λ . Allow one end error.
	= 0.920 (3 sfs)	A1	
		3	
7(ii)	$P(X > 3) = 1 - e^{-1.5}(1+1.5+ \frac{1.5^2}{2} + \frac{1.5^3}{3!})$	M1	Allow any λ . Allow one end error
	= 0.0656	A1	
		2	
iii(a)	Incorrectly concluding that more absences than usual when there are not oe	B1	In context
		1	
iii(b)	$H_0: \lambda = 1.5$ (or 0.3) $H_1: \lambda > 1.5$ (or 0.3)	B1	Or μ Both
	$P(X > 4) = "0.0656" - e^{-1.5} \times \frac{1.5^4}{4!}$ = 0.0186 (3 sf)	M1	or $1 - e^{-1.5}(1+1.5+ \frac{1.5^2}{2} + \frac{1.5^3}{3!} + \frac{1.5^4}{4!})$
	$P(\text{Type I}) = 0.0186$ or 0.0185	A1ft	Ft their $P(X > 4)$ if less than 0.05
		3	
iii(c)	$P(X > 3) = "0.0656"$	B1ft	Ft their (ii)
	$0.0656 > 0.05$	M1	
	No evidence of more than usual male absences	A1ft	Ft their $P(X > 3)$. Correct conclusion. No contradictions.

Question 58

(i)	$\frac{14-14.2}{\frac{3.1}{\sqrt{50}}}$ (= -0.456)	M1	For stand'n; must have $\sqrt{50}$
	$1 - \Phi("0.456")$	M1	for area consistent with their working
	= 0.324 (3 sfs)	A1	
		3	
(ii)	No because n large	B1	Accept $n > 30$
		1	
(iii)	$H_0: \mu = 14.2$ $H_1: \mu < 14.2$	B1	or 'pop mean', but not just 'mean'
	$\frac{13.5-14.2}{\frac{3.1}{\sqrt{100}}}$	M1	For stand'n; must have $\sqrt{100}$
	= -2.258	A1	
	comp -2.054 (or -2.055)	M1	Valid comparison of z values or areas (0.0119 < 0.02)
	There is evidence (at 2% level) that mean mass in this area < 14.2	A1ft	Ft their z. Correct conclusion no contradictions
		5	

Question 59

(i)	$H_0: \mu = 51$ $H_1: \mu < 51$	B1	Or popn mean ...
	$\bar{x} = \frac{7480}{150} = 49.8667 = 49.9$	B1	
	$s^2 = \frac{150}{149} \left(\frac{380000}{150} - \left(\frac{748}{15} \right)^2 \right)$ = 46.9620 = 47.0 or $s = 6.85$	M1	Correct subst in s^2 or $\sqrt{s^2}$ formula Biased var scores M0
	$\frac{49.8667-51}{\sqrt{\frac{46.962}{150}}}$ allow $\frac{49.9-51}{\sqrt{\frac{47}{150}}}$	M1	Allow 49.8667 to 49.9 in numerator Need sqrt 150
	= (-) 2.025 = (-) 1.965	A1	Accept 2.02 or 2.03 Accept -2.0264 -1.9651 provided correct working
	comp $z = 1.96$	M1	or comp $1 - \Phi(2.025)$ with 0.025
	There is evidence that $\mu < 51$	A1 ft	no contradictions biased var B1B1M0M1A0M1A1ft (max 5/7) accept cv method $x_{crit} = 49.9028$ M1A1 $49.867 < 49.9 \dots$ M1A1

(ii)	$\frac{\bar{x}-51}{\frac{6.856}{\sqrt{150}}} = -1.96$	M1	Need 51 and sqrt 150 and correct form
	$\bar{x} = 51 - 1.097 = 49.9$ Rejection region is $\bar{x} < 49.9$	A1	This may have been found in part (i)
	$\frac{49.9-49}{\frac{6.856}{\sqrt{150}}}$ (= 1.608 to 1.614)	M1	Need 49 and sqrt 150 and correct form
	$P(\bar{x} > 49.9 \mid \mu = 49) = 1 - \Phi('1.608')$	M1	
	$P(\text{Type II error}) = 0.0539$	A1	Allow 0.0533 to 0.0539
		5	

Question 60

(i)	$H_0: p = \frac{1}{3}$ $H_1: p < \frac{1}{3}$	B1	
		1	
(ii)	$0.0084 < 0.01$	B1	Allow $P(N \leq 36) < 0.01$ or 1%
	There is evidence that p has decreased	B1 dep	Allow ' p has decreased' or $p < \frac{1}{3}$
		2	
(iii)	150	B1	
		1	

Question 61

(i)	$H_0: p = 0.15$ $H_1: p < 0.15$ $(N(60 \times 0.15, 60 \times 0.15 \times 0.85))$ $= N(9, 7.65)$	B1	Accept $H_0: \mu = 9$ $H_1: \mu < 9$ Use of Normal approximation: $(N(0.15, \frac{0.15 \times 0.85}{60}))$ $= N(0.15, 0.002125)$
	$\frac{6.5 - 9}{\sqrt{7.65}}$	M1	For standardising (or $\frac{\frac{6.5 - 9}{60} - 0.15}{\sqrt{0.002125}} = -0.904$) Allow wrong or no cc
	$= -0.904$	A1	Accept \pm
	'0.904' < 1.282	M1	Valid comparison of z values or $\Phi(-0.904) = 0.183 > 0.1$ ft their 0.904
	No evidence train late less often	A1ft	Use of Bin (60,0.15) to give $\Pr(<= 6) = 0.1848$ M1A1 Valid comparison with 0.1 M1 Conclusion A1ft
		5	
(ii)	$0.1 + z \times \sqrt{\frac{0.1 \times 0.9}{60}} = 0.150$	M1	For $\sqrt{(0.1 \times 0.9 / 60)}$ seen
		M1	for $0.1 + z \times \dots = 0.150$ or $2z \dots = 0.1$
	$z = 1.291$	A1	
	$\Phi(1.291) (= 0.90(16))$	M1	for correct method to find α
	$\alpha = 80$	A1ft	ft their z. Must be a +ve non-zero integer < 100
		5	

Question 62

(i)	$\hat{\mu} = \frac{126}{70}$ or $\frac{9}{5}$ or 1.8 oe	B1	
	$\Sigma x^2 f = 286$	B1	Seen or implied
	$\text{Est}(\sigma^2) = \frac{70}{69} (\frac{\Sigma x^2 f}{70} - 1.8^2)$	M1	oe attempted
	$= 0.858$ or $296 / 345$	A1	Note: Final answer for var 0.846 (biased) and no working implies B1 for 286
		4	
(ii)	$H_0: \mu = 1.9$ $H_1: \mu < 1.9$	B1	Or 'pop mean'; not just 'mean'
	$\frac{1.8 - 1.9}{\sqrt{\frac{0.858}{70}}}$	M1	Standardise with their values from (i). Must have sqrt 70. No SD / Var mix
	$= -0.903$	A1	Accept \pm
	$0.903 < 1.645$	M1	comp 1.645 allow comp 1.96 if $H_1: \mu \neq 1.9$ or comp $1 - \Phi(0.903) = 0.182$ or 0.183 with 0.05 (or 0.025 if $H_1: \mu \neq 1.9$)
	No evidence that mean no courts in S is less than in N	A1ft	No contradictions. ft their 0.903, but not comp 1.96 i.e. no ft for a 2 tail test Accept cv method: cv = 1.718 M1A1 1.718 < 1.8 M1 conclusion A1 (cv centred on 1.8 gives 1.982 M1A1 and M1 for 1.982 > 1.9 A1 conclusion)
		5	
(iii)	Type II because H_0 was not rejected	B1ft	ft their conclusion, i.e. if H_0 rejected, 'Type I because H_0 rejected' B1 Answer must be consistent with their conclusion. No conclusion in (ii) will score B0
		1	

Question 63

(i)	Test is for “difference” oe	B1	Test is not for ‘increase’ or ‘decrease’ oe No contradictions
		1	
(ii)	0.05	B1	
	Conclude mean time is different when it is not	B1	oe, in context
(iii)	Assume $\sigma = 6.4$	2	
	Assume $\sigma = 6.4$	B1	
	H_0 : pop mean = 91.4 H_1 : pop mean \neq 91.4	B1	Allow μ , but not ‘mean’
	$\bar{x} = \frac{568.5}{6} (= 94.75)$	B1	
	$\frac{94.75 - 91.4}{\frac{6.4}{\sqrt{6}}}$	M1	Must have $\sqrt{6}$
	= 1.282 cv of $z = 1.96$	A1	
	‘1.282’ < 1.96	M1	Valid comparison or comp $\Phi(“1.282”) with 0.975$ 0.9(001) < 0.975 or 0.0999 (or 0.1) > 0.025 consistent use of one tail test can score M1 for comparison with 1.645oe but not A1ft oe. No contradictions. ft their z.
No evidence mean time different	A1 ft	CV method $x = 96.52$ M1 A1 $94.75 < 96.52$ M1 Conc A1	
	7		

Question 64

$H_0: \lambda = 32$ $H_1: \lambda < 32$	B1	Accept ‘population mean’ (μ)
$X \sim N(32, 32)$	B1	seen or implied
$\frac{21.5 - 32}{\sqrt{32}}$	M1	Standardise with their values. Allow with no or wrong cc
= -1.856 cv of $z = -2.054$ (or -2.055 or -2.053)	A1	
‘1.856’ < 2.054	M1	Valid comparison or comp $\Phi(“1.856”) with 0.98$ i.e. $0.9682 < 0.98$ oe
No evidence that fewer accidents at B than at A	A1f	No contradictions Note Use of CV method $x = 20.38$ M1 A1 comparison $21.5 > 20.38$ M1 conc A1
	6	

Question 65

(i)	$H_0: p = \frac{1}{4}$ $H_1: p > \frac{1}{4}$	B1	
	${}^{10}C_6 \left(\frac{1}{4}\right)^6 \left(\frac{3}{4}\right)^4 + {}^{10}C_7 \left(\frac{1}{4}\right)^7 \left(\frac{3}{4}\right)^3 + {}^{10}C_8 \left(\frac{1}{4}\right)^8 \left(\frac{3}{4}\right)^2 + {}^{10}C_9 \left(\frac{1}{4}\right)^9 \left(\frac{3}{4}\right) + \left(\frac{1}{4}\right)^{10}$	M1	Correct terms, allow one term incorrect or omitted or extra or summing all correct terms from 0 to 5 allow one term incorrect or omitted or extra
	= 0.0197	A1	or 0.9803
	comp '0.0197' with 0.01	M1	Valid comparison with 0.01 or valid comparison with 0.99
	No evidence to conclude $p > \frac{1}{4}$	A1	FT No contradictions Use of two-tail test can score BOM1A1M1(comparison with 0.005) A0
		5	
(ii)	${}^{10}C_7 \left(\frac{1}{4}\right)^7 \left(\frac{3}{4}\right)^3 + {}^{10}C_8 \left(\frac{1}{4}\right)^8 \left(\frac{3}{4}\right)^2 + {}^{10}C_9 \left(\frac{1}{4}\right)^9 \left(\frac{3}{4}\right) + \left(\frac{1}{4}\right)^{10}$	M1	Their $P(X \geq 6) - {}^{10}C_6 (0.25)^6 (0.75)^4$
	P(Type I) = 0.00351 (3 sf)	A1	Accept 0.00348 to 0.00351
		2	
(iii)	C.R is $X \geq 7$ $P(\text{Type II}) = 1 - P(X \geq 7 p = \frac{3}{5}) =$	M1	May be implied
	$1 - ({}^{10}C_7 \left(\frac{3}{5}\right)^7 \left(\frac{2}{5}\right)^3 + {}^{10}C_8 \left(\frac{3}{5}\right)^8 \left(\frac{2}{5}\right)^2 + {}^{10}C_9 \left(\frac{3}{5}\right)^9 \left(\frac{2}{5}\right) + \left(\frac{3}{5}\right)^{10})$	M1	Accept $1 - P(X \geq 8 p = \frac{3}{5})$ or $1 - P(X \geq 6 p = \frac{3}{5})$
	= 0.618	A1	
		3	

Question 66

$H_0: \text{Pop mean} = 20$ $H_1: \text{Pop mean} \neq 20$	B1	Accept μ
$\frac{\sum x}{6} \quad (= \frac{126.9}{6} = 21.15)$	M1	Attempted or 126.9 and 11.64 attempted
$\frac{21.15 - 20}{\sqrt{\frac{1.94}{6}}}$	M1	Must have $\sqrt{6}$ or $\frac{120 - 126.9}{\sqrt{11.64}}$ no mixed method
= 2.022	A1	
$2(1 - \Phi(2.022)) = 2(1 - 0.9784) = 0.0432$	M1	FT $2 \times (1 - 0.9784)$
$\alpha = 4.32$ (3 sf)	A1	FT Allow 4.3 or 4, if correct working seen, or clearly implied, as far as 0.0216 FT their z, no error seen One-tail test scores maximum 3/6
	6	

Question 67

(i)	Max no. of passengers plane can take oe	B1	oe e.g. No of passengers who bought tickets
		1	
(ii)	$\lambda = 3.2$	B1	
	$e^{-3.2} \left(\frac{3.2^3}{3!} + \frac{3.2^4}{4!} + \frac{3.2^5}{5!} \right)$	M1	Any λ . Allow one end error
	$= 0.5146 = 0.515$ (3 sfs)	A1	SR Use of Bin(640,0.005) scores B1 (only) for 0.516
		3	
(iii)	$n > 50$	B1	Accept n is large
	$np = 1.6$, which is < 5 or $p=0.005$ which is < 0.1	B1	Allow $np = 3.2$
		2	
(iv)	H_0 : Pop mean (for 5 days) = 8 H_1 : Pop mean (for 5 days) < 8	B1	or Pop mean (for 1 day) = 1.6 Pop mean (for 1 day) < 1.6 Allow λ or μ but not just 'mean'
	$e^{-8} \left(1 + 8 + \frac{8^2}{2!} \right)$	M1	Any λ ($\neq 1.6$) No end errors. Accept use of Bin(1600,0.005) $P(0,1,2)=0.0136$
	$= 0.0138$	A1	
	Comp 0.025	M1	Valid comparison
	Evidence that mean no. failing to arrive has decreased	A1	FT their '0.0138' or '0.0136'. No contradictions
		5	

Question 68

(i)	$H_0: p = 0.1$ $H_1: p < 0.1$	B1	
		1	
(ii)	B(40, 0.1) stated or implied by use of	B1	e.g. by ${}^{40}C_x$ or $0.9^p \times 0.1^q$ ($p + q = 40$)
	$0.9^{40} + 40 \times 0.9^{39} \times 0.1$	M1	Correct working (if seen). If working not seen, M1 may be implied by 0.0805
	$= 0.0805$	A1	
		3	
(iii)	$z = 1.645$	B1	seen
	$\frac{6}{80} \pm z \sqrt{\frac{\frac{6}{80} \times (80-6)}{80}}$	M1	Formula of correct form. Must be a 'z'
	$= 0.0266$ to 0.123 (3 sfs)	A1	Allow 0.03 to 0.12 or better Must be an interval
		3	
(iv)	10% (or manufacturer's claim) is within CI Hence no reason to question claim	B1	FT Allow '10% is within CI, accept claim' oe Must include both parts. No contradictions. FT their CI Note if CI is centred on 0.1 allow ft 0.075 is within CI, accept claim
		1	

Question 69

(i)	Est(μ) = 1.85	B1	
	Est(σ^2) = $\frac{50}{49} \left(\frac{175.25}{50} - 1.85^2 \right)$	M1	Allow $\sqrt{\frac{50}{49} \left(\frac{175.25}{50} - 1.85^2 \right)}$ or 0.0290 for M1
	= 0.0842 (3 sf) or $\frac{33}{392}$	A1	Cao If $\frac{50}{49}$ omitted (giving var = 0.0825 or sd = 0.287) M0A0
		3	
(ii)	H ₀ : Pop mean time = 1.9 (h) H ₁ : Pop mean time < 1.9 (h)	B1	Allow ' μ ' but not just 'mean'
	$\pm \frac{1.85 - 1.9}{\sqrt{\frac{0.0842}{50}}}$	M1	$\pm \frac{1.85 - 1.9}{\sqrt{0.290}}$ Accept totals method (92.5-95) / $\sqrt{4.21}$
	= -1.22	A1	= -1.22
	comp $z = -1.645$	M1	Or other valid comparison 0.888 or 0.889 < 0.95 OR 0.111 or 0.112 > 0.05
	No evidence that mean time < 1.9 h	A1	FT their z. Correct conclusion. No contradictions If $\frac{50}{49}$ not used in (1): var = 0.8225, sd = 0.907, cr = 1.17 can score all marks in (ii) Note- 2 tail test can score B0 M1 A1 M1 (comparison with 1.96) A0 (no ft) max3/5
		5	

Question 70

(i)	H ₀ : P(6) = $\frac{1}{6}$ H ₁ : P(6) < $\frac{1}{6}$	B1	
	$\left(\frac{5}{6}\right)^{30} + 30\left(\frac{1}{6}\right) \times \left(\frac{5}{6}\right)^{29} + {}^{30}C_2\left(\frac{1}{6}\right)^2 \times \left(\frac{5}{6}\right)^{28}$	M1	Allow one term incorrect, omitted or extra
	= 0.103	A1	
	'0.103' > 0.05	M1	
	No evidence (at 5% level) that die biased	A1ft	oe No contradictions
		5	
(ii)	$\left(\frac{5}{6}\right)^{30} + 30\left(\frac{1}{6}\right) \times \left(\frac{5}{6}\right)^{29}$	M1	
	P(Type I) = 0.0295	A1	
		2	

Question 71

(i)	Assume sd still 4.8 or is unchanged	B1	or Assume the 150 times can be treated as a random sample / are independent
	H ₀ : Pop mean = 26.5 H ₁ : Pop mean > 26.5	B1	Allow ' μ ' but not just 'mean'
	$\frac{27.5 - 26.5}{\frac{4.8}{\sqrt{150}}}$	M1	Standardise, with $\sqrt{\quad}$ Accept CV method
	= 2.552	A1	
	Comp with z-value '2.552' > 2.326	M1	or comp $1 - \Phi('2.552')$ with 0.01 $1 - 0.9946 = 0.0054 < 0.01$
	There is evidence time has increased	A1ft	oe No contradictions (2 tail test scores max. B1 B0 M1 A1 M1 (for comparison with 2.576) A0 no ft)
		6	

(ii)	No because pop is normal so distr of \bar{X} is normal	B1	Condone just 'No because pop is normal'
		1	

Question 72

(i)	H ₀ : Pop mean=546 H ₁ : Pop mean>546	B1	Both. Allow just μ , but not just 'mean'
	$\frac{581 - 546}{\frac{120}{\sqrt{40}}}$	M1	Standardising. Need $\frac{120}{\sqrt{40}}$
	=1.845 allow 1.844	A1	Allow 1.84 or 1.85 AWRT
	1.845 < 1.96	M1	OE. Or area comparison 0.0325 > 0.025 or large probabilities
	No evidence that mean weekly income has increased	A1FT	No contradictions. If H ₁ : ≠, and 2.241 used, max B0M1A1M1A0
		5	
(ii)	$\frac{a - 546}{\frac{120}{\sqrt{40}}} = 1.96$	M1	Standardise to find a . Need $\frac{120}{\sqrt{40}}$ and 546 and a value of z
	$a = 583.19$	A1	Allow 583 to 3sf
	$\frac{583.19 - 595}{\frac{120}{\sqrt{40}}} (= -0.622)$	M1	Standardise. Need $\frac{120}{\sqrt{40}}$ and 595
	$\phi(-0.622) = 1 - \phi(0.622)$	M1	Consistent area
	0.267	A1	
		5	

Question 73

Assume trains are independent OR probability of being on time is constant	B1	Must be in context
H ₀ : P(on time)=0.92 H ₁ : P(on time)<0.92	B1	Both. Allow 'p' or π
1 - $({}^{20}C_{17} \times 0.92^{17} \times 0.08^3 + {}^{20}C_{18} \times 0.92^{18} \times 0.08^2 + 20 \times 0.92^{19} \times 0.08 + 0.92^{20})$	M1	Allow one end error Must have 1 - ...
=0.0706 (3 sf)	A1	
Compare with 0.05	M1	Valid comparison needed
No evidence that percentage less than 92%	A1FT	OE No contradictions. <u>Method using normal approximation:</u> If the first B1B1 is earned then: $CV - 1.566 \left(\text{from } \frac{16.5 - 20 \times 0.92}{\sqrt{20 \times 0.92 \times 0.08}}, \text{ with continuity correction} \right)$ or CV=1.978 (without continuity correction) comp $z=1.645$ No evidence that % decreased (1.566) or evidence that % decreased (1.978) is awarded SC2 after B marks
	6	

Question 74

(i)	H ₀ : Pop mean (or λ or μ) is 1.1 H ₁ : Pop mean (or λ or μ) is more than 1.1	B1	
	$P(X \geq 4) = 1 - e^{-1.1} \left(1 + 1.1 + \frac{1.1^2}{2} + \frac{1.1^3}{3!} \right)$	M1	Correct expression for either $P(X \geq 4)$ or $P(X \geq 5)$
	0.0257	A1	Correct value of either $P(X \geq 4)$ or $P(X \geq 5)$
	$P(X \geq 5) = 0.0257 - e^{-1.1} \times \frac{1.1^4}{4!} = 0.00544$	B1	B1 for the other value (Note use of $P(X < 4) = 0.9743$ and $P(X < 5) = 0.99456$ can score only if comparison with 0.99 seen)
	$0.00544 < 0.01 < 0.0257$	M1	OE stated (valid comparison)
	There is evidence mean has increased	B1	SC $P(X \geq 6) = 0.000968$ M1A1 Conclusion B1
		6	
(ii)	Concluding mean has increased when it has not	B1	In context
	'0.00544'	B1FT	FT <i>their</i> $P(X \geq 5)$, dep < 0.01
		2	
iii)	$e^{-7.0} \left(1 + 7 + \frac{7^2}{2} + \frac{7^3}{3!} + \frac{7^4}{4!} \right)$	M1	Correct expression for $P(X \leq 4 \lambda = 7.0)$
	0.173 (3 sf)	A1	
		2	

Question 75

(i)(a)	Assume standard deviation for the region is 7.1	B1	Or standard deviation is same as for whole population OE
	$\frac{63.2 - 65.2}{\frac{7.1}{\sqrt{n}}} = -2.182$	M1	Attempt to find correct equation (accept +2.182)
	$n = \{-2.182 \times 7.1 \div (-2)\}^2$	A1	Any correct expression for n or \sqrt{n} . SOI
	$n = 60$	A1	CWO. Must be an integer
		4	
(i)(b)	H ₀ : population mean (or μ) = 65.2 H ₁ : population mean (or μ) < 65.2	B1	Not just 'mean'
	$2.182 > 1.751$	M1	Or valid area comparison.
	There is evidence that animals are shorter in this region	A1	CWO. No contradictions
		3	
(ii)	Population unknown or population not given as normal	B1	Allow population not normal. Accept distribution of X unknown.
		1	

Question 76

(a)	$P(X \leq n)$ ($n \leq 20$) attempted, using B(20, 0.95)	M1	OE
	$P(X \leq 17)$ <u>or</u> $P(X \leq 16)$ attempted, using B(20, 0.95)	M1	OE
	$(P(X \leq 17)) = 0.0755$ <u>and</u> $(P(X \leq 16)) = 0.0159$	A1	OE (0.925 and 0.984) both correct
	Rej region is $X \leq 16$ or $X < 17$	A1	Dependent on M1M1 and previous answers correct to at least 0.075/0.076 and 0.016 or 0.92/0.93 and 0.98 Correct unsupported answers of 0.0755 and 0.0159 OE scores M1 M1 A0
		4	
(b)	0.0159	B1	FT <i>their</i> rejection region, from Binomial in a , if $P(X$ in rejection region) < 0.025
		1	
(c)	Use of B(20, 0.7)	M1	
	$P(X > 16 p = 0.7)$	M1	Correct method using B(20, 0.7)
	$= 0.107$	A1	
		3	

Question 77

(a)	$\text{est}(\mu) = 37.6$ or $\frac{1504}{40}$ or $\frac{188}{5}$	B1	
	$\text{est}(\sigma^2) = \frac{40}{39} \left[\frac{57760}{40} - 37.6^2 \right] = 31.0154 = \frac{2016}{65}$	M1	Correct substitution in any correct formula $\frac{1}{39} \left[57760 - \frac{1504^2}{40} \right]$
	$= 31.(0)$ (3 sf)	A1	Accept $\frac{2016}{65}$ or $31\frac{1}{65}$
		3	
(b)	H_0 : Pop mean (or μ) = 39.2 H_1 : Pop mean (or μ) < 39.2	B1	Both. Not just 'mean'
	$\frac{37.6 - 39.2}{\frac{\sqrt{31.0154}}{\sqrt{40}}}$	M1	Allow use of biased variance (30.2), must have $\sqrt{40}$
	$= -1.817$	A1	SC FT use of biased = -1.840 for A1
	'1.817' > 1.645 OE	M1	Valid comparison ' <i>their</i> 1.817' with 1.645 or valid area comparison $0.0346 < 0.05$ OE
	There is evidence that mean time has decreased	A1FT	FT <i>their</i> 1.817; in context, not definite, no contradictions SC For 2 tail test: $H_1: \mu \neq 39.2$ and comp 1.96, max B0M1A1M1A0 (no FT for final mark)
		5	

Question 78

(a)	Later customers might spend times different from first ones	B1
		1
(b)	0.02	B1
	Concluding that $\mu \neq 6.0$, when actually $\mu = 6.0$	B1
		2
(c)	$H_0: \mu = 6.0$ $H_1: \mu \neq 6.0$	B1
	$\frac{6.8 - 6.0}{\sqrt{\frac{4.8}{50}}}$	M1
	2.582	A1
	comp 2.326	M1
	Evidence that $\mu \neq 6.0$	A1
		5
(d)	Population distribution unknown	B1
		1

Question 79

$H_0: \lambda = 104$ (or 5.2) $H_1: \lambda > 104$ (or 5.2)	B1
N(104, 104) stated or implied	B1
$\frac{124.5 - 104}{\sqrt{104}}$	M1
2.010	A1
$2.010 > 1.96$	M1
There is evidence that λ has increased	A1
	6

Question 80

(a)	$E(X) = 2$	B1
	$0.2 \times 1 + 0.4 \times 2^2 + 0.2 \times 3^2 + 0.1 \times 4^2 - 2^2 (= 1.2)$ AG	B1
		2
(b)	$\frac{\sigma - 2}{\sqrt{1.2 \div 200}} = \Phi^{-1}(0.9)$ (M1 for LHS, M1 for RHS)	M1
	$\sigma = 2 + \sqrt{1.2 \div 200} \times 1.282$	M1
	2.10 (3 sf)	A1
		4
(c)	Yes, because X is not normally distributed.	B1
		1
(d)	H_0 : pop mean = 2 H_1 : pop mean < 2	B1
	$\frac{1.86 - 2}{\sqrt{1.2 \div 200}}$	M1
	1.807	A1
	comp $z = 1.645$	M1
	There is evidence that the spinner is biased so that mean is less than 2	A1
		5

Question 81

(a)	H_0 : Proportion = 0.05 H_1 : Proportion > 0.05	B1
		1
(b)	$1 - (0.95^{25} + 25 \times 0.95^{24} \times 0.05 + {}^{25}C_2 \times 0.95^{23} \times 0.05^2 + {}^{25}C_3 \times 0.95^{22} \times 0.05^3)$	M1
	Completely correct expression	A1
	0.0341	A1
		3
(c)	Type II	B1
	Will conclude proportion not increased	B1
		2

Question 82

(a)	$(1^2 + 2^2 + 3^2 + 4^2 + 5^2) \div 5 - 3^2$ (= 2 AG)	B1
		1
(b)	N(3, 2)	M1
	$\frac{2.6 - "3"}{\sqrt{\frac{2}{40}}}$ (= -1.789)	M1
	$\Phi(" -1.789") = 1 - \Phi("1.789")$	M1
	0.0367 to 0.0368	A1
		4
(c)	Concluding that spinner is unbiased when it is biased	B1
		1

Question 83

(a)	Assume standard deviation unchanged or standard deviation = 0.08	B1
	Assume yields normally distributed	B1
		2
(b)	H_0 : Population mean yield (or μ) = 0.56 H_1 : Population mean yield (or μ) > 0.56	B1
	$\frac{0.61 - 0.56}{\frac{0.08}{\sqrt{10}}}$	M1
	1.976	A1
	Comp 1.96	M1
	There is evidence that mean yield has increased	A1
		5

Question 84

(a)	$\frac{40 - 38.4}{\frac{6.9}{\sqrt{30}}} = 1.270$ $\frac{38 - 38.4}{\frac{6.9}{\sqrt{30}}} = -0.3175$	M1
		A1
		A1
	$\Phi('1.270') - (1 - \Phi('0.3175'))$	M1
	= 0.523 (3 sf) or 0.522	A1
		5

(b)(i)	2-tail because looking for 'change', not decrease or increase	B1
		1
(b)(ii)	H_0 : Population mean journey time (or μ) = 38.4 H_1 : Population mean journey time (or μ) \neq 38.4	B1
	$\frac{40.2 - 38.4}{\frac{6.9}{\sqrt{30}}}$	M1
	= 1.429	A1
	'1.429' < 1.645	M1
	There is no evidence that mean journey time has changed.	A1 FT
(b)(iii)	Yes, because population distribution unknown.	B1
		1

Question 85

(a)	$\sqrt{2.1}$ or 1.45 (3 sf)	B1
		1
(b)	$\lambda = 4.2$	B1
	$1 - e^{-4.2}(1 + 4.2)$	M1
	$= 0.922$ (3 sf)	A1
		3
(c)	$\lambda = 6.3$ $e^{-6.3} \left(\frac{6.3^5}{5!} + \frac{6.3^6}{6!} + \frac{6.3^7}{7!} \right)$	M1
	$= 0.455$ (3 sf)	A1
		2
(d)	$H_0: \lambda = 6.3$ $H_1: \lambda < 6.3$	B1
	$P(X \leq 2) = e^{-6.3} \left(1 + 6.3 + \frac{6.3^2}{2!} \right)$	M1
	$= 0.0498$ or 0.0499	A1
	'0.0498' < 0.1	M1
	There is evidence that mean number of absences has decreased.	A1 FT
	5	
(e)	H_0 rejected	*B1 FT
	Hence Type I error possible	DB1 FT
		2

Question 86

i(a)	$H_0: P(\text{contains offer}) = \frac{1}{3}$ $H_1: P(\text{contains offer}) < \frac{1}{3}$	B1
	$P(0,1 \text{ or } 2 \text{ offers in } 20 \mid H_0)$ $= \left(\frac{2}{3}\right)^{20} + 20 \left(\frac{2}{3}\right)^{19} \left(\frac{1}{3}\right) + {}^{20}C_2 \left(\frac{2}{3}\right)^{18} \left(\frac{1}{3}\right)^2$	M1
	= 0.0176 (3sf)	A1
	'0.0176' < 0.1	M1
	(Reject H_0) No evidence (at 10% level) to support manufacturers claim	A1 FT
		5
(b)	$1 - P(X \leq 3)$	M1
	$= 1 - \left[\left(\frac{6}{7}\right)^{20} + 20 \left(\frac{6}{7}\right)^{19} \left(\frac{1}{7}\right) + {}^{20}C_2 \left(\frac{6}{7}\right)^{18} \left(\frac{1}{7}\right)^2 + {}^{20}C_3 \left(\frac{6}{7}\right)^{17} \left(\frac{1}{7}\right)^3 \right]$	A1
	= 0.318 (3sf)	A1
		3
i(c)	Concluding that prop is 1 in 3 when it is actually less(1 in 7)	B1
		1

Question 87

$\text{est}(\mu) = \frac{1850}{200}$ or 9.25	B1
$\text{est}(\sigma^2) = \frac{200}{199} \left(\frac{17850}{200} - \left(\frac{1850}{200} \right)^2 \right)$ or $\frac{1}{199} \left(17850 - \frac{1850^2}{200} \right)$	M1
$= 3.71$ or 3.7060 or $\frac{1475}{398}$	A1
$H_0: \mu = 8.9$ $H_1: \mu \neq 8.9$	B1
$\frac{\frac{1850}{200} - 8.9}{\sqrt{\frac{3.706}{200}}}$	M1
$= 2.57(3\text{sf})$ (or using areas 0.00507 – 0.0051)	A1
$2.24 < 2.57$ or $0.00507 < 0.0125$	M1
(Reject H_0) There is evidence that μ is not 8.9	A1 FT
	8

Question 88

i(a)	H ₀ : population proportion = 0.08 OE H ₁ : population proportion > 0.08 OE	B1	Allow 'p = 0.08' etc.
	$P(X \geq 4) = 1 - P(X \leq 3) =$ $1 - (0.92^{25} + 25 \times 0.92^{24} \times 0.08 + {}^{25}C_2 \times 0.92^{23} \times 0.08^2 + {}^{25}C_3 \times 0.92^{22} \times 0.08^3)$	M1	Allow 1 - (one term omitted or extra or wrong).
	0.135 (3 sf)	A1	
	0.135 > 0.05	M1	Valid comparison. Note: '0.865' < 0.95 can score M1 A1 and can recover previous M1 A1 for 0.865.
	There is no evidence that proportion owning Chantor has increased	A1 FT	In context. Not definite, e.g. not 'Proportion not increased'. No contradictions.
		5	
i(b)	H ₀ was not rejected.	*B1 FT	H ₀ was rejected (consistent with (a)).
	Hence Type II might have been made.	DB1 FT	Type I error.
		2	
i(c)	$P(X \geq 5) = 1 - P(X \leq 4)$ $= 1 - ((1 - 0.1351) + {}^{25}C_4 \times 0.92^{21} \times 0.08^4) [= 0.0451]$	*M1	Attempted. Note: If critical region method used in (a) marks can be awarded here.
	0.0451 < 0.05	A1	Comparison of 0.045[1] with 0.05. Note: If critical region method used in (a) marks can be awarded here.
	P(Type I error) = 0.0451 or 0.0452	A1	Dependent on M1* only. SC Unsupported answers score: B1 for 0.0451 < 0.05 and B1 for final answer 0.0451 only.
		3	

Question 89

i(a)	One-tail because investigating whether "higher"	B1	OE. Must have both parts.
		1	
i(b)	H ₀ : Population mean (or μ) in city same as for others H ₁ : Population mean (or μ) in city greater than for others	B1 FT	If (a) two-tail: H ₀ : Pop mean (or μ) in city same as for others. H ₁ : Pop mean (or μ) in region different from others.
	2.41 > 2.326 or 0.008 < 0.01 or 0.992 > 0.99	M1	If (a) two-tail: 2.41 < 2.576 or 0.992 < 0.995.
	There is evidence that buildings are higher [on average].	A1 FT	In context, not definite. No contradictions. If (a) two-tail: There is no evidence that the [average] height of buildings is different.
		3	

Question 90

(a)(i)	$H_0: \lambda = 2.4$ $H_1: \lambda > 2.4$	B1	Accept λ or μ Accept 2.4 or 0.8 (per year)
		1	
(a)(ii)	$1 - e^{-2.4}(1 + 2.4 + \frac{2.4^2}{2} + \frac{2.4^3}{3!} + \frac{2.4^4}{4!})$	M1	Any λ ; allow one end error
	0.0959 (3 sf)	A1	SC unsupported answer 0.0959 scores B1 only not M1A1
	$0.0959 > 0.05$	M1	Valid comparison Use of $0.9041 < 0.95$ can recover either M1A1 or B1
	There is evidence that Jane's claim not justified or There is insufficient evidence to support Jane's claim	A1 FT	OE. In context, not definite, e.g. not 'Jane is wrong', no contradictions. Condone omission of Jane.
		4	
3(b)	Mean not constant so Poisson model not valid	B1	
		1	

Question 91

(a)	Conclude (mean) (journey) time has not decreased when in fact it has.	B1	OE in context
		1	
(b)	H_0 : Pop mean (or μ) = 1.4 H_1 : Pop mean (or μ) < 1.4	B1	May be seen in (a)
	$\frac{1.36 - 1.4}{\frac{0.12}{\sqrt{50}}}$	M1	Accept totals method $\frac{68 - 70}{\sqrt{50} \times 0.12}$ No mixed methods or no standard deviation/variance mixes
	-2.357 or -2.36	A1	Correct z or correct area if used
	-2.357 < -1.96 or 0.0092 < 0.025 or 0.9908 > 0.975 Or CV method 1.36 < 1.367	M1	valid comparison
	There is evidence that (mean) (journey) times have decreased	A1 FT	in context not definite no contradictions NB use of two tail test scores max B0M1A1M1A0 no ft for two tail test
		5	
(c)	H_0 was rejected OE	*B1 FT	FT H_0 was accepted OE
	Type I	DB1 FT	FT Type II
		2	

Question 92

(a)	Conclude that (population) mean time has changed (or is not 42.4) although μ has not changed (or is still 42.4)	B1	OE. In context.
		1	
(b)	H_0 : population mean (or μ) = 42.4 H_1 : population mean (or μ) \neq 42.4	B1	Not just 'mean'. (could be seen in (a))
	$\pm \frac{45.6 - 42.4}{\sqrt{38.2 \div 20}}$	M1	For standardising (must have $\sqrt{20}$)
	± 2.315	A1	
	$2.240 < '2.315'$	M1	For valid comparison (accept 2.241) or $P(z > 2.315) = 0.0103 < 0.0125$ oe
	There is evidence that μ or mean time has changed	A1 FT	FT <i>their z</i> In context, not definite. No contradictions. Note: Accept correct alternative methods SC: One tail test no FT. Can score B0 M1 A1 M1 (comparison with 1.96) A0 (maximum 3 out of 5)
		5	

Question 93

(a)	$H_0: p = \frac{1}{4}$ $H_1: p \neq \frac{1}{4}$	B1	or $H_0: \mu = 25$ or $H_1: \mu \neq 25$
		1	
(b)	$N\left(25, \frac{75}{4}\right)$	B1	SOI. Allow B1 for $N\left(25, \frac{75}{4}\right)$ or $N(0.25, 0.001875)$ SOI.
	$\pm \frac{15.5 - 25}{\sqrt{\frac{75}{4}}}$ or $\frac{\frac{15.5}{100} - 0.25}{\sqrt{\frac{0.25 \times 0.75}{100}}}$	M1	Standardise with <i>their</i> $N(25, \dots)$ Allow with no or wrong continuity correction.
	± -2.194 (2.19)	A1	
	$-2.326 < -2.194$ or $0.0141 > 0.01$ or $0.9859 < 0.99$	M1	For valid comparison (accept 2.326 to 2.329)
	No evidence to reject that the probability is $\frac{1}{4}$	A1 FT	OE must be in context and not definite, e.g. not 'Claim untrue'. No contradictions. FT <i>their z</i> ; dependent on two-tailed test (one-tailed test can score B1 M1 A1 M1 A0) SC for use of Binomial $B(100, 0.25)$ $P = 0.0111$ for B1 and then comparison with 0.01 and correct conclusion for B1, maximum 2 out of 5 marks.
		5	

Question 94

(a)	Not representative (of all students in the school)	B1	OE idea of 'not being representative' e.g. different grades in the school have different characteristics/proportions.... Don't accept 'not random' or 'biased' without further explanation.
		1	
(b)	H ₀ : P(not correct uniform) = 0.15 H ₁ : P(not correct uniform) < 0.15	B1	Allow "p"
		1	
(c)	Any two probs attempted using B(50,0.15)	M1	
	$P(X \leq 3) = 0.85^{50} + 50 \times 0.85^{49} \times 0.15 + {}^{50}C_2 \times 0.85^{48} \times 0.15^2 + {}^{50}C_3 \times 0.85^{47} \times 0.15^3$	M1	Attempt the tail probability P(0,1,2,3) with B(50,0.15) must be added.
	$P(X \leq 4) = 0.04605 + {}^{50}C_4 \times 0.85^{46} \times 0.15^4$	M1	OE. <i>Their</i> P(X ≤ 3) + P(X = 4) or P(0,1,2,3,4) with B(50,0.15) must be added.
	P(X ≤ 3) = 0.0460 or 0.0461 [<0.05] P(X ≤ 4) = 0.112 or [>0.05]	A1	Both correct. OR if P(X ≤ 4) not seen; P(4)=0.06606 and 0.06606 > 0.05 and P(X ≤ 3)=0.0460 scores M1 A1
	P(Type I) = 0.0460 or 0.0461 (3 sf)	A1	Dependent on second M1. SC If M1M1M1A0 scored allow A1FT for incorrect P(X ≤ 3) as long as <0.05
(d)	4 is outside critical region (≤3) OE or P(X ≤ 4) = 0.112 which is > 0.05	M1	FT working from (c).
	No evidence that proportion not wearing the correct uniform has decreased (Accept H ₀)	A1	In context not definite, e.g. not 'Proportion has not decreased'. No contradiction.
		2	
(e)	Not rejected H ₀	*B1 FT	FT If Reject H ₀ in (d)
	Type II	DB1 FT	FT Type I
		2	

Question 95

(a)	$H_0: \mu = 64.6$ $H_1: \mu < 64.6$	B1	Allow population mean, not just 'mean'.
	$[\pm] \frac{63.5 - 64.6}{5.2 \div \sqrt{100}}$	M1	Standardising. Must have $\sqrt{100}$.
	$[\pm] -2.115$	A1	Accept -2.12 (3sf)
	'2.115' > 1.96 or '-2.115' < -1.96 [do not accept H_0]	M1	Valid comparison (0.0172 < 0.025 for area comparison).
	There is evidence that $\mu < 64.6$	A1 FT	Not definite, e.g. not ' $\mu < 64.6$ '. in context. No contradictions. Accept critical value method leading to $63.5 < 63.58$ or $64.6 > 64.52$.
		5	
(b)	$\frac{m - 64.6}{5.2 \div \sqrt{100}} = -1.96$	M1	Finding the critical value using $N\left(64.6, \frac{5.2}{\sqrt{100}}\right)$ and a z value.
	$m = 63.5808$	A1	
	$\frac{63.5808 - 62.7}{5.2 \div \sqrt{100}} [= 1.694]$	M1	Standardising using $N\left(62.7, \frac{5.2}{\sqrt{100}}\right)$ and a critical value.
	$1 - \Phi('1.694')$	M1	For area consistent with <i>their</i> values.
	0.0451	A1	Accept answers that round to 0.045.
		5	

Question 96

(a)(i)	Po(0.025)	B1	For Poisson and correct parameter.
	$n = 2500 > 50, np = 0.025 < 5$	B1	Must show 2500 and 0.025. Accept $p = \frac{1}{100000} < 0.1$ in place of $np = 0.025 < 5$.
		2	
(a)(ii)	$1 - e^{-0.025}$	M1	Allow any λ . FT <i>their</i> (a)(i) if normal; must have continuity correction.
	0.0247 (3sf)	A1	Must be from Poisson. Unsupported correct answer scores B1 instead of M1 A1 .
		2	
(b)	$H_0: p = 0.3$ $H_1: p < 0.3$	B1	
	$0.7^{28} + 28 \times 0.7^{27} \times 0.3 + {}^{28}C_2 \times 0.7^{26} \times 0.3^2 + {}^{28}C_3 \times 0.7^{25} \times 0.3^3 + {}^{28}C_4 \times 0.7^{24} \times 0.3^4$	M1	Use of B(28, 0.3). Addition of terms must be intended. Allow one term wrong or omitted or extra.
	0.0474	A1	Unsupported correct answer scores B1 instead of M1 A1 .
	$0.0474 > 0.02$ [Not reject H_0]	M1	Valid comparison.
	No evidence that suspicion is true.	A1 ft	Not definite e.g. not 'Suspicion is not true', in context, no contradictions. SC use of N(8.4, 5.88) leading to $0.054 > 0.2$ OE can score B1 only for comparison and correct conclusion. Correct hypotheses with p will also score B1.
		5	

Question 97

(a)	$H_0: P(0) = \frac{1}{10}$ $H_1: P(0) < \frac{1}{10}$	B1	Accept p.
		1	
(b)	For B(30,0.1)	M1	Used not just stated.
	$P(X = 0) = 0.9^{30} [= 0.0424] [<0.1]$	M1	
	$P(X = 0 \text{ or } 1) = 0.9^{30} + 30 \times 0.9^{29} \times 0.1 = 0.184 [>0.1]$	B1	Accept 0.184 or 0.183.
	Rejection region is 0 zeros	A1	Dependent on M1 M1 and at least one comparison, no errors seen. SC One unsupported correct answer 0.0424/0.184(or 0.183) and correct rejection region scores B1 ; with comparison with 0.1 scores B2 . Two unsupported correct answers 0.0424 and 0.184(or 0.183) and correct rejection region scores B2 or if with one comparison with 0.1 scores B3 .
		4	
(c)	0.0424	B1	FT <i>their (b)</i> must have a critical region (only follow though Binomial), dependent on answer < 0.1.
		1	
(d)	Bin(30, $\frac{1}{40}$)	B1	SOI
	$1 - 0.975^{30}$	M1	FT <i>their</i> rr and with Bin(30, 1/40)).
	0.532 (3dp)	A1	SC Unsupported correct answer scores B2 only.
		3	
(e)	Not concluding that the probability is less than $\frac{1}{10}$, when in fact it is.	B1	In context.
		1	

Question 98

(a)	Fireworks are destroyed when tested.	B1	
		1	
(b)	$H_0: \text{Pop mean time lasted (or } \mu) = 30$ $H_1: \text{Pop mean time lasted (or } \mu) < 30$	B1	Not just 'mean'.
	$\pm \frac{29-30}{\frac{5}{\sqrt{100}}}$	M1	For standardising. Must have $\sqrt{100}$. Use of totals N(3000,2500) giving $\frac{(2900-3000)}{\sqrt{2500}}$ scores M1 . No mixed methods.
	± -2	A1	
	$-2 > -2.326$ [Do not reject H_0]	M1	Accept -2.326 to -2.329 . Valid comparison or area comparison $0.0228 > 0.01$ or $0.9772 < 0.99$. Accept CR method $28.837 < 29$ or $30.163 > 30$.
	There is not enough evidence that mean time lasted is less than 30 seconds OR Not enough evidence to support the inspector's suspicion	A1 FT	In context (if used need mean or time / condone average instead of mean), not definite, e.g. not 'mean time lasted is not less than 30 seconds'. No contradictions. Note 2 tailed test can score B0 M1 A1 M1 (comparison with 2.574–2.579) A0 (no FT).
		5	
(c)	Yes. Because population distribution is unknown [condone not Normal].	B1	Both needed. Condone X for parent population.
		1	

Question 99

(a)	$H_0: \mu = 64.6$ $H_1: \mu < 64.6$	B1	Allow population mean, not just 'mean'.
	$[\pm] \frac{63.5 - 64.6}{5.2 \div \sqrt{100}}$	M1	Standardising. Must have $\sqrt{100}$.
	$[\pm] -2.115$	A1	Accept -2.12 (3sf)
	'2.115' > 1.96 or '-2.115' < -1.96 [do not accept H_0]	M1	Valid comparison (0.0172 < 0.025 for area comparison).
	There is evidence that $\mu < 64.6$	A1 FT	Not definite, e.g. not ' $\mu < 64.6$ '. in context. No contradictions. Accept critical value method leading to $63.5 < 63.58$ or $64.6 > 64.52$.
		5	
(b)	$\frac{m - 64.6}{5.2 \div \sqrt{100}} = -1.96$	M1	Finding the critical value using $N\left(64.6, \frac{5.2}{\sqrt{100}}\right)$ and a z value.
	$m = 63.5808$	A1	
	$\frac{63.5808 - 62.7}{5.2 \div \sqrt{100}} [= 1.694]$	M1	Standardising using $N\left(62.7, \frac{5.2}{\sqrt{100}}\right)$ and a critical value.
	$1 - \Phi('1.694')$	M1	For area consistent with <i>their</i> values.
	0.0451	A1	Accept answers that round to 0.045.
		5	

Question 100

(a)(i)	Po(0.025)	B1	For Poisson and correct parameter.
	$n = 2500 > 50, np = 0.025 < 5$	B1	Must show 2500 and 0.025. Accept $p = \frac{1}{100000} < 0.1$ in place of $np = 0.025 < 5$.
		2	
(a)(ii)	$1 - e^{-0.025}$	M1	Allow any λ . FT <i>their</i> (a)(i) if normal; must have continuity correction.
	0.0247 (3sf)	A1	Must be from Poisson. Unsupported correct answer scores B1 instead of M1 A1 .
		2	
(b)	$H_0: p = 0.3$ $H_1: p < 0.3$	B1	
	$0.7^{28} + 28 \times 0.7^{27} \times 0.3 + {}^{28}C_2 \times 0.7^{26} \times 0.3^2 + {}^{28}C_3 \times 0.7^{25} \times 0.3^3 + {}^{28}C_4 \times 0.7^{24} \times 0.3^4$	M1	Use of B(28, 0.3). Addition of terms must be intended. Allow one term wrong or omitted or extra.
	0.0474	A1	Unsupported correct answer scores B1 instead of M1 A1 .
	$0.0474 > 0.02$ [Not reject H_0]	M1	Valid comparison.
	No evidence that suspicion is true.	A1 ft	Not definite e.g. not 'Suspicion is not true', in context, no contradictions. SC use of N(8.4, 5.88) leading to $0.054 > 0.2$ OE can score B1 only for comparison and correct conclusion. Correct hypotheses with p will also score B1.
		5	

Question 101

(a)	$H_0: \mu = 25.5$ $H_1: \mu < 25.5$	B1	
	$\frac{23.7 - 25.5}{5.2 \div \sqrt{40}}$	M1	Must have $\sqrt{40}$
	= -2.189	A1	
	'2.189' < 2.326	M1	For valid comparison For two-tailed test: allow compare 2.576 if $H_1: \mu \neq 25.5$
	[Accept H_0] No evidence that mean time has decreased	A1 FT	In context, not definite, no contradictions FT <i>their</i> 2.189 but no FT for two-tailed test N.B. Use of two-tailed test can score max B0 M1 A1 M1 A0 Condone use of critical value method (23.59 M1 A1 and $23.7 > 23.59$ M1 A1 correct conclusion or 25.612 M1 A1 and $25.5 < 25.612$ M1 A1 with correct conclusion)
		5	
(b)	No, because H_0 was not rejected	B1 FT	FT <i>their</i> conclusion in (a)
		1	

Question 102

	$B(300, \frac{1}{5}) \rightarrow N(60, 48)$	B1	SOI
	$\frac{45.5 - 60}{\sqrt{48}}$	M1	Condone with wrong or no continuity correction
	= -2.093	A1	
	'2.093' > 1.96	M1	Valid comparison Note: $\phi(-2.093)$ (= 0.0182), $0.0182 < 0.025$
	[Evidence to reject H_0] There is evidence that $P(\text{landing on blue}) \neq \frac{1}{5}$	A1 FT	Allow 'There is evidence that the spinner is biased.' In context, not definite, no contradictions Condone critical values method (critical value 46.42 M1 A1 and $45.5 < '46.42'$ M1 for valid comparison A1 for correct conclusion)
			SC: 0.0182 unsupported: $0.0182 < 0.025$ And there is evidence that the spinner is biased. In context, not definite B1 only
		5	

Question 103

(a)	Batteries unusable after testing or Population too big or too costly or too time consuming to use the whole population oe	B1	
		1	
(b)	$H_0: \mu = 150$ $H_1: \mu < 150$	B1	Or population mean = 150; not just 'mean' = 150
	$\frac{147 - 150}{\sqrt{225} \div \sqrt{120}}$	M1	Allow with continuity correction Need $\sqrt{120}$
	-2.191	A1	Condone - 2.19
	-2.191 < -2.054 [or -2.055]	M1	OE. For valid comparison with 2.054 or 2.055 Or 0.0143 (or 0.0142) < 0.02 For two tail test allow comp -2.326 OE if $H_1: \mu \neq 150$ (can score B0M1A1M1A0 max 3/5)
	[Reject H_0] There is evidence that the (mean) life of type B is less than type A (or less than 150)	A1 FT	In context, not definite with no contradictions Accept critical value method 147.19 M1A1 147 < 147.19 M1 conclusion A1 Or 150 > 149.81
		5	
(c)	$147 \pm z \times \frac{15}{\sqrt{120}}$	M1	Expression of correct form must be a z value
	$z = 1.881$ [or 1.882]	B1	
	144 to 150 (3 s.f.)	A1	Must be an interval Incorrect z value can only score M1B0A0
		3	

Question 104

(a)	Conclude more than 10% of the students are left handed when this is not true	B1	OE. Must be in context (accept use of p). Need the context of one tail test.
		1	
(b)	$1 - (0.9^{20} + 20 \times 0.9^{19} \times 0.1 + {}^{20}C_2 \times 0.9^{18} \times 0.1^2 + {}^{20}C_3 \times 0.9^{17} \times 0.1^3 + {}^{20}C_4 \times 0.9^{16} \times 0.1^4)$	M2	M2: fully correct M1: attempt $1 - P(X = 0, 1, 2, 3, 4)$; allow $1 - P(X = 0, 1, 2, 3, 4, 5)$ or $1 - P(X = 0, 1, 2, 3)$ need 1 - ... the method mark cannot be implied
	0.0432 (3 s.f.)	A1	If M0 awarded allow SC B2 for 0.0432
		3	
(c)	$0.7^{20} + 20 \times 0.7^{19} \times 0.3 + {}^{20}C_2 \times 0.7^{18} \times 0.3^2 + {}^{20}C_3 \times 0.7^{17} \times 0.3^3 + {}^{20}C_4 \times 0.7^{16} \times 0.3^4$	M1	Attempt to find $P(\leq 4)$ using B(20,0.3) Allow one end error The method mark cannot be implied
	0.238 or 0.237 (3 s.f.)	A1	If M0 awarded allow SC B1 for 0.238 or 0.237
		2	

Question 105

(a)	H_0 : Pop mean = 4.6 [or 9.2] H_1 : Pop mean < 4.6 [or 9.2]	B1	or $\lambda = 4.6$ or μ (Not just 'mean') or $\lambda < 4.6$
		1	
(b)	Use of Poisson with $\lambda = 9.2$	B1	SOI
	$P(X \leq 3) = e^{-9.2}(1 + 9.2 + \frac{9.2^2}{2} + \frac{9.2^3}{3!}) = 0.0184$ or 0.018 [< 0.02] $P(X \leq 4) = 0.0184 + e^{-9.2} \times \frac{9.2^4}{4!} = 0.0486$ or 0.049 [> 0.02]	M1	At least one of these attempted correct λ (with Poisson expression seen not implied)
	CR is $X \leq 3$	*A1	Both correct SC Use of $\lambda = 4.6$ scores B1 for $P(X = 0) = 0.01[0][1]$ and $P(X \leq 1) = 0.056[3]$ only
		DA1	From CWO and at least one comparison seen SC If M0 awarded allow *B1 for both 0.018 and 0.049 or better and DB1 for correct critical region from CWO and at least one comparison seen.
		4	
(c)	5 is not in critical region OR $P(X \leq 5) = 0.104 > 0.02$ so [not reject H_0] no evidence that number of cars arriving is now fewer	M1 A1 FT	For a comparison (i.e. $5 > 3$) OE In context, not definite No contradictions e.g. not 'No. of cars arriving is not fewer' ft <i>their</i> critical region if used (but must be from Poisson and integers)
		2	
(d)	No, because H_0 was not rejected	B1 FT	OE, FT <i>their</i> (c)
		1	
(e)	$N(276, 276)$	B1	SOI
	$\frac{300.5 - 276}{\sqrt{276}} [= 1.475]$	M1	Standardising with <i>their</i> values Allow with wrong or no continuity correction
	$1 - \Phi(1.475) = 0.0701$ (3 s.f.)	A1	SC Use of Poisson: B1 for answer 0.0727 (3 sf)
		3	

Question 106

H_0 : Pop mean height = 2.3 H_1 : Pop mean height > 2.3	B1	Not just 'mean' Allow μ
$\frac{2.4 - 2.3}{\frac{0.4}{\sqrt{60}}}$	M1	For standardising, must have $\sqrt{60}$
1.936 or 1.937 or 1.94	A1	
'1.936' < 1.96	M1	Valid comparison with 1.96 Or $2.64\% > 2.5\%$ OE Accept $1.936 < 2.24$ or $2.64\% > 1.25\%$ OE if $H_1 \mu \neq 2.3$
[Do not reject H_0] No evidence that (mean) height (with fertiliser) is more than without	A1 FT	FT <i>their</i> z In context, not definite. E.g. not 'Mean height is not greater' with no contradictions No FT for 2 tail test (max B0 M1 A1 M1 A0 3/5) Accept critical values method 2.401 (M1 A1) $2.4 < 2.401$ (M1) Condone 2.299 (M1 A1) < 2.3 (M1) A1 conclusion
	5	

Question 107

(a)	H_0 : pop mean run time = 28.2 mins H_1 : pop mean run time < 28.2 mins	B1	Allow ' μ '. Not 'mean journey time'
		1	
(b)	$\frac{27-28.2}{4/\sqrt{40}} [= -1.897]$	M1	For standardising Must have $\sqrt{40}$
	$\Phi(< -1.897) = 1 - \Phi(1.897)$	M1	For correct area consistent with these values
	0.0289 (3 sf)	A1	
		3	
(c)	H_0 is not rejected so...	M1	
	Type II error can be made and Type I error cannot be made	A1	Both needed (accept 'only a Type II error could be made')
		2	

Question 108

H_0 : P(correct) = $\frac{1}{6}$ H_1 : P(correct) > $\frac{1}{6}$	B1	Allow $p = \frac{1}{6}$ Allow $p > \frac{1}{6}$
$1 - ({}^{15}C_4 \times (\frac{5}{6})^{11} \times (\frac{1}{6})^4 + {}^{15}C_3 \times (\frac{5}{6})^{12} \times (\frac{1}{6})^3 + {}^{15}C_2 \times (\frac{5}{6})^{13} \times (\frac{1}{6})^2 + 15 \times (\frac{5}{6})^{14} \times \frac{1}{6} + (\frac{5}{6})^{15})$	M1	Expression must be seen Allow one end error
0.0898 or 0.0897 (3 sf)	A1	SC if M0 scored allow SCB1 for 0.0898 or 0.0897
0.0898 < 0.1	M1	Valid comparison For valid comparison with 0.9 (0.9102 > 0.9 seen the previous M1 and A1 can be recovered)
[Reject H_0] There is evidence (at the 10% level) that Arvind can predict scores	FTA1	Not definite, e.g. not 'He can predict' or 'Claim true' In context and no contradictions
	5	

Question 109

(a)	H_0 : P(red) = 0.2 H_1 : P(red) < 0.2	B1	Allow H_0 : $p = 0.2$ H_1 : $p < 0.2$.
	$P(X \leq 4) = 0.8^{40} + 40 \times 0.8^{39} \times 0.2 + {}^{40}C_2 \times 0.8^{38} \times 0.2^2 + {}^{40}C_3 \times 0.8^{37} \times 0.2^3 + {}^{40}C_4 \times 0.8^{36} \times 0.2^4$	M1	For full expression seen. Allow one term omitted, incorrect or extra.
	0.0759	A1	SC 0.0759 without working B1.
	their '0.0759' > 0.05	M1	Valid comparison (from binomial probs) of their $P(X \leq 4)$ with 0.05.
	[Do not reject H_0]. Not enough evidence that it lands on red fewer times than if it were fair or not enough evidence to suggest that the spinner is biased	A1 FT	FT their 0.0759. In context, not definite, no contradictions.
		5	
(b)	$P(X \leq 3) = 0.0759 - {}^{40}C_4 \times 0.8^{36} \times 0.2^4$	M1	OE Attempted. Must be using $B(40, 0.2)$. Method could be implied by correct answer here.
	= 0.0285 or 0.0284	*A1	
	Largest value of r is 3	DA1	
		3	

Question 110

(a)	H ₀ : Population mean time (or μ) = 32.5 H ₁ : Population mean time (or μ) < 32.5	B1	Not just "mean".
	$\pm \frac{31.8 - 32.5}{3.1 \div \sqrt{50}}$	M1	Must have $\sqrt{50}$. Could be implied.
	= ± -1.597	A1	
	'-1.597' < -1.406 [or '1.597' > 1.406]	M1	Valid comparison of their z_{calc} with ± 1.406 . or $0.0551 < 0.08$ (or $0.0552 < 0.08$).
	[reject H ₀] There is evidence that [population] [mean] time has decreased	A1 FT	In context, not definite, no contradictions. Note: Accept critical value method 31.88 (31.9) M1 A1 and $31.8 < 31.88$ M1 conclusion A1 .
		5	
(b)	$\frac{a - 32.5}{3.1 \div \sqrt{50}} = -1.406$	M1	Standardise with 32.5 and $\sqrt{50}$ and z value on RHS.
	$a = 31.88$ or 31.9	A1	May be seen in part (a). Can score M1A1 here as well using a similar approach to (a).
	$\frac{\text{their '31.88} - 31.5}{3.1 \div \sqrt{50}}$ [= 0.8668 to 0.8760]	M1	Standardise with <i>their</i> cv and mean = 31.5. Must have $\sqrt{50}$.
	$1 - \Phi('0.8668')$	M1	For area consistent with their working.
	= 0.190 to 0.193 (3 sf)	A1	
		5	

Question 111

	H ₀ : Population mean length = 10.3 cm H ₁ : Population mean length < 10.3 cm	B1	or $\mu = 10.3$ (not just "mean"). $\mu < 10.3$
	$\pm \frac{9.8 - 10.3}{2.6 / \sqrt{100}}$	M1	If ± 1.923 (or 0.0272) seen allow M1 implied.
	= -1.923	A1	Accept \pm . Accept 3sf.
	-1.923 > -2.054 or -2.055	M1	OE For a valid comparison. Or compare $1 - \phi('1.923')$ with 0.02 e.g. $0.0272 > 0.02$ Use of CV $9.8 > 9.766$ scores M1 A1 for 9.766 and M1 for comparison.
	[Not reject H ₀] No evidence that [mean] length has decreased	A1 FT	FT <i>their</i> z . No contradictions, not definite, in context.
		5	

Question 112

(a)	H_0 : Population mean = 7.2 or 2.4 H_1 : Population mean < 7.2 or 2.4	B1	or λ or $\mu = 7.2$ or 2.4 (Not just 'mean'). or λ or $\mu < 7.2$ or 2.4
		1	
(b)	$\lambda = 7.2$	B1	SOI
	$[P(X \leq 2)] = e^{-7.2} \left(1 + 7.2 + \frac{7.2^2}{2} \right)$ or $e^{-7.2}(1 + 7.2 + 25.92)$ or $0.0007465 + 0.0053754 + 0.01935 [= 0.0255]$ $[P(X \leq 3)] = '0.0255' + e^{-7.2} \times \frac{7.2^3}{3!}$ or $'0.0255' + e^{-7.2} (62.21)$ or $'0.0255' + 0.04644 [= 0.0719]$	M1	Both expressions needed, allow any λ If $\lambda \neq 7.2$ allow $P(X \leq n)$ for 2 consecutive values of n with $P(X \leq n) < 0.05$ and $P(X \leq n + 1) > 0.05$.
	$P(\text{Type I}) = 0.02547$ or 0.0255 (3 sf)	B1	
		3	
(c)	$3 > 2$ or $P(X \leq 3) > 0.05$ or $'0.0719' > 0.05$	M1	For a valid comparison or 3 outside critical region. FT <i>their</i> CR in (b).
	[Not reject H_0] No evidence that [mean] number of faults has decreased	A1 FT	No contradictions. In context, not definite.
		2	
(d)	$1 - e^{-1.5}(1 + 1.5 + 1.5^2 / 2)$ or $1 - e^{-1.5}(1 + 1.5 + 1.125)$ or $1 - (0.2231 + 0.3347 + 0.2510)$	M1	Must see expression. FT <i>their</i> CR in (b).
	$= 0.191$ (3 sf)	A1	
		2	

Question 113

(a)	H_0 : Population mean time (or μ) = 32.5 H_1 : Population mean time (or μ) < 32.5	B1	Not just "mean".
	$\pm \frac{31.8 - 32.5}{3.1 \div \sqrt{50}}$	M1	Must have $\sqrt{50}$. Could be implied.
	$= \pm -1.597$	A1	
	$'-1.597' < -1.406$ [or $'1.597' > 1.406$]	M1	Valid comparison of their z_{calc} with ± 1.406 . or $0.0551 < 0.08$ (or $0.0552 < 0.08$).
	[reject H_0] There is evidence that [population] [mean] time has decreased	A1 FT	In context, not definite, no contradictions. Note: Accept critical value method 31.88 (31.9) M1 A1 and $31.8 < 31.88$ M1 conclusion A1 .
		5	
(b)	$\frac{a - 32.5}{3.1 \div \sqrt{50}} = -1.406$	M1	Standardise with 32.5 and $\sqrt{50}$ and z value on RHS.
	$a = 31.88$ or 31.9	A1	May be seen in part (a). Can score M1A1 here as well using a similar approach to (a).
	$\frac{\text{their}' 31.88 - 31.5}{3.1 \div \sqrt{50}} [= 0.8668 \text{ to } 0.8760]$	M1	Standardise with <i>their</i> cv and mean = 31.5. Must have $\sqrt{50}$.
	$1 - \Phi('0.8668')$	M1	For area consistent with their working.
	$= 0.190$ to 0.193 (3 sf)	A1	
		5	

Question 114

(a)	$H_0: P(\text{red}) = 0.2$ $H_1: P(\text{red}) < 0.2$	B1	Allow $H_0: p = 0.2$ $H_1: p < 0.2$.
	$P(X \leq 4) = 0.8^{40} + 40 \times 0.8^{39} \times 0.2 + {}^{40}C_2 \times 0.8^{38} \times 0.2^2 + {}^{40}C_3 \times 0.8^{37} \times 0.2^3 + {}^{40}C_4 \times 0.8^{36} \times 0.2^4$	M1	For full expression seen. Allow one term omitted, incorrect or extra.
	0.0759	A1	SC 0.0759 without working B1.
	<i>their</i> '0.0759' > 0.05	M1	Valid comparison (from binomial probs) of their $P(X \leq 4)$ with 0.05.
	[Do not reject H_0]. Not enough evidence that it lands on red fewer times than if it were fair or not enough evidence to suggest that the spinner is biased	A1 FT	FT <i>their</i> 0.0759. In context, not definite, no contradictions.
		5	
(b)	$P(X \leq 3) = 0.0759 - {}^{40}C_4 \times 0.8^{36} \times 0.2^4$	M1	OE Attempted. Must be using $B(40, 0.2)$. Method could be implied by correct answer here.
	= 0.0285 or 0.0284	*A1	
	Largest value of r is 3	DA1	
		3	

Question 115

(a)	He is expecting a decrease (in μ)	B1	OE
		1	
(b)	$-2.02 < -1.96$	M1	For valid comparison. Allow $2.02 > 1.96$ or $0.0217 < 0.025$ or $0.9783 > 0.975$
	(Reject H_0) There is evidence to suggest that this year's (mean) time is less than 25	A1	OE (such as evidence to support Akash's belief), in context, not definite. No contradictions.
		2	
(c)	$1 - \Phi(2.14) [= 0.0162]$	M1	
	1.62	A1	Allow 1.62% or 1.6 or 1.6%.
	$\alpha \geq 1.62$ (3 sf)	A1ft	FT <i>their</i> 1.62. Allow $\alpha \geq 1.62\%$ or 1.6 or 1.6%. Condone >.
		3	
(d)	$\frac{24.8-m}{3.9-10}$	M1	For standardising.
	$\frac{24.8-m}{3.9-10} = -1.645$	M1	Equate <i>their</i> standardised value to -1.645 (signs must be consistent).
	$m = 25.4$ (3 sf)	A1	
		3	

Question 116

(a)	$e^{-5.7}(1 + 5.7 + \frac{5.7^2}{2!})$ or $e^{-5.7}(1 + 5.7 + 16.245)$ or $0.003346 + 0.01907 + 0.05436$	M1	Allow one end error. Must see this expression.
	= 0.0768 (3 sf)	A1	SC B1 for unsupported answer of 0.0768.
		2	
(b)	$e^{-0.9}(1 + 0.9 + \frac{0.9^2}{2!})$	M1	Attempted; allow one end error (must see expression).
	$= 1 - e^{-0.9}(1 + 0.9 + \frac{0.9^2}{2!}) = 1 - e^{-0.9}(1 + 0.9 + 0.405) = 1 - (0.4066 + 3659 + 0.1647)$	A1	Correct expression $P(X \geq 3)$ no end errors (must see expression).
	= 0.0629 (3 sf)	A1	SC B2 for unsupported answer of 0.0629.
		3	

Question 117

(a)	H ₀ : Pop mean no. people = 3.03 or 1.01 (per 20 min) H ₁ : Pop mean no. people > 3.03 or 1.01 (per 20 min)	B1	These must not just be 'mean', but allow just ' λ ' or ' μ '.
	Use of Po(3.03)	M1	
	$= 1 - e^{-3.03} \left(1 + 3.03 + \frac{3.03^2}{2} + \frac{3.03^3}{3!} + \frac{3.03^4}{4!} + \frac{3.03^5}{5!} \right)$ $= 1 - e^{-3.03} (1 + 3.03 + 4.5905 + 4.6364 + 3.5120 + 2.128)$ $= 1 - (0.04832 + 0.1464 + 0.2218 + 0.2240 + 0.1697 + 0.1028)$	M1	Allow incorrect λ . Allow one end error. Must see Poisson expression used.
	= 0.0870 (3sf) [0.0869727]	A1	Allow 0.087 .
	0.0870 > 0.05	M1	For a valid comparison.
	(Do not reject H ₀) Insufficient evidence to believe (mean) number of people has increased	A1FT	Conclusion stated must be in context, not definite and include no contradictions (e.g. not 'mean number people has not increased').
		6	If only P(x = 6) award max 2/6 (single term not valid). SC No working B1 B2 M1 A1. Award maximum 5/6.
(b)	"0.0869727" - $e^{-3.03} \times \frac{3.03^6}{6!}$ or 0.869727 - $e^{-3.03} (1.0748)$ or 0.869727 - 0.05193 or $1 - e^{-3.03} \left(1 + 3.03 + \frac{3.03^2}{2} + \frac{3.03^3}{3!} + \frac{3.03^4}{4!} + \frac{3.03^5}{5!} + \frac{3.03^6}{6!} \right)$	M1	OE. Must see Poisson expression (may be in part (a)).
	0.0350 or 0.0351	A1	Accept 0.035. SC no working seen, award B1 for 0.0350, 0.0351 or 0.035.
		2	
(c)	Concluding that the (mean) number of people (using the path per 20 mins in the evening) has increased when it has not	B1	OE. Conclusion must be in context.
		1	
(d)	A value for the true mean	B1	Allow without context for this mark.
	Number of people using the path per 20 mins in the evening.	B1	Condone equivalent comment on three randomly chosen 20-minute periods.
		2	

Question 118

(a)	$\bar{x} = 1700/50 = 34$	B1	
	$\text{Est}(\sigma^2) = \frac{50}{49} \left(\frac{59050}{50} - 34^2 \right)$ or $\frac{1}{49} \left(59050 - \frac{1700^2}{50} \right)$	M1	$\text{Est}(\sigma^2) = \frac{59050}{50} - 34^2$ biased scores M0.
	$= 25.5$ (3 sf) or $\frac{1250}{49}$	A1	$= 25$ scores A0.
		3	
(b)	H_0 : Population mean time = 32.4 H_1 : Population mean time \neq 32.4	B1	Not just 'mean' but allow just ' μ '.
	$\frac{34 - 32.4}{\frac{\sqrt{25.5}}{\sqrt{50}}}$	M1	Must have $\sqrt{50}$ and not 50. FT <i>their</i> mean and var. Can be implied.
	$= 2.24$ (3 sf)	A1	or $P(\bar{T} > 34) = 0.0125$. SC use of biased var (25) $z = 2.26$ or $p = 0.0119$, allow M1A1.
	$'2.24' < 2.326$	M1	Or $0.0125 > 0.01$ for a valid comparison.
	[Not reject H_0] Insufficient evidence that (mean) time has changed	A1FT	In context, not definite, e.g. not 'Time not changed'. No contradictions. Note: accept CV method $x_{\text{crit}} = 34.06$ for M1A1. Compares $34 < 34.06$ for M1, conclusion for A1. Condone $x = 32.34$ M1A1: compares $32.4 > 32.34$ for M1, conclusion for A1.
		5	SC for using a one-tail method. Award max 3/5 (B0 M1 A1 M1 A0).

Question 119

(a)	$\left(1 - \frac{1}{3}\right)^{10}$	M1	
	$= 0.0173$ (3 sf)	A1	No working scores SC B1 .
		2	
(b)	$1 - (1 - p)^{10} = 0.8926$	M1	Accept $1 - q^{10} = 0.8926$. Equation must be in p or in q but not both.
	$1 - p = 0.1074^{0.1}$ [= 0.800]	M1	For valid attempt to solve their (binomial) equation in p^{10} or q^{10} .
	$p = 0.200$ (3 sf) or 0.2	A1	
		3	

Question 120

(a)	Est (μ) = 3.25 = 13/4 or 1625/500	B1	
	Est(σ^2) = $\frac{500}{499} \left(\frac{5663.5}{500} - "3.25"{}^2 \right)$ or $\frac{1}{499} \left(5663.5 - \frac{1625^2}{500} \right)$	M1	Expression of correct form.
	= 0.766 (3 sf) or 1529/1996	A1	Biased variance of 0.7645 scores M0A0.
		3	
(b)	H ₀ : Pop mean (or μ) = '3.25' H ₁ : Pop mean (or μ) \neq '3.25'	B1FT	Not just 'mean'. FT their 3.25 .
	$\frac{2.95 - "3.25"}{\sqrt{"0.766" \div 60}}$	M1	Standardising with their values. Must have $\sqrt{60}$.
	= -2.655	A1	Or $P(\bar{X} < 2.95) = 0.0039$ or 0.00396 or 0.00397 . SC FT their biased est(σ^2), i.e. 0.7645 to give $z = 2.658$ A1.
	'2.655' > 2.576 or '-2.655' < -2.576	M1	For valid comparison, e.g. 0.0039 or 0.00396 or 0.00397 < 0.005, or 0.0078 < 0.01, or 0.00792 < 0.01 .
	[Reject H ₀] There is evidence that (mean) mass in (country B) is different (from country A).	A1FT	OE. Must be in context and not definite, e.g., not 'Mean mass is not different', No contradictions. Context needs either 'mass' or 'countries' OE.
			SC , Use of one-tail test. '2.655' > 2.326 or 0.0039 < 0.01 M1A0 (Max B0M1A1M1A0 3/5).
			Accept critical value method. Either: X _{crit} =2.959 M1A1 2.95<2.959 M1A1FT with correct conclusion, or X _{crit} =3.241 M1A1 3.25>3,241 M1A1FT with correct conclusion.
		5	

Question 121

(a)	$H_0: \lambda = 7.6$ [or 1.9] $H_1: \lambda < 7.6$ [or 1.9]	B1	Or Population mean = 7.6 or μ (not just 'mean'). Or Population mean < 7.6 or μ .
		1	
(b)	Mean = 7.6	B1	Seen.
	$P(X \leq 2) = e^{-7.6} (1 + 7.6 + \frac{7.6^2}{2})$ [= 0.0188 or 0.0187]	M1	OE.
	$P(X \leq 3) = e^{-7.6} (1 + 7.6 + \frac{7.6^2}{2} + \frac{7.6^3}{3!})$ [= 0.0554 or 0.0553]	M1	OE. Expression must be seen in at least one probability calculation.
	0.0188 or 0.0187 and 0.0554 or 0.0553	A1	A1 for both values.
	Critical region is $X \leq 2$	A1	Dep on both M marks. SC No Poisson expression seen in either prob scores B1 for 0.0188 or 0.0187 and B1 for 0.0554 or 0.0553 and B1 for CR.
	$P(\text{Type I error}) = P(X \leq 2) = 0.0188$ or 0.0187 (3 sf)	B1FT	FT <i>their</i> $P(X \leq 2)$ or <i>their</i> CR.
		6	
(c)	Concluding that the (mean) no. of accidents has reduced when it has not.	B1	OE. Must be in context. Accept: 'It is believed that the booklet has helped to improve safety when actually it has not'.
		1	
(d)	3 not in critical region.	M1	FT their CR or $P(X \leq 3) = 0.0554 > 0.05$.
	No evidence mean number of accidents has decreased.	A1FT	In context. Cannot be a definite statement, e.g., 'mean number accidents has not decreased'.
		2	

Question 122

Assume SD still = 5.2	B1	OE i.e. 'Assume the SD remains unchanged'.
$H_0: \mu = 24.0$ $H_1: \mu > 24.0$	B1	Or population mean; not just mean.
$\frac{25.8 - 24.0}{\frac{5.2}{\sqrt{50}}}$	M1	For standardising (could be implied). Must have $\sqrt{50}$.
= 2.448	A1	Or $P(\bar{X} > 25.8) = 0.0071$.
'2.448' > 2.326	M1	Or $0.0071 < 0.01$. For valid comparison.
[Reject H_0] There is evidence that (mean) amount of wheat is greater.	A1FT	OE. FT their z_{calc} . In context, not definite, eg not 'Mean amount of wheat is greater' No contradictions CV method: CV = 25.71 M1A1 25.71 < 25.8 M1 A1FT or CV = 24.09 M1 A1 24.09 > 24 M1 A1FT.
	6	