

Markscheme

November 2024

Mathematics: analysis and approaches

Standard level

Paper 1

19 pages



© International Baccalaureate Organization 2024

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2024

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2024

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *M0* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies A3, M2 etc., do not split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this working is incorrect and/or suggests a misunderstanding of the question. This will encourage a uniform approach to marking, with less examiner discretion. Although some candidates may be advantaged for that specific question item, it is likely that these candidates will lose marks elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used **in a subsequent part**. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	8√2	5.65685 (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)
2.	$\frac{35}{72}$	0.468111… (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)

3 Implied marks

Implied marks appear in **brackets e.g.** (*M1*), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is *(M1)A1*, it is possible to award full marks for *their* correct answer, **without working being seen**. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (*MR*). A candidate should be penalized only once for a particular misread. Use the *MR* stamp to indicate that this has been a misread and do not award the first mark, even if this is an *M* mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** . . . **OR**.

7 Alternative forms

Unless the question specifies otherwise, accept equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, *M* marks and intermediate *A* marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come *"from the use of 3 sf values"*.

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an *A* mark to be awarded, arithmetic should be completed, and any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$. An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or written as $\frac{5}{2}$.

However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate A marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".

Section A

1.	(a)	(i)	27	A1
		(ii)	30	A1
		(iii)	35	A1
				[3 marks]
,	(b)	$Q_1 =$	31 and $Q_3 = 46$	(A1)
		atter	npt to subtract their upper and lower quartiles	(M1)
	(b)	$Q_1 =$ atter te: A	31 and $Q_3 = 46$ npt to subtract their upper and lower quartiles ward M1 only for correct values, or for values clearly indicated as ca	(A1 $(M1)$ Indidate's Q_1 and
		Q	B. PR	

46-31

IQR = 15

A1 [3 marks] [Total: 6 marks]

2. (a)
$$\frac{1}{2}r^2(\theta) = 6$$
 OR $\frac{1}{2}r^2(0.75) = 6$ (A1)

attempt to solve their equation to find
$$r$$
 or r^2 (M1)
Note: To award the M1, candidate's equation must include r^2 and $\theta = 1.5$, and they
must isolate r^2 or r .
 $r^2 = 16$
 $r = 4$ (cm) A1
[3 marks]
(b) evidence of summing the two radii and the arc length (M1)
perimeter $= 2r + r\theta$
 $= 8 + 4(0.75)$
 $= 11$ (cm) A1
[2 marks]
Total [5 marks]

3. (a) use of
$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$
 (M1)

$$P(A \cap B) = 0.45 + 0.65 - 0.8$$
 (or equivalent) (A1)
= 0.3 A1

[3 marks]

(b)
$$P(A' \cap B') = 0.2$$
 (may be seen in Venn diagram) (A1)

attempt to substitute their values into
$$P(A' | B') = \frac{P(A' \cap B')}{P(B')}$$
 (M1)

$$P(A' | B') = \frac{0.2}{0.35}$$
$$= \frac{20}{35} \left(= \frac{4}{7} \right)$$

A1

[3 marks] Total [6 marks]

4. METHOD 1

attempt to expand	$(2n+5)^2 - (2n-5)^2$	$5)^{2}$
-------------------	-----------------------	----------

Note: Award *M0* for invalid attempts such as $(2n+5)^2 = 4n^2 + 25$.

$$= 4n^{2} + 20n + 25 - (4n^{2} - 20n + 25)$$
 or equivalent **A1**

$$=40n \text{ OR } 20n+20n$$
 A1

=
$$10(4n)$$
 OR $\frac{40}{10} = 4$ OR $\frac{40n}{4} = 10n$ OR $20n + 20n = 10(2n + 2n)$ (or equivalent) **R1**
so is a multiple of 10 **AG**

Note: Do not award the R1 unless both A marks have been awarded.

METHOD 2

use of $a^2 - b^2 = (a+b)(a-b)$ where $a = 2n+5, b = 2n-5$	M1
= (2n+5+2n-5)(2n+5-2n+5)	A1A1
Note: Award A1 for each correct bracket.	
$=4n\times10(=40n)$	R1
so is a multiple of 10	AG
Note: Do not award the <i>R1</i> unless both <i>A</i> marks have been awarded.	

[4 marks]

(M1)

(M1)

5. attempt to substitute into cosine rule

$$(\cos 2\theta =)\frac{3^2 + 10^2 - 8^2}{2 \times 3 \times 10}$$
 OR $8^2 = 3^2 + 10^2 - 2 \times 3 \times 10 \times \cos 2\theta$

$$\left(\cos 2\theta =\right)\frac{45}{60}\left(=\frac{3}{4}\right) \tag{A1}$$

attempt to use $\cos 2\theta = 2\cos^2 \theta - 1$

$$\cos^{2}\theta = \frac{1 + \frac{45}{60}}{2} \left(= \frac{1 + \frac{3}{4}}{2} \right)$$

$$\cos^{2}\theta = \frac{105}{120} \left(= \frac{7}{8} \right)$$

$$\cos\theta = (\pm)\sqrt{\frac{105}{120}} \left(= \sqrt{\frac{7}{8}} \right)$$

$$= \frac{\sqrt{7}}{2\sqrt{2}}$$

$$= \frac{\sqrt{14}}{4} (p = 14, q = 4)$$
Note: The final answer must be positive.

[6 marks]

(M1)

(M1)

(M1)

6. attempt to use $u_n = u_1 + (n-1)d$ or $S_n = \frac{n}{2} [2u_1 + (n-1)d]$ or $S_n = \frac{n}{2} [u_1 + u_n]$ to set up at least one equation in u_1 and d

$$14 = u_1 + 9d$$
 and $200 = \frac{25}{2} [2u_1 + 24d]$ **A1**

attempt to solve their two linear equations in u_1 and d simultaneously (must eliminate one variable)

$$d = -2 (\Longrightarrow u_1 = 32) \tag{A1}$$

attempt to solve $u_k = 0$ with their d (or with their d and u_1)

$$\Rightarrow k = 17$$
 A1





– 13 –

Section B

7. (a)	f(x) = -2x + 12 (accept $y = -2x + 12$, accept $m = -2$ and $c = 12$)	A1A1
Note: A	ward A1 for correct gradient, A1 for correct y-intercept	
		[2 marks]
(b)	METHOD 1 Axis of Symmetry	
	axis of symmetry is $\frac{-2+6}{2}(=2)$	(A1)
	equating $\frac{-b}{2a}$ to their axis of symmetry	(<i>M1</i>)
	$\frac{-b}{-2} = 2$	
	<i>b</i> = 4	A1
	METHOD 2 substitution	
	attempt to substitute $(-2,0)$ or $(6,0)$ into g	(<i>M1</i>)
	$-(-2)^2 - 2b + 12 = 0 \text{ or } -(6)^2 + 6b + 12 = 0$	(A1)
	<i>b</i> = 4	A1
	METHOD 3 factored form	
	attempt to write g in factored form	(M1)
	g(x) = (x+2)(6-x)	
	$-x^2 + 4x + 12$	(A1)
	<i>b</i> = 4	A1
	METHOD 4 quadratic formula	
	attempt to substitute into quadratic formula and set equal to -2 or 6	(M1)
	$6 = \frac{-b \pm \sqrt{b^2 - 4(-1)(12)}}{2(-1)} \text{OR} -2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(12)}}{2(-1)}$	
	$b^2 - 24b + 144 = b^2 + 48$ OR $b^2 + 8b + 16 = b^2 + 48$ (or equivalent, must nervice) radical)	ot contain (A1)
	<i>b</i> = 4	A1
		[3 marks]
		continued

Question 7 continued

(c) recognizing to subtract
$$g - f$$
 (in correct order) (M1)

$$\int_{0}^{6} \left(-x^{2} + 4x + 12 \right) - \left(-2x + 12 \right) dx$$
 A1

$$\int_0^6 \left(-x^2 + 6x\right) \mathrm{d}x \qquad \qquad \mathbf{AG}$$

$$-\frac{1}{3}x^3 + 3x^2(+C)$$
 A1

attempt to substitute limits into their integrated function and find difference (M1)

$\left(-\frac{1}{3} \cdot 6^{3} + 3 \cdot 6^{2}\right) - \left(-\frac{1}{3} \cdot 0^{3} + 3 \cdot 0^{2}\right)$	
36	A1
	[4 marks]

(e)	g'(x) = -2x + 4	A1
	attempt to equate their derivative of g to their gradient of f	(M1)
	-2x + 4 = -2	A1
	<i>x</i> = 3	A1
	<i>y</i> = 15	A1
		[5 marks]
		Total [16 marks]

– 15 –

8. (a) (i) substitution of x = 2 in part (i) or $x = \frac{1}{8}$ in part (ii) (M1)

$$\log_2(8 \times 2) \text{ OR } \log_2(16)$$

 $\log_2(16) = 4$ A1

(ii)
$$f\left(\frac{1}{8}\right) = 0$$
 A1

[3 marks]

(b) swap
$$x$$
 and y (M1)

$$x = \log_2(8y) \text{ OR } x = 3 + \log_2 y$$

attempt to write as exponential (*M1*)

$$2^{x} = 8y$$

$$(f^{-1}(x) =) \frac{2^x}{8} (= 2^{x-3}) \left(\text{accept } y = \frac{2^x}{8} \text{ or } y = 2^{x-3} \right)$$
 A1

[4 marks]

A1

(A1)

[1 mark]

continued...

 $\left(f^{-1}
ight)$

(c) $\frac{1}{8}$

A2

(M1)

(M1)

A1

Question 8 continued

(d) METHOD 1

$f(4x^2) = \log_2(8 \times 4x^2)$	
attempt to use addition rule for logs	(M1)
$\log_2 8 + \log_2 4 + \log_2 x^2 \text{ OR } \log_2 32 + \log_2 x^2$ (or equivalent)	(A1)
attempt to use exponent property for logarithms	(<i>M1</i>)
$f(4x^2) = 5 + 2\log_2 x$ (or equivalent)	A1
the graph of g must be vertically stretched (dilated) by a scale factor of 2 and then	

the graph of g must be vertically stretched (dilated) by a scale factor of 2 and then vertically translated (shifted) 5 units upwards.

METHOD 2

$$f(4x^2) = \log_2(8 \times 4x^2)$$

attempt to write argument as a power

$$\log_2(32x^2) = \log_2\left(\left(\sqrt{32x}\right)^2\right) \text{ (or equivalent)}$$
(A1)

attempt to use exponent property for logarithms

$$f(4x^2) = 2\log_2(\sqrt{32}x)$$
 (or equivalent) A1

EITHER

the graph of g must be vertically stretched (dilated) by a scale factor of 2 and stretched (dilated) horizontally by a scale factor of $\frac{1}{\sqrt{32}}$.

OR

the graph of g must be stretched (dilated) horizontally by a scale factor of $\frac{1}{\sqrt{32}}$ and

vertically stretched (dilated) by a scale factor of 2.

Note: In this method, the final mark is **A1**, as the question specifically asks for a translation and a stretch.

[6 marks] [Total: 14 marks] – 17 –

9. (a) outer curved surface area is $2\pi(4r)h$ AND inner curved surface area is $2\pi rh$ (A1) area of each base (top and bottom) is $\pi(4r)^2 - \pi r^2$ (A1)

$$S = 2\left[\pi(4r)^{2} - \pi r^{2}\right] + 2\pi(4r)h + 2\pi rh$$
 A1

$$=30\pi r^2 + 10\pi rh$$

(b) $30\pi r^2 + 10\pi rh = 240\pi$

attempt to solve their equation for h or rh in terms of r (must isolate h or rh) (M1)

$$h = \frac{240 - 30r^2}{10r} \left(= \frac{24 - 3r^2}{r} \right) \text{OR } rh = \frac{240 - 30r^2}{10} \left(= 24 - 3r^2 \right) \text{ (or equivalent)}$$
 A1

uses volume = large cylinder – small cylinder (M1)

$$V = \pi (4r)^2 h - \pi r^2 h \quad (= 16\pi r^2 h - \pi r^2 h = 15\pi r^2 h)$$

attempt to substitute in for h or rh

$$V = 15\pi r^2 \left(\frac{24 - 3r^2}{r}\right) \text{ OR } V = 15\pi r \left(\frac{240 - 30r^2}{10}\right) \left(=15\pi r \left(24 - 3r^2\right)\right) \text{ OR}$$

$$384\pi r - 48\pi r^3 - 24\pi r + 3\pi r^3 \text{ A1}$$

 $=360\pi r - 45\pi r^3$

[6 marks]

(M1)

(c)
$$\frac{dV}{dr} = 360\pi - 135\pi r^2$$
 A1A1

[2 marks]

continued...

Question 9 continued

(d) METHOD 1 (working with r) recognition that (for a maximum) $\frac{dV}{dr} = 0$ М1 $360\pi - 135\pi r^2 = 0$ $r^2 = \frac{360}{135} \left(= \frac{8}{3} \right)$ $r = \sqrt{\frac{360}{135}} \left(= \sqrt{\frac{8}{3}} \right)$ A1 $p = 2 \text{ OR } r = 2\sqrt{\frac{2}{3}}$ A1 **METHOD 2** (working with $p\sqrt{\frac{2}{3}}$) recognition that (for a maximum) $\frac{\mathrm{d}V}{\mathrm{d}r} = 0$ М1 $360\pi - 135\pi \left(p\sqrt{\frac{2}{3}}\right)^2 = 0$ $360 - 90p^2 = 0$ $p^2 = 4$ A1 $p = 2 \text{ OR } r = 2\sqrt{\frac{2}{3}}$ A1 [3 marks]

continued...

Total [17 marks]

Question 9 continued

(e)

attempt to substitute their value of r into $V = 360\pi r - 45\pi r^3$ M1 $V = 360\pi \times 2\sqrt{\frac{2}{3}} - 45\pi \times \left(2\sqrt{\frac{2}{3}}\right)^3$ $360\pi \times 2\sqrt{\frac{2}{3}} - 45\pi \times 8 \times \frac{2}{3} \times \sqrt{\frac{2}{3}} \quad \left(=720\pi\sqrt{\frac{2}{3}} - 240\pi\sqrt{\frac{2}{3}}\right)$ (A1) $= 480\pi\sqrt{\frac{2}{3}}$ (A1) (q = 480) [3 marks]

Satprep.co.



Markscheme

November 2024

Mathematics: analysis and approaches

Standard level

Paper 1



19 pages

© International Baccalaureate Organization 2024

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2024

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2024

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *M0* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more *A* marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award *A0A1A1*.
- Where the markscheme specifies *A3*, *M2 etc.*, do **not** split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this working is incorrect and/or suggests a misunderstanding of the question. This will encourage a uniform approach to marking, with less examiner discretion. Although some candidates may be advantaged for that specific question item, it is likely that these candidates will lose marks elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used **in a subsequent part**. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then

used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	8√2	5.65685 (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)
2.	$\frac{35}{72}$	0.468111 (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)

3 Implied marks

Implied marks appear in **brackets e.g.** (*M1*), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is (*M1*)*A1*, it is possible to award full marks for *their* correct answer, without working being seen. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (e.g. probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).

- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a misread (MR). A candidate should be penalized only once for a particular misread. Use the MR stamp to indicate that this has been a misread and do not award the first mark, even if this is an M mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** ... OR.

7 Alternative forms

Unless the question specifies otherwise, accept equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, **M** marks and intermediate **A** marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "from the use of 3 sf values".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an *A* mark to be awarded, arithmetic should be completed,

and any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be

written as $\frac{5}{2}$. An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate *A* marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".

SECTION A

A1	22	(i)	(a)	1.
A1	20	(ii)		
A1) 30	(iii)		
[3 marks]				

$Q_1 = 26$ and $Q_3 = 38$	(A1)
	$Q_1 = 26$ and $Q_3 = 38$

attempt to subtract their upper and lower quartiles			
Note: Award <i>M1</i> only for correct values, or for values clearly indicated as cal Q1 and Q3.	ndidate's		

38–26 IQR =12

A1

[3 marks] [Total: 6 marks]

2. (a)
$$\frac{1}{2}r^2\theta = 48$$
 OR $\frac{1}{2}r^2(1.5) = 48$ (A1)
attempt to solve their equation to find r or r^2 (M1)
Note: To award the M1, candidate's equation must include r^2 and $\theta = 1.5$, and they
must attempt to isolate r^2 or r .
 $r^2 = 64$
 $r = 8$ (cm) A1
[3 marks]
(b) evidence of summing the two radii and the arc length (M1)
perimeter $= 2r + r\theta$
 $= 16 + 8(1.5)$
 $= 28$ (cm) A1
[2 marks]
Total [5 marks]

3. (a) use of
$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$
 (M1)
 $P(A \cap B) = 0.65 + 0.45 - 0.85$ (or equivalent) (A1)

= 0.25

(b)
$$P(A' \cap B') = 0.15$$
 (may be seen in Venn diagram) (A1)

attempt to substitute their values into $P(A' | B') = \frac{P(A' \cap B')}{P(B')}$ (M1)

$$=\frac{15}{55}\left(=\frac{3}{11}\right)$$

[3 marks] Total [6 marks]



4. **METHOD 1**

attempt to expand $(3n+2)^2 - (3n-2)^2$

Note: Award *M0* for invalid attempts such as $(3n+2)^2 = 9n^2 + 4$.

$$=9n^{2}+12n+4-(9n^{2}-12n+4) \text{ or equivalent}$$
 A1
= 24n OR 12n+12n A1

$$= 12(2n) \text{ OR } \frac{24}{12} = 2 \text{ OR } \frac{24n}{2} = 12n \text{ OR } 12n + 12n = 12(n+n) \text{ (or equivalent)}$$
R1
p is a multiple of 12
R3

so is a multiple of 12

Note: Do not award the R1 unless both A marks have been awarded.

METHOD 2

use of $a^2 - b^2 = (a+b)(a-b)$ where $a = 3n+2, b = 3n-2$	M1
= (3n+2+3n-2)(3n+2-3n+2)	
$=6n \times 4$	A1
= 24 <i>n</i>	A1
=12(2 <i>n</i>) OR $\frac{24n}{12}$ = 2 <i>n</i> OR $\frac{24n}{2}$ = 12 <i>n</i> OR 12 <i>n</i> +12 <i>n</i> = 12(<i>n</i> + <i>n</i>) (or equiv	alent)
	R1
so is a multiple of 12	AG

Note: Do not award the R1 unless both A marks have been awarded.

[4 marks]

М1

– 10 –

(M1)

(M1)

5. attempt to substitute into cosine rule

$$(\cos 2\theta =) \frac{4^2 + 6^2 - 5^2}{2 \times 4 \times 6} \text{ OR } 5^2 = 4^2 + 6^2 - 2 \times 4 \times 6 \times \cos 2\theta$$
$$(\cos 2\theta =) \frac{27}{48} \left(= \frac{9}{16} \right)$$
(A1)

attempt to use $\cos 2\theta = 2\cos^2 \theta - 1$

$$\cos^{2} \theta = \frac{1 + \frac{27}{48}}{2} \left(= \frac{1 + \frac{9}{16}}{2} \right)$$

$$\cos^{2} \theta = \frac{75}{96} \left(= \frac{25}{32} \right)$$

$$\cos \theta = (\pm) \sqrt{\frac{75}{96}} \left(= \sqrt{\frac{25}{32}} = \frac{5}{\sqrt{32}} \right)$$

$$= \frac{5}{4\sqrt{2}}$$

$$= \frac{5\sqrt{2}}{8} (p = 5, q = 8)$$
A1

Note: The final answer must be positive.

[6 marks]

(M1)

6. attempt to use $u_n = u_1 + (n-1)d$ or $S_n = \frac{n}{2} [2u_1 + (n-1)d]$ or $S_n = \frac{n}{2} [u_1 + u_n]$ to set up at least one equation in u_1 and d (M1)

$$16 = u_1 + 9d$$
 and $100 = \frac{25}{2} [2u_1 + 24d]$ (A1)

attempt to solve their two linear equations in u_1 and d simultaneously (must eliminate one variable) (M1)

$$d = -4 (\Longrightarrow u_1 = 52)$$

attempt to solve $u_k = 0$ with their d (or with their d and u_1)

$$\Rightarrow k = 14$$
 A1 [6 marks]



SECTION B

the: Award A1 for correct gradient, A1 for correct y-intercept (b) METHOD 1 Axis of Symmetry axis of symmetry is $\frac{-2+4}{2}(=1)$ equating $\frac{-b}{2a}$ to their axis of symmetry $\frac{-b}{-2}=1$ b=2 METHOD 2 substitution attempt to substitute (-2,0) or (4,0) into g $-(-2)^2 - 2b + 8 = 0$ or $-(4)^2 + 4b + 8 = 0$ b=2 METHOD 3 factored form attempt to write g in factored form g(x) = (x+2)(4-x) $-x^2 + 2x + 8$ b=2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ OR $-2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32$ OR $b^2 + 8b + 16 = b^2 + 32$ (or equivalent, must not contain radical)	A1A1
(b) METHOD 1 Axis of Symmetry axis of symmetry is $\frac{-2+4}{2}(=1)$ equating $\frac{-b}{2a}$ to their axis of symmetry $\frac{-b}{-2}=1$ b=2 METHOD 2 substitution attempt to substitute (-2,0) or (4,0) into g $-(-2)^2 - 2b + 8 = 0$ or $-(4)^2 + 4b + 8 = 0$ b=2 METHOD 3 factored form attempt to write g in factored form g(x) = (x+2)(4-x) $-x^2 + 2x + 8$ b=2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ OR $-2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32$ OR $b^2 + 8b + 16 = b^2 + 32$ (or equivalent, must not contain radical)	
(b) METHOD 1 Axis of Symmetry axis of symmetry is $\frac{-2+4}{2}(=1)$ equating $\frac{-b}{2a}$ to their axis of symmetry $\frac{-b}{-2} = 1$ b=2 METHOD 2 substitution attempt to substitute (-2,0) or (4,0) into g $-(-2)^2 - 2b + 8 = 0$ or $-(4)^2 + 4b + 8 = 0$ b=2 METHOD 3 factored form attempt to write g in factored form g(x) = (x+2)(4-x) $-x^2 + 2x + 8$ b=2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ OR $-2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32$ OR $b^2 + 8b + 16 = b^2 + 32$ (or equivalent, must not contain radical)	marks]
axis of symmetry is $\frac{-2+4}{2}(=1)$ equating $\frac{-b}{2a}$ to their axis of symmetry $\frac{-b}{-2} = 1$ b = 2 METHOD 2 substitution attempt to substitute (-2,0) or (4,0) into g $-(-2)^2 - 2b + 8 = 0$ or $-(4)^2 + 4b + 8 = 0$ b = 2 METHOD 3 factored form attempt to write g in factored form g(x) = (x+2)(4-x) $-x^2 + 2x + 8$ b = 2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ OR $-2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32$ OR $b^2 + 8b + 16 = b^2 + 32$ (or equivalent, must not contain radical)	
equating $\frac{-b}{2a}$ to their axis of symmetry $\frac{-b}{-2} = 1$ b = 2 METHOD 2 substitution attempt to substitute (-2,0) or (4,0) into g $-(-2)^2 - 2b + 8 = 0$ or $-(4)^2 + 4b + 8 = 0$ b = 2 METHOD 3 factored form attempt to write g in factored form g(x) = (x+2)(4-x) $-x^2 + 2x + 8$ b = 2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ OR $-2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32$ OR $b^2 + 8b + 16 = b^2 + 32$ (or equivalent, must not contain radical)	(A1)
$\frac{-b}{-2} = 1$ b = 2 METHOD 2 substitution attempt to substitute (-2,0) or (4,0) into g $-(-2)^2 - 2b + 8 = 0$ or $-(4)^2 + 4b + 8 = 0$ b = 2 METHOD 3 factored form attempt to write g in factored form g(x) = (x+2)(4-x) $-x^2 + 2x + 8$ b = 2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ OR $-2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32$ OR $b^2 + 8b + 16 = b^2 + 32$ (or equivalent, must not contain radical)	(M1)
b = 2 METHOD 2 substitution attempt to substitute (-2,0) or (4,0) into g $-(-2)^{2} - 2b + 8 = 0 \text{ or } -(4)^{2} + 4b + 8 = 0$ $b = 2$ METHOD 3 factored form attempt to write g in factored form $g(x) = (x+2)(4-x)$ $-x^{2} + 2x + 8$ $b = 2$ METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)} OR -2 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)}$ $b^{2} - 16b + 64 = b^{2} + 32 OR b^{2} + 8b + 16 = b^{2} + 32 (\text{or equivalent, must not contain radical)}$	
METHOD 2 substitution attempt to substitute (-2,0) or (4,0) into g $-(-2)^2 - 2b + 8 = 0$ or $-(4)^2 + 4b + 8 = 0$ b = 2 METHOD 3 factored form attempt to write g in factored form g(x) = (x+2)(4-x) $-x^2 + 2x + 8$ b = 2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ OR $-2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32$ OR $b^2 + 8b + 16 = b^2 + 32$ (or equivalent, must not contain radical)	A1
attempt to substitute (-2,0) or (4,0) into g $-(-2)^{2} - 2b + 8 = 0 \text{ or } -(4)^{2} + 4b + 8 = 0$ $b = 2$ METHOD 3 factored form attempt to write g in factored form $g(x) = (x+2)(4-x)$ $-x^{2} + 2x + 8$ $b = 2$ METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)} \text{OR} -2 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)}$ $b^{2} - 16b + 64 = b^{2} + 32 \text{OR} b^{2} + 8b + 16 = b^{2} + 32 \text{(or equivalent, must not contain radical)}$	
$-(-2)^{2} - 2b + 8 = 0 \text{ or } -(4)^{2} + 4b + 8 = 0$ b = 2 METHOD 3 factored form attempt to write g in factored form g(x) = (x+2)(4-x) $-x^{2} + 2x + 8$ b = 2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)} \text{OR} -2 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)}$ $b^{2} - 16b + 64 = b^{2} + 32 \text{OR} b^{2} + 8b + 16 = b^{2} + 32 \text{ (or equivalent, must not contain radical)}$	(M1)
b = 2 METHOD 3 factored form attempt to write <i>g</i> in factored form $g(x) = (x+2)(4-x)$ $-x^{2} + 2x + 8$ $b = 2$ METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)} OR -2 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)}$ $b^{2} - 16b + 64 = b^{2} + 32 OR b^{2} + 8b + 16 = b^{2} + 32 \text{ (or equivalent, must not contain radical)}$	(A1)
METHOD 3 factored form attempt to write g in factored form g(x) = (x+2)(4-x) $-x^2 + 2x + 8$ b = 2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ OR $-2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32$ OR $b^2 + 8b + 16 = b^2 + 32$ (or equivalent, must not contain radical)	A1
attempt to write g in factored form $g(x) = (x+2)(4-x)$ $-x^{2} + 2x + 8$ $b = 2$ METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)} \text{OR} -2 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)}$ $b^{2} - 16b + 64 = b^{2} + 32 \text{OR} b^{2} + 8b + 16 = b^{2} + 32 \text{ (or equivalent, must not contain radical)}}$	
g(x) = (x+2)(4-x) $-x^{2} + 2x + 8$ b = 2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)} \text{OR} -2 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)}$ $b^{2} - 16b + 64 = b^{2} + 32 \text{OR} b^{2} + 8b + 16 = b^{2} + 32 \text{ (or equivalent, must not contain radical)}}$	(M1)
$-x^{2} + 2x + 8$ b = 2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)} \text{OR} -2 = \frac{-b \pm \sqrt{b^{2} - 4(-1)(8)}}{2(-1)}$ $b^{2} - 16b + 64 = b^{2} + 32 \text{OR} b^{2} + 8b + 16 = b^{2} + 32 \text{ (or equivalent, must not contain radical)}}$	
<i>b</i> = 2 METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)} \text{OR} -2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32 \text{OR} b^2 + 8b + 16 = b^2 + 32 \text{ (or equivalent, must not contain radical)}}$	(A1)
METHOD 4 quadratic formula attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)} \text{OR} -2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32 \text{OR} b^2 + 8b + 16 = b^2 + 32 \text{(or equivalent, must not contain radical)}}$	A1
attempt to substitute into quadratic formula and set equal to -2 or 4 $4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)} \text{OR} -2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32 \text{OR} b^2 + 8b + 16 = b^2 + 32 \text{ (or equivalent, must not contain radical)}}$	
$4 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)} \text{OR} -2 = \frac{-b \pm \sqrt{b^2 - 4(-1)(8)}}{2(-1)}$ $b^2 - 16b + 64 = b^2 + 32 \text{OR} b^2 + 8b + 16 = b^2 + 32 \text{ (or equivalent, must not contain radical)}}$	(M1)
$b^2 - 16b + 64 = b^2 + 32$ OR $b^2 + 8b + 16 = b^2 + 32$ (or equivalent, must not contain radical)	
	(A1)
b = 2	A1
[3]	narks]

Question 7 continued.

(c) recognizing to subtract g - f (in correct order) (M1)

$$\int_{0}^{4} (-x^{2} + 2x + 8) - (-2x + 8) dx$$
 A1

$$\int_0^4 \left(-x^2 + 4x \right) \mathrm{d}x \qquad \qquad \mathsf{AG}$$

(d) attempt to integrate (M1)

$$-\frac{1}{3}x^3 + 2x^2(+C)$$
 A1

attempt to substitute limits into their integrated function and find difference (M1)

$\left(-\frac{1}{3} \cdot 4^{3} + 2 \cdot 4^{2}\right) - \left(-\frac{1}{3} \cdot 0^{3} + 2 \cdot 0^{2}\right)$	
$\frac{32}{3}$	A1
	[4 marks]

A1	g'(x) = -2x + 2	(e)
(M1)	attempt to equate their derivative of g to their gradient of f	
A1	-2x + 2 = -2	
A1	<i>x</i> = 2	
A1	<i>y</i> = 8	
[5 marks]		
Total [16 marks]		

(a)	(i) substitution of $x = 8$	8 in part (i) or $x = \frac{1}{4}$ in part (ii)	(<i>M1</i>)
	$\log_2(4 \times 8) OR \log_2$	₂ (32)	
	$\log_2(32) = 5$		A1
	(ii) $f\left(\frac{1}{4}\right) = 0$		A1
			[3 marks]
(b)	swap x and y		(M1)
	$x = \log_2(4y)$ OR $x = 2 + \log_2(4y)$	$\log_2 y$	(114)
	$2^x = 4y$	lentia	(A1)
	$(f^{-1}(x) =) \frac{2^x}{4} (= 2^{x-2})$		A1
			[4 marks]
(c)	$\frac{1}{4}$		A1
			[1 mark]
			continued

8.

Question 8 continued.

(d) METHOD 1

$f(16x^3) = \log_2(4 \times 16x^3)$	
attempt to use addition rule for logs	(M1)
$\log_2 4 + \log_2 16 + \log_2 x^3 \text{OR} \log_2 64 + \log_2 x^3$ (or equivalent)	(A1)
attempt to use exponent property for logarithms	(M1)
$f(16x^3) = 6 + 3\log_2 x$ (or equivalent)	A1

the graph of g must be vertically stretched (dilated) by a scale factor of 3 and then vertically translated (shifted) 6 units upwards. **A2**

METHOD 2 (5 marks maximum)	
$f(16x^3) = \log_2(4 \times 16x^3)$	
attempt to write argument as a power	(M1)
$\log_2(64x^3) = \log_2((4x)^3)$ (or equivalent)	(A1)
attempt to use exponent property for logarithms	(M1)
$f(16x^3) = 3\log_2(4x)$ (or equivalent)	A1
EITHER	
the graph of g must be vertically stretched (dilated) by a scale factor of 3	and
stretched (dilated) horizontally by a scale factor of $rac{1}{4}$.	A1
OR	

the graph of g must be stretched (dilated) horizontally by a scale factor of $\frac{1}{4}$ and vertically stretched (dilated) by a scale factor of 3.

Note: In this method, the final mark is **A1**, as the question specifically asks for a translation and a stretch.

[6 marks] [Total: 14 marks] 9. (a) outer curved surface area is $2\pi(4r)h$ AND inner curved surface area is $2\pi rh$ (A1)

area of each base (top and bottom) is
$$\pi (4r)^2 - \pi r^2$$
 (A1)

$$S = 2\left[\pi(4r)^{2} - \pi r^{2}\right] + 2\pi(4r)h + 2\pi rh$$
 A1

$$=30\pi r^2 + 10\pi rh$$

(b) $30\pi r^2 + 10\pi rh = 240\pi$

attempt to solve their equation for h or rh in terms of r (must isolate h or rh) (M1)

$$h = \frac{240 - 30r^2}{10r} \left(= \frac{24 - 3r^2}{r} \right) \text{OR } rh = \frac{240 - 30r^2}{10} \left(= 24 - 3r^2 \right) \text{ (or equivalent)}$$
A1

$$V = \pi (4r)^2 h - \pi r^2 h \quad \left(= 16\pi r^2 h - \pi r^2 h = 15\pi r^2 h \right)$$

attempt to substitute in for h or rh (M1)

$$V = 15\pi r^2 \left(\frac{24 - 3r^2}{r}\right) \text{ OR } V = 15\pi r \left(\frac{240 - 30r^2}{10}\right) \left(=15\pi r \left(24 - 3r^2\right)\right) \text{ OR}$$

$$384\pi r - 48\pi r^3 - 24\pi r + 3\pi r^3 \text{ A1}$$

$$=360\pi r - 45\pi r^3 \qquad \qquad \mathbf{AG}$$

[6 marks]

(c)
$$\frac{\mathrm{d}V}{\mathrm{d}r} = 360\pi - 135\pi r^2$$
 A1A1

[2 marks]
Question 9 continued.

(d) **METHOD 1** (working with *r*) recognition that (for a maximum) $\frac{dV}{dr} = 0$ **M1**

$$360\pi - 135\pi r^{2} = 0$$

$$r^{2} = \frac{360}{135} \left(= \frac{8}{3} \right)$$

$$r = \sqrt{\frac{360}{135}} \left(= \sqrt{\frac{8}{3}} \right)$$
A1
$$n = 2 \text{ OP } r = 2 \sqrt{\frac{2}{3}}$$
A1

$$p = 2 \text{ OR } r = 2\sqrt{\frac{2}{3}}$$
 A1

METHOD 2 (working with
$$p\sqrt{\frac{2}{3}}$$
)
recognition that (for a maximum) $\frac{dV}{dr} = 0$

$$360\pi - 135\pi \left(p\sqrt{\frac{2}{3}}\right)^2 = 0$$

$$360 - 90p^2 = 0$$

$$p^2 = 4$$

$$p = 2 \text{ OR } r = 2\sqrt{\frac{2}{3}}$$
A1
[3 marks]

[3 marks]

Question 9 continued.

(e) attempt to substitute their value of *r* into $V = 360\pi r - 45\pi r^3$ М1

$$V = 360\pi \times 2\sqrt{\frac{2}{3}} - 45\pi \times \left(2\sqrt{\frac{2}{3}}\right)^{3}$$

= $360\pi \times 2\sqrt{\frac{2}{3}} - 45\pi \times 8 \times \frac{2}{3} \times \sqrt{\frac{2}{3}} \quad \left(=720\pi\sqrt{\frac{2}{3}} - 240\pi\sqrt{\frac{2}{3}}\right)$ (A1)
= $480\pi\sqrt{\frac{2}{3}} \quad (q = 480)$ A1





Markscheme

May 2024

Mathematics: analysis and approaches

Standard level

Paper 1

22 pages



© International Baccalaureate Organization 2024

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2024

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2024

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *M0* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies A3, M2 etc., do not split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this
 working is incorrect and/or suggests a misunderstanding of the question. This will encourage a
 uniform approach to marking, with less examiner discretion. Although some candidates may be
 advantaged for that specific question item, it is likely that these candidates will lose marks elsewhere
 too.
- An exception to the previous rule is when an incorrect answer from further working is used **in a subsequent part**. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	8√2	5.65685 (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)
2.	$\frac{35}{72}$	0.468111… (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)

3 Implied marks

Implied marks appear in **brackets e.g.** (*M1*), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is *(M1)A1*, it is possible to award full marks for *their* correct answer, **without working being seen**. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (MR). A candidate should be penalized only once for a particular misread. Use the MR stamp to indicate that this has been a misread and do not award the first mark, even if this is an M mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** ... OR.

7 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, *M* marks and intermediate *A* marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "*from the use of 3 sf values*".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an *A* mark to be awarded, arithmetic should be completed, and

any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$. An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or

written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate A marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".

Section A

1. (a) valid method to find the common difference (M1)

$$d = \frac{22-10}{2} \text{ OR } 10+2d = 22 \text{ OR } u_1 + d = 10, u_1 + 3d = 22 \text{ OR } u_3 = 16$$

$$d = 6$$
(A1)

$$u_1 = 10-6(=4)$$
(A1)

$$u_n = 4+6(n-1) \text{ OR } u_n = 6n-2$$
(A1)

$$u_n = 4+6(n-1) \text{ OR } u_n = 6n-2$$
(A1)
[2 marks]
Total [4 marks]
2. (a) attempt to form equation for the sum of frequencies=16 or mean=3 (M1)

$$p+q+4+2+3=16(\Rightarrow p+q=7)$$
A1

$$\frac{p+2q+12+8+18}{16} = 3(\Rightarrow p+2q=10) \text{ OR } \frac{p+2q+12+8+18}{9+p+q} = 3(\Rightarrow 2p+q=11) \text{ A1}$$
attempt to eliminate one variable from their equations (M1)

$$p+2(7-p)+38=48 \text{ OR } 2(7-q)+q=11$$

$$p=4 \text{ and } q=3$$
A1
Note: Award M1A0A0M0A1 for $p=4, q=3$ with no working.
[5 marks]

- - 4

A1

(b) mean final score =30

[1 mark] Total [6 marks]

3. (a)
$$\log_{10} 1 - \log_{10} a$$
 OR $\log_{10} a^{-1} = -\log_{10} a$ OR $\log_{10} 10^{-\frac{1}{3}}$ OR $10^x = \frac{1}{10^{\frac{1}{3}}}$ (A1)

$$=-\frac{1}{3}$$

[2 marks]

(b)
$$\frac{\log_{10} a}{\log_{10} 1000}$$
 OR $\frac{1}{3} \log_{1000} 10$ OR $\log_{1000} \sqrt[3]{1000^{\frac{1}{3}}}$ OR $10^{\frac{1}{3}} = 1000^{x} \left(= \left(10^{3}\right)^{x}\right)$ (A1)

 $=\frac{1}{9}$

$$\frac{\log_{10} a}{3} \text{ OR } \frac{1}{3} \log_{1000} 1000^{\frac{1}{3}} \text{ OR } \log_{1000} 1000^{\frac{1}{9}} \text{ OR } 3x = \frac{1}{3}$$
 (A1)

A1

[3 marks] Total [5 marks]

М1

A1

4. (a)
$$2r + r\theta = 10$$
 A1

$$\frac{1}{2}r^2\theta = 6.25$$

attempt to eliminate $\,\theta\,\,$ to obtain an equation in $\,r$

correct intermediate equation in r

$$10 - 2r = \frac{25}{2r} \text{ OR } \frac{10}{r} - 2 = \frac{25}{2r^2} \text{ OR } \frac{1}{2}r^2 \left(\frac{10}{r} - 2\right) = 6.25 \text{ OR } 12.5 + 2r^2 = 10r$$

$$4r^2 - 20r + 25 = 0$$
AG [4 marks]

(b) attempt to solve quadratic by factorizing or use of formula or completing the square (M1)

$$(2r-5)^{2} = 0 \quad \text{OR} \quad r = \frac{20 \pm \sqrt{(-20)^{2} - 4(4)(25)}}{2(4)} \left(= \frac{20 \pm \sqrt{400 - 400}}{8} \right)$$
$$r = \frac{5}{2}$$

attempt to substitute their value of r into their perimeter or area equation (M1)

$$\theta = \frac{10 - 2\left(\frac{5}{2}\right)}{\left(\frac{5}{2}\right)} \text{ or } \theta = \frac{25}{2\left(\frac{5}{2}\right)^2}$$

$$\theta = 2$$
A1

[4 marks] Total [8 marks]

(M1)

5. (a) recognising $\cos x = 2\sin x \cos x$

$$(\cos x \neq 0)$$
 so $\sin x = \frac{1}{2}$ OR one correct value (accept degrees) (A1)

x - coordinates
$$\frac{\pi}{6}$$
 and $\frac{5\pi}{6}$ A1

Note: Award (*M1*)(A1)A0 for solutions of 30° and 150° .

[3 marks]

(b) METHOD 1

attempt to integrate $\pm(\cos x - \sin 2x)$ (M1)

$$\int_{\frac{\pi}{2}}^{\frac{5\pi}{6}} (\cos x - \sin 2x) dx \quad OR \quad \int_{\frac{\pi}{2}}^{\frac{5\pi}{6}} (\cos x - 2\sin x \cos x) dx$$
$$= \left[\sin x + \frac{1}{2} \cos 2x \right]_{\frac{\pi}{2}}^{\frac{5\pi}{6}} \quad OR \quad = \left[\sin x - \sin^2 x \right]_{\frac{\pi}{2}}^{\frac{5\pi}{6}} \qquad A1$$

Note: Award A1 for ± correct integration. Condone incorrect or absent limits up to this point.

attempt to substitute their limits into their integral and subtract M1

$$= \left(\sin\left(\frac{5\pi}{6}\right) + \frac{1}{2}\cos\left(\frac{5\pi}{3}\right)\right) - \left(\sin\left(\frac{\pi}{2}\right) + \frac{1}{2}\cos(\pi)\right) OR \left(\sin\left(\frac{5\pi}{6}\right) - \sin^2\left(\frac{5\pi}{6}\right)\right) - \left(\sin\left(\frac{\pi}{2}\right) - \sin^2\left(\frac{\pi}{2}\right)\right)$$

$$= \left(\frac{1}{2} + \frac{1}{4}\right) - \left(1 - \frac{1}{2}\right) OR = \left(\frac{1}{2} - \frac{1}{4}\right) - (1 - 1)$$

$$\operatorname{area} = \frac{1}{4}$$
A1

Note: Award all corresponding marks as appropriate for finding the area between A and B. Accept work done in degrees.

Question 5 continued

METHOD 2

$$\int_{\frac{\pi}{2}}^{\frac{5\pi}{6}} \cos x \, dx = \left[\sin x\right]_{\frac{\pi}{2}}^{\frac{5\pi}{6}} \quad \text{and} \quad \int_{\frac{\pi}{2}}^{\frac{5\pi}{6}} \sin 2x \, dx = \left[-\frac{1}{2}\cos 2x\right]_{\frac{\pi}{2}}^{\frac{5\pi}{6}}$$

Note: Award A1 for correct integration. Condone incorrect or absent limits up to this point.

attempt to substitute their limits into their integral and subtract (for both integrals) M1

$$\sin\left(\frac{5\pi}{6}\right) - \sin\left(\frac{\pi}{2}\right)$$
 and $-\frac{1}{2}\cos\left(\frac{5\pi}{3}\right) + \frac{1}{2}\cos(\pi)$

attempt to subtract the two integrals in either order (seen anywhere) (M1)

$$\left(\sin\left(\frac{5\pi}{6}\right) - \sin\left(\frac{\pi}{2}\right)\right) - \left(-\frac{1}{2}\cos\left(\frac{5\pi}{3}\right) + \frac{1}{2}\cos\left(\pi\right)\right) \text{ OR } \int_{\frac{\pi}{2}}^{\frac{5\pi}{6}} \cos x \, dx - \int_{\frac{\pi}{2}}^{\frac{5\pi}{6}} \sin 2x \, dx$$
$$= \left(\frac{1}{2} - 1\right) - \left(\frac{1}{4} - \frac{1}{2}\right) \left(= -\frac{1}{4}\right)$$
$$\text{area} = \frac{1}{4}$$

Note: Award all corresponding marks as appropriate for finding the area between A and B. Accept work done in degrees.

[4 marks] Total [7 marks]

A1

6. (a)
$$S_n = \frac{10^n - 1}{9}$$

(a = 10, b = 9) A1

[1 mark]

(b) METHOD 1

$$S_{1} + S_{2} + S_{3} + \dots + S_{n}$$

$$= \frac{10 - 1}{9} + \frac{10^{2} - 1}{9} + \dots + \frac{10^{n} - 1}{9}$$

$$= \frac{10 - 1 + 10^{2} - 1 + 10^{3} - 1 + \dots + 10^{n} - 1}{9} \quad \text{OR} \quad \frac{9(10 - 1 + 10^{2} - 1 + 10^{3} - 1 + \dots + 10^{n} - 1)}{81}$$
(A1)

attempt to use geometric series formula on powers of 10, and collect -1's together M1

$$10+10^2+10^3+\ldots+10^n = \frac{10(10^n-1)}{10-1}$$
 and $-1-1-1\ldots=-n$

$$=\frac{\frac{10(10^{n}-1)}{10-1}-n}{9} \text{ OR } \frac{9\left(\frac{10(10^{n}-1)}{10-1}\right)-9n}{81}$$

Note: Award A1 for any correct intermediate expression.

$$=\frac{10(10^n-1)-9n}{81}$$

Question 6 continued

METHOD 2

attempt to create sum using sigma notation with $\boldsymbol{S}_{\boldsymbol{n}}$ М1

$$\sum_{i=1}^{n} \frac{10^{i} - 1}{9} \left(= \frac{1}{9} \left(\sum_{i=1}^{n} 10^{i} - \sum_{i=1}^{n} 1 \right) \right)$$

$$\sum_{i=1}^{n} 10^{i} = \frac{10(10^{n} - 1)}{9}$$
A1
$$\sum_{i=1}^{n} 1 = n$$
A1
$$= \frac{1}{9} \left(\frac{10(10^{n} - 1)}{9} - n \right) \text{ OR } \frac{1}{9} \left(\frac{10(10^{n} - 1) - 9n}{9} \right)$$
A1
$$= \frac{10(10^{n} - 1) - 9n}{81}$$
AG

AG

[4 marks] Total [5 marks]

- 14 -

Section B

7. (a) (i)
$$\frac{dy}{dx} = 3x^2 - 2x - 1$$
 A1A1
Note: Award A1 for $3x^2 - 2x$ and A1 for -1 .

(ii)
$$\frac{\mathrm{d}^2 y}{\mathrm{d}x^2} = 6x - 2$$

[3 marks]

(b) setting their (quadratic)
$$\frac{dy}{dx}$$
 to 0 and solve using valid method (M1)
 $(3x+1)(x-1)=0$ OR $x = \frac{2\pm\sqrt{(-2)^2-4(3)(-1)}}{2(3)}\left(=\frac{2\pm\sqrt{4+12}}{6}\right)$
 $x = -\frac{1}{3}$ (OR $x = 1$) A1
substituting one of their x values into $\frac{d^2y}{dx^2}$ M1
EITHER
at $x = -\frac{1}{3}$, $\frac{d^2y}{dx^2} = 6\left(-\frac{1}{3}\right) - 2\left(=-4\right)$ <0 so local max R1
OR

at
$$x = 1$$
, $\frac{d^2 y}{dx^2} = 6(1) - 2(=4) > 0$ so local min hence local max at $x = -\frac{1}{3}$ **R1**

Note: Award **R1** only if the previous **M1** has been awarded and there is reference to < 0 or > 0, as appropriate.

(M1)

Question 7 continued

THEN

substituting their *x*-coordinate of A into *y*

$$y = \left(-\frac{1}{3}\right)^3 - \left(-\frac{1}{3}\right)^2 - \left(-\frac{1}{3}\right) + 1$$

$$y = \frac{32}{27}$$

So coordinates of A are $\left(-\frac{1}{3}, \frac{32}{27}\right)$
A1

Note: This (M1)A1 may be awarded independently of the previous M1R1.

[6 marks]

(c)	setting their $\frac{d^2y}{dx^2}$ to 0	(M1)
	6x - 2 = 0	
	$x = \frac{1}{3}$	A1
		[2 marks]
(d)	gradient of tangent = -1	(A1)
	negative reciprocal of their gradient	(M1)
	gradient of normal = 1	
	equation is $y = x + 1$ (accept point/slope form $y - 1 = (x - 0)$)	A1

Note: Do not accept L = x + 1.

[3 marks] Total [14 marks] 8. (a)



Note: These marks may be awarded independently.

(b) $y \neq 4$ (or equivalent)

[5 marks]

A1

A1

A1

A1

[1 mark]

Question 8 continued

(c)
$$2 + \frac{5}{2} \text{ OR } \left(-\frac{1}{2}\right) + 2 \times \frac{5}{2} \text{ OR } \frac{-\frac{1}{2} + p}{2} = 2 \text{ OR } -4 = -p + \frac{1}{2}$$
 A1

$$p = \frac{9}{2}$$

(d) METHOD 1

attempt to substitute both roots to form a quadratic (M1)
EITHER

$$\left(x + \frac{1}{2}\right) \left(x - \frac{9}{2}\right) \text{ OR } x^2 - \left(-\frac{1}{2} + \frac{9}{2}\right) x + \left(-\frac{1}{2} \times \frac{9}{2}\right)$$

$$= x^2 - 4x - \frac{9}{4}$$

$$\left(b = -4, c = -\frac{9}{4}\right)$$

$$A1A1$$

Note: Award **A1** for each correct value. They may be embedded or stated explicitly.

OR

$$(2x+1)(2x-9) = 4\left(x^2 - 4x - \frac{9}{4}\right)$$

b = -4, c = -\frac{9}{4}
A1A1

Note: Award **A1** for each correct value. They must be stated explicitly.

METHOD 2

$$-\frac{b}{2} = 2 \text{ OR } 4 + b = 0 \Longrightarrow b = -4$$

attempt to form a valid equation to find c using their b

$$\left(-\frac{1}{2}\right)^{2} + -4\left(-\frac{1}{2}\right) + c = 0 \text{ OR } \left(\frac{9}{2}\right)^{2} + -4\left(\frac{9}{2}\right) + c = 0$$

$$c = -\frac{9}{4}$$
A1

continued...

(M1)

Question 8 continued

METHOD 3

attempt to form two valid equations in b and c (M1)

$$\left(-\frac{1}{2}\right)^{2} + b\left(-\frac{1}{2}\right) + c = 0, \ \left(\frac{9}{2}\right)^{2} + b\left(\frac{9}{2}\right) + c = 0$$

$$b = -4, c = -\frac{9}{4}$$

A1A1

METHOD 4

attempt to write g(x) in the form $(x-h)^2 + k$ and substitute for x, h and g(x) (M1)

$$\left(-\frac{1}{2}-2\right)^{2}+k=0 \Rightarrow k=-\frac{25}{4}$$

$$(x-2)^{2}-\frac{25}{4}$$

$$=x^{2}-4x-\frac{9}{4}$$

$$\left(b=-4,c=-\frac{9}{4}\right)$$
A1A1

Note: Award A1 for each correct value. They may be embedded or stated explicitly.

[3 marks]

(e) attempt to substitute x = 2 into their g(x) OR

complete the square on their g(x) (may be seen in part (d)) (M1)

$$y = -\frac{25}{4}$$
[2 marks]

Question 8 continued





3 solutions A1
Note: Exception to *FT*: If the candidate's graph in part (a) is incorrect, the *M1* may be awarded for a

sketch of their graph from part (a) and g(x) . Do not award the final **A1** in this case.

[2 marks] Total [14 marks]

(M1)

(M1)

9. (a) evidence of understanding that there are now 3R and 2B

$$p = \frac{3}{5}, q = \frac{2}{5}$$
 A1

(b) attempt to add two products (M1)

$$P(same) = P(RR \text{ or } BB) = \frac{3}{4} \times \frac{4}{5} + \frac{1}{4} \times \frac{2}{5}$$

$$= \frac{14}{20}$$

$$= \frac{7}{10}$$
AG [2 marks]

(c) attempt to use conditional probability formula in context (M1) $P(RR|same) = \frac{P(RR)}{P(same)}$

Note: Award *M0* if candidate only writes P(A|B) formula and nothing else.

$$=\frac{\left(\frac{12}{20}\right)}{\left(\frac{14}{20}\right)}$$

$$=\frac{12}{14}\left(=\frac{6}{7}\right)$$
[3 marks]

Question 9 continued

.

(d)
$$a = \frac{6}{20} \left(= \frac{3}{10} \right), \ b = \frac{12}{20} \left(= \frac{6}{10} \right)$$
 A1A1

[2 marks]

(e) attempt to use the formula for E(X)(M1)

$$E(X) = 0 \times \frac{1}{10} + 1 \times \frac{6}{20} + 2 \times \frac{12}{20}$$
$$= \frac{30}{20} \left(= \frac{3}{2} \right)$$
A1

A1

[2 marks]

[1 mark]

(f)
$$\frac{1}{6}$$

METHOD 1 (g)

$$P(n-1 \text{ reds }) = \frac{3}{4} \times \frac{4}{5} \times ... \times \frac{n+1}{n+2} \left(= \frac{3}{n+2} \right)$$
(A1)

P(next one blue) = $\frac{1}{n+3}$	(A1)

 $P(n-1 \text{ reds then } 1 \text{ blue}) = P(n-1 \text{ reds}) \times P(\text{next one blue})$ (M1) 3 1 3

$$\frac{3}{n+2} \times \frac{1}{n+3} = \frac{3}{56}$$
(A1)
(n+2)(n+3) = 56
n = 5
A1

$$n=5$$

Note: If no working shown, award *M1A0A0A0A1* for n = 5.

Question 9 continued

METHOD 2

Let *X* be the number of selections in total made when first blue picked

attempt to establish pattern for X = 1, 2, 3, ... with at least 3 cases (M1)

$$P(X=1) = \frac{1}{4}$$
 and $P(\text{second pick}) = \frac{3}{4} \times \frac{1}{5}$ (A1)

$$P(X=3)\left(=\frac{3}{4}\times\frac{4}{5}\times\frac{1}{6}\right)=\frac{3}{5}\times\frac{1}{6}$$
(A1)

$$P(X=5) = \frac{3}{7} \times \frac{1}{8} \left(= \frac{3}{56} \right)$$
(A1)

so n = 5

METHOD 3

P(next one blue) = $\frac{1}{n+3}$	(A1)
------------------------------------	------

recognising $P(n-1 \text{ R then } 1B) = P(n-1 \text{ R}) \times P(\text{next one } B)$ OR $\frac{3}{4} \times \frac{4}{5} \times ... \times \frac{1}{n+3}$ (M1)

$$\frac{3}{4} \times \frac{4}{5} \times \frac{5}{6} \times \frac{6}{7} \times \frac{1}{8} \left(= \frac{3}{56} \right)$$
(A1)(A1)

Note: Award **A1** for $\frac{3}{4} \times \frac{4}{5} \times \frac{5}{6} \times \frac{6}{7}$ (seen anywhere) and **A1** for $\times \frac{1}{8}$.

so n = 5

A1

A1

[5 marks] Total [17 marks]



Markscheme

May 2024

Mathematics: analysis and approaches

Standard level

Paper 1

27 pages



© International Baccalaureate Organization 2024

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2024

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2024

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *M0* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies A3, M2 etc., do not split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this
 working is incorrect and/or suggests a misunderstanding of the question. This will encourage a
 uniform approach to marking, with less examiner discretion. Although some candidates may be
 advantaged for that specific question item, it is likely that these candidates will lose marks
 elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used **in a subsequent part**. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	8√2	5.65685 (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)
2.	$\frac{35}{72}$	0.468111… (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)

3 Implied marks

Implied marks appear in **brackets e.g.** (*M1*), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is *(M1)A1*, it is possible to award full marks for *their* correct answer, **without working being seen.** For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (*MR*). A candidate should be penalized only once for a particular misread. Use the *MR* stamp to indicate that this has been a misread and do not award the first mark, even if this is an *M* mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** . . . **OR**.

7 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, **M** marks and intermediate **A** marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "*from the use of 3 sf values*".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an *A* mark to be awarded, arithmetic should be completed, and

any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$. An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or 5 10

written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate A marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".

SECT	ION A
------	-------

1.	(a)	(i)	f(4) = 1	A1
		(ii)	$f \circ f(4) = 3$	A1
		(iii)	$f^{-1}(3) = 1$	A1
				[3 marks]

(b)



curve passes through (2,2) OR through (1, 4) and (3, 1)

Note: Do not award the second mark unless the first mark has been awarded.(Do not award **A0A1**).

[2 marks]

A1

Total [5 marks]

(M1)

2.	$\tan^{-1}1 = 45^{\circ}$ or equivalent	(A1)

attempt to equate $2x - 5^{\circ}$ to their reference angle

Note: Do not accept $2x - 5^\circ = 1$.

$$2x - 5^{\circ} = 45^{\circ}, (225^{\circ})$$

 $x = 25^{\circ}, 115^{\circ}$

A1A1

Note: Do not award the final A1 if any additional solutions are seen.





$$(3m-1)(m+2) = 0 \text{ OR } m = \frac{-5 \pm \sqrt{25+24}}{6} \text{ (or equivalent)}$$
 (A1)

$$m = \frac{1}{3}$$
, $m = -2$ **A1**

[3	marks]
----	--------

(M1)

A1

(b) setting their *m*-value(s) = 3^x OR recognising a quadratic in 3^x

$$3^{x} = \frac{1}{3}$$
 (or $3^{x} = -2$) OR $3 \times (3^{x})^{2} + 5 \times 3^{x} - 2 = 0$

$$x = -1$$

Note: Award the final **A1** if candidate's answer includes x = -1 and $x = \log_3(-2)$. Award **A0** if other incorrect answers are given.

[2 marks]

Total [5 marks]

4. (a) (i)
$$\left(\frac{9}{2}, \frac{3\sqrt{3}}{2}\right)$$
 (accept $x = \frac{9}{2}$ and $y = \frac{3\sqrt{3}}{2}$) A1

(ii) using
$$m = \frac{\text{changein } y}{\text{changein } x}$$
 with their midpoint OR gradient perpendicular to AC

$$OR \ m = \tan 30^{\circ} \tag{M1}$$

$$m = \frac{\sqrt{3}}{3} \tag{A1}$$

$$y = \frac{\sqrt{3}}{3}x$$
 OR $y - \frac{3\sqrt{3}}{2} = \frac{\sqrt{3}}{3}\left(x - \frac{9}{2}\right)$ (must be written as an equation) A1

(b) substituting
$$x = 6$$
 into their equation (M1)
so at B $y=2\sqrt{3}$ (A1)
area of triangle OAB $=\frac{1}{2} \times 6 \times 2\sqrt{3} = 6\sqrt{3}$

area of quadrilateral OABC = $12\sqrt{3}$

A1

[3 marks]

Total [7 marks]





Question 5 continued.

(b) (i) multiplying the two correct branches (A1)

$$(1-k)\left(1-\frac{k}{2}\right)$$

attempt to expand and equate to $\frac{5}{9}$ (M1)
 $1-k-\frac{k}{2}+\frac{k^2}{2}=\frac{5}{9}$
 $18-18k-9k+9k^2=10$ OR $\frac{k^2}{2}-\frac{3k}{2}+\frac{4}{9}=0$ OR $\frac{9k^2}{2}-\frac{27k}{2}+4=0$ A1
 $9k^2-27k+8=0$ AG
(ii) $(k=\frac{1}{3}$ is the only valid solution as) $\frac{8}{3} > 1$ R1

Note: Accept any valid reasoning indicating that any probability cannot be greater than 1 and/or probability cannot be less than 0.

[4 marks]

Total [6 marks]
6.	(a)	$y = \frac{2}{3}$ (must be written as equation with $y = $,)	A1
----	-----	---	----

[1 mark]

(ii) **EITHER**

$$\frac{2(x+3)}{3(x+2)} = mx+1$$

attempt to expand to obtain a quadratic equation	(M1)
$2x + 6 = 3mx^2 + 6mx + 3x + 6$	
$3mx^{2} + (6m+1)x = 0$ OR $3mx^{2} + 6mx + x = 0$	A1
recognition that discriminant $\Delta = 0$ for one solution	(M1)
$\left(6m+1\right)^2=0$	

Question 6 continued.

OR

$$\frac{2(x+3)}{3(x+2)} = mx+1$$

attempt to ex	pand to obtain a	quadratic equation	(M1)
			•

$$2x + 6 = 3mx^{2} + 6mx + 3x + 6$$

$$3mx^{2} + (6m+1)x = 0 \quad \text{OR} \quad 3mx^{2} + 6mx + x = 0$$

attempt to solve their quadratic for
$$x$$
 and equating their solutions (M1)

$$x(3mx+6m+1)=0$$

$$x = 0 \text{ OR } x = -\frac{6m+1}{3m}(=0)$$

 $-\frac{6m+1}{3m} = 0$

OR

attempt to find f'(x) using the quotient rule

(M1)

$$f'(x) = \frac{2}{3} \left(\frac{(x+2) - (x+3)}{(x+2)^2} \right) = \left(\frac{-2}{3(x+2)^2} \right) \text{ OR } \frac{2(3x+6) - 3(2x+6)}{(3x+6)^2} \text{ or}$$

equivalent A1

equivalent

recognition that *m* is the derivative of f(x) at x = 0(**M1**)

THEN

$$\Rightarrow m = -\frac{1}{6}$$

Question 6 continued.

(iii)

Note: In this part, FT may be awarded only for values of *m* between -1 and 0.

$$-\frac{1}{6} < m < 0$$

A2

Note: Award **A1** for only $m > -\frac{1}{6}$. Award **A1** for only m < 0.





SECTION B

7. (a)
$$12b = 2\pi$$
 OR $(b =)\frac{2\pi}{12}$ OR $12 = \frac{2\pi}{b}$ A1
 $b = \frac{\pi}{6}$ AG

[1 mark]



(c) (i) attempt to substitute
$$x = 5$$
 into $g(x)$ (M1)

$$g(5) = 3.5 \sin \frac{5\pi}{6} + 11$$

$$\sin \frac{5\pi}{6} = \frac{1}{2}$$

$$g(5) = 3.5 \times \frac{1}{2} + 11$$
(A1)

$$g(5) = 12.75\left(=\frac{51}{4}\right)$$
 A1

(ii) METHOD 1 (finding maximum temperature)

considering the maximum value of $\sin \frac{\pi}{6} x (=1)$ OR g'(x) = 0 at maximum

OR maximum = vertical shift + amplitude (may be seen on a graph) (M1)

 $g_{\text{max}} = 3.5 + 11$ OR $\frac{\pi}{6} \cdot 3.5 \cos\left(\frac{\pi}{6}x\right) = 0$ OR x = 3

 $g_{\rm max} = 14.5$ A1

14.5 < 15 (hence the midday water temperature is never warm enough for Alex to swim)

Note: Do not award the R mark unless the previous marks been awarded (Do not award **M1A0R1** or **M0A0R1**).

Worded conclusions are acceptable for the R1, as long as the reasoning is clear that the water does not reach 15° , so not warm enough for Alex.

continued...

R1

Question 7 continued.

METHOD 2 (working with inequality)

$$3.5\sin\left(\frac{\pi}{6}x\right) + 11 \ge 15\tag{M1}$$

$$\sin\!\left(\frac{\pi}{6}x\right) \ge \frac{8}{7}$$

sine values can never be greater than 1 (hence the midday water temperature is never warm enough for Alex to swim)

Note: Do not award the R mark unless the previous marks been awarded (Do not award *M1A0R1* or *M0A0R1*).

If candidate works with an equation throughout, the *M1* and *A1* may be awarded, if appropriate. A correct inequality is required for the *R1* to be awarded.

[6 marks]

R1

Question 7 continued.

(d) **EITHER**

attempt to find
$$0.7 f(x)$$
 OR $0.7 f(x) + q$ (M1)
 $0.7 f(x) = 3.5 \sin \frac{\pi}{6} x + 10.5$ OR $0.7 f(x) + q = 3.5 \sin \frac{\pi}{6} x + 10.5 + q$ OR
 $10.5 + q = 11$ (A1)

OR

attempt to find $0.7 f(x)$ for a particular value of x	(<i>M1</i>)
eg maximum 20×0.7=14	(A1)

THEN

q = 0.5

A1

[3 marks]

Total [12 marks]

8.

(a) (i)

METHOD 1

$$x^{2} + 2x + 2 = (x+1)^{2} + 1$$
 A1
 $(x+1)^{2} \ge 0$ R1

$$(x+1)^2+1>0$$
 AG

METHOD 2

discriminant $\Delta = 4 - 8(= -$	4) A1
-----------------------------------	-------

 $\Delta < 0$ and concave up OR $\Delta < 0$ and coefficient of $x^2 > 0$ (may be seen in diagram)

hence $x^2 + 2x + 2 > 0$

(ii) x > -1 (accept $x \ge -1$)

A1

R1

AG

[3 marks]

(M1)

Question 8 continued.

(b) (i)
$$x = -1$$
 A1

$$f''(x) = \frac{2(x^2 + 2x + 2) - (2x + 2)(2x + 2)}{(x^2 + 2x + 2)^2}$$
A1A1

Note: Award *A1* for first term in numerator, *A1* for second term in numerator (including the minus sign). If the numerator is correct, but the denominator is incorrect or missing, award *M1A1A0*.

$$f''(x) = \frac{2x^2 + 4x + 4 - 4x^2 - 8x - 4}{(x^2 + 2x + 2)^2} \quad \text{OR} \quad f''(x) = \frac{2x^2 + 4x + 4 - (4x^2 + 8x + 4)}{(x^2 + 2x + 2)^2} \quad \text{A1}$$

$$f''(x) = \frac{-2x^2 - 4x}{(x^2 + 2x + 2)^2}$$

AG

Question 8 continued.

Note: Do not award AOR1.

(iii) substituting x = -1 into f''(x), clearly leading to positive numerator **A1** $f''(-1) = \left(\frac{-2+4}{2}\right)(-2)$

$$f''(-1) = \left(\frac{1}{1}\right) (=2)$$

 $f''(-1) > 0$ **R1**

therefore this is a local minimum point

AG

[7 marks]



Question 8 continued.

J

(c) recognition to integrate
$$f'(x)$$
 (M1)
 $f(x) = \int \frac{2x+2}{x^2+2x+2} dx$ OR $\int \frac{1}{u} du$ with $u = x^2 + 2x + 2$

$$f(x) = \ln(x^2 + 2x + 2)(+c)$$
 A1

using initial conditions $x = 2, y = 3 + \ln 10$ to find *c* (M1)

$$f(2) = \ln(10) + c = 3 + \ln(10) \Longrightarrow c = 3$$

$$f(x) = \ln(x^2 + 2x + 2) + 3$$
 A1

[4 marks]

(d)
$$f'(2) = \frac{6}{10}$$
 A1

attempt to take the negative reciprocal of their f'(2) M1

gradient of normal $= -\frac{10}{6}$

$$y - (\ln 10 + 3) = -\frac{5}{3}(x - 2) \qquad \left(y = -\frac{5}{3}x + \frac{19}{3} + \ln 10\right) \quad \left(\frac{y - \ln 10 - 3}{x - 2} = -\frac{5}{3}\right)$$

[3 marks]

Total [17 marks]

A1

(a) attempt to find a difference (M1)

$$d = p - a, 2d = q - a, d = q - p$$
 OR $p = a + d, q = a + 2d, q = p + d$

correct equation

9.

$$p-a = q-p$$
 OR $q-a = 2(p-a)$ OR $p = \frac{a+q}{2}$ (or equivalent)
 $2p-q = a$ AG

[2 marks]

A1

$$r = \frac{s}{a}, r^2 = \frac{t}{a}, r = \frac{t}{s}$$
 OR $s = ar, t = ar^2, t = sr$

correct equation

$$\left(\frac{s}{a}\right)^2 = \frac{t}{a}$$
 OR $\frac{s}{a} = \frac{t}{s}$ (or equivalent)
 $s^2 = at$
[2 marks]
continued...

Question 9 continued

(c) **EITHER**

$$2p-1=s^2$$
 (or equivalent) **A1**

$$(s^2 > 0) \Rightarrow 2p - 1 > 0 \text{ OR } s = \sqrt{2p - 1} \Rightarrow 2p - 1 > 0 \text{ OR } p = \frac{s^2 + 1}{2} \text{ (and } s^2 > 0 \text{)}$$

OR

$$2p-1=a$$
 and $s^2=a$ A1

$$(s^2 > 0, so) \ a > 0 \implies 2p-1 > 0 \ OR \ p = \frac{a+1}{2} \text{ and } a > 0$$

$$\Rightarrow p > \frac{1}{2}$$

AG

Note: Do not award AOR1.



Question 9 continued

(d) (i)
$$9,5,1,-3$$
 A1A1
Note: Award A1 for each of 2nd term and 4th term
(ii) $9,3,1,\frac{1}{3}$ A1A1
Note: Award A1 for each of 2nd term and 4th term
[4 marks]
(e) (i) attempt to find the difference between two consecutive terms
 $d = u_2 - u_1 = 5 + \ln 3 - 9 - \ln 9$ OR $d = u_3 - u_2 = 1 + \ln 1 - 5 - \ln 3$
 $\ln 9 = 2\ln 3$ OR $\ln 1 = 0$ OR $\ln 3 - \ln 9 = \ln \frac{1}{3} (= \ln 3^{-1} = -\ln 3)$ (seen anywhere) (A1)

$$d = -4 - \ln 3$$
 A1

Question 9 continued.

(ii) METHOD 1

attempt to substitute first term and their common difference into $S_{\rm 10}$ (M1)

$$\frac{10}{2} \left(2(9 + \ln 9) + 9(-4 - \ln 3) \right) \text{OR} \ \frac{10}{2} \left(2(9 + 2\ln 3) + 9(-4 - \ln 3) \right) \text{ (or equivalent)}$$

$$=5(-18-5\ln 3)$$
 (or equivalent in terms of $\ln 3$) A1

$$\sum_{i=1}^{10} u_i = -90 - 25 \ln 3$$

METHOD 2

$$u_{10} = 9 + \ln 9 + 9(-4 - \ln 3)(= -27 + \ln 9 - 9\ln 3)$$

attempt to substitute first term and their u_{10} into S_{10} (M1)

$$\frac{10}{2} (2(9+\ln 9)+9(-4-\ln 3)) \text{ OR } \frac{10}{2} (9+\ln 9-27+\ln 9-9\ln 3) \text{ OR}$$

$$\frac{10}{2} \left(2(9+2\ln 3)+9(-4-\ln 3) \right) \text{ OR } \frac{10}{2} \left(9+\ln 9-27-7\ln 3 \right) \text{ (or equivalent)}$$
 A1

 $=5(-18-5\ln 3)$ (or equivalent in terms of $\ln 3$) A1

$$\sum_{i=1}^{10} u_i = -90 - 25 \ln 3$$
 AG

[6 marks]

Total [16 marks]



Markscheme

November 2023

Mathematics: analysis and approaches

Standard level

Paper 1 TZ2

19 pages



© International Baccalaureate Organization 2023

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2023

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2023

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *M0* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where **M** and **A** marks are noted on the same line, *e.g.* **M1A1**, this usually means **M1** for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and **A1** for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies A3, M2 etc., do not split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this
 working is incorrect and/or suggests a misunderstanding of the question. This will encourage a
 uniform approach to marking, with less examiner discretion. Although some candidates may be
 advantaged for that specific question item, it is likely that these candidates will lose marks
 elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used **in a subsequent part**. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	8√2	5.65685 (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)
2.	$\frac{35}{72}$	0.468111… (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)

3 Implied marks

Implied marks appear in **brackets e.g.** (*M1*) and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is *(M1)A1*, it is possible to award full marks for *their* correct answer, **without working being seen.** For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (*MR*). A candidate should be penalized only once for a particular misread. Use the *MR* stamp to indicate that this has been a misread and do not award the first mark, even if this is an *M* mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** ... OR.

7 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, M marks and intermediate A marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come *"from the use of 3 sf values"*.

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an **A** mark to be awarded, arithmetic should be completed, and any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$. An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or written as $\frac{5}{2}$.

However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate A marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".

Section A

1. (a)
$$a=5$$

(b) (i) period = π
(ii) $b = \frac{2\pi}{\pi} \text{ OR } \pi = \frac{2\pi}{b}$
(A1)
 $= 2$
(A1)
 $= 2$
(A1)
 $f(\pi) = 5\cos(\frac{\pi}{3})$
 $\cos(\frac{\pi}{3}) = \frac{1}{2}$
(A1)
 $f(\pi) = 5\cos(\frac{\pi}{3})$
 $\cos(\frac{\pi}{3}) = \frac{1}{2}$
(A1)
 $f(\pi) = 5\cos(\frac{\pi}{3})$
 $\cos(\frac{\pi}{3}) = \frac{1}{2}$
(A1)
 $f(\pi) = 5\cos(\frac{\pi}{3})$
(A1)
(A1

Total [7 marks]

2. (a) attempt to form $(g \circ f)(x)$ (M1)

$$((g \circ f)(x)) = (x-3)^2 + k^2 \quad (=x^2 - 6x + 9 + k^2)$$
 A1

[2 marks]

(b) substituting x = 2 into their $(g \circ f)(x)$ and setting their expression =10 (M1) $(2-3)^2 + k^2 = 10$ OR $2^2 - 6(2) + 9 + k^2 = 10$ $k^2 = 9$ $k = \pm 3$ (A1) and setting their expression =10 (M1) and setting their expression =10 (M1) (and setting their expression =10 (M1)(and setting their expression

3. (a) $(P(A \cup B) =)0.65 + 0.75 - 0.6 \text{ OR } 0.05 + 0.6 + 0.15$ (A1) = 0.8 [2 marks]

(b) recognition that $A' \cap B' = (A \cup B)'$ OR $A' \cap B' = 1 - A \cup B$ (region/value may be seen in a correctly shaded/labeled Venn diagram) (M1) (=1-0.8)= 0.2 A1 Note: For the final mark, 0.2 must be stated as the candidate's answer, or labeled as $P(A' \cap B')$ in their Venn diagram. Just seeing an unlabeled 0.2

in the correct region of their diagram earns M1A0.

[2 marks] Total [4 marks]

(M1)

4. (a) METHOD 1

attempt to form at least one equation, using e	either S_4 or S_5	(M1)
--	-----------------------	------

$$65 = 25p - 5q$$
 $(13 = 5p - q)$ and $40 = 16p - 4q$ $(10 = 4p - q)$ (A1)

valid attempt to solve simultaneous linear equations in p and q by substituting or eliminating one of the variables.

$$p = 3, q = 2$$
 A1A1

Note: If candidate does not explicitly state their values of *p* and *q*, but gives $S_n = 3n^2 - 2n$, award final two marks as **A1A0**.

METHOD 2

attempt to form at least one equation, using either S_4 or S_5 (M1)

$$65 = \frac{5}{2}(2u_1 + 4d) \quad (26 = 2u_1 + 4d) \text{ and } 40 = 2(2u_1 + 3d) \quad (20 = 2u_1 + 3d) \quad (A1)$$

valid attempt to solve simultaneous linear equations in u_1 and d by substituting or eliminating one of the variables.

$$u_1 = 1, \ d = 6$$

 $S_n = \frac{n}{2} (2 + 6(n - 1)) = 3n^2 - 2n$

$$p = 3$$
 and $q = 2$ A1
Note: If candidate does not explicitly state their values of p and q , do not award the final mark.

[5 marks]

(M1)

(b)
$$u_5 = S_5 - S_4$$
 OR substituting their values of u_1 and d into $u_5 = u_1 + 4d$
OR substituting their value of u_1 into $65 = \frac{5}{2}(u_1 + u_5)$ (M1)

$$(u_5 =)65 - 40$$
 OR $(u_5 =)1 + 4 \times 6$ OR $65 = \frac{5}{2}(1 + u_5)$
= 25
[2 marks]
Total [7 marks]

5. METHOD 1 EITHER

attempt to use Pythagoras' theorem in a right-angled triangle. (M1)

$$\left(\sqrt{5^2 - 1^2}\right) = \sqrt{24} \tag{A1}$$

OR

attempt to use the Pythagorean identity $\cos^2 \alpha + \sin^2 \alpha = 1$ (M1)

$$\sin^2 \mathbf{B} \hat{\mathbf{A}} \mathbf{C} = 1 - \left(\frac{1}{5}\right)^2 \tag{A1}$$

THEN

 $\sin B\hat{A}C = \frac{\sqrt{24}}{5} \text{ (may be seen in area formula)}$ attempt to use 'Area = $\frac{1}{2}ab\sin C'$ (must include their calculated value of $\sin B\hat{A}C$) (M1) = $\frac{1}{2} \times 10 \times \sqrt{6} \times \frac{\sqrt{24}}{5}$ (A1) = 12 (cm²) A1 [6 marks] continued...

Question 5 continued

METHOD 2

attempt to find perpendicular height of triangle BAC	(M1)
EITHER	

height =
$$\sqrt{6} \times \sin \hat{BAC}$$

attempt to use the Pythagorean identity $\cos^2 \alpha + \sin^2 \alpha = 1$ (M1)

height =
$$\sqrt{6} \times \sqrt{1 - \left(\frac{1}{5}\right)^2}$$
 (A1)

$$=\sqrt{6} \times \frac{\sqrt{24}}{5} \left(=\frac{12}{5}\right)$$

OR

ı. √6	(64)
$ad_{a}cent =$	(A1)
5	()

attempt to use Pythagoras' theorem in a right-angled triangle. (M1)

height =
$$\sqrt{6 - \frac{6}{25}} \left(= \frac{12}{5} \right)$$
 (may be seen in area formula) (A1)

THEN

attempt to use 'Area = $\frac{1}{2}$ base×height' (must include their calculated value for height) (M1)

$$=\frac{1}{2} \times 10 \times \frac{12}{5}$$
$$= 12 \text{ (cm}^2)$$

[6 marks]

(M1)

6. attempt to apply binomial expansion

 $(1+kx)^n = 1 + {}^nC_1kx + {}^nC_2k^2x^2 + \dots$ OR ${}^nC_1k = 12$ OR ${}^nC_2 = 28$

$$nk = 12$$
 (A1)

$$\frac{n(n-1)}{2} = 28 \quad \text{OR} \quad \frac{n!}{(n-2)!2!} = 28 \tag{A1}$$

$$n^2 - n - 56 = 0$$
 OR $n(n-1) = 56$

valid attempt to solve

(**M1)**

(n-8)(n+7) = 0 OR 8(8-1) = 56 OR finding correct value in Pascal's triangle

$$\Rightarrow n = 8$$

$$\Rightarrow k = \frac{3}{2}$$

Note: If candidate finds n = 8 with no working shown, award **M1A0A0M1A1A0**. If candidate finds n = 8 and $k = \frac{3}{2}$ with no working shown, award **M1A0A0M1A1A1**.

[6 marks]

Section **B**

(i)	p = 9	A1
(ii)	$600 < n \le 800$	A1
te:Awa	ard A0 if candidate answers 700.	
		[2 marks]
(i)	median = 600	A1
(ii)	80% of $800 = 640$	(A1)
	40 (performances less than 80% of tickets sold)	(A1)
	20 (performances)	A1
		[4 marks]
-	(i) (ii) te: Awa (i) (ii)	 (i) p = 9 (ii) 600 < n ≤ 800 te:Award <i>A0</i> if candidate answers 700. (i) median = 600 (ii) 80% of 800 = 640 40 (performances less than 80% of tickets sold) 20 (performances)

any reasonable answer which suggests a biased sample (must include reason, (c) (i) do not accept reasons such as "sample size is too small", or answers that simply say "not representative of entire audience" without a valid reason) A1

> eg likely to come from the same part of the theatre OR be part of same group OR be from priority seating OR it is convenience sampling

(ii) every 20 th person	A1A1
Note: Award A1 for recognizing that sampling occurs at regular intervals eg	
Award A1 for interval length is 20.	
(iii) quota (sampling method)	A1

quota (sampling method) (iii)

[4 marks]

Question 7 continued

(M1)	75% (of 36000 spent between \$3 and \$25)	(i)	(d)
A1	= 27000		
A1	<i>a</i> = 7	(ii)	
[3 marks]			

(e)	(i)	METHOD 1	
		old mean is 600 (tickets)	(A1)
		recognising new mean is old mean + 17	(M1)
		600+17	
		=617 (tickets)	A1
		METHOD 2	
		new total number of tickets = $36000 + 60 \times 17(= 37020)$	(A1)
		new mean = $\frac{36000 + 60 \times 17}{60} (= \frac{37020}{60})$	(M1)
		=617 (tickets)	A1
	(ii)	no effect on the variance	A1
			[4 marks]
			Total [17 marks]

8. (a) *x* = 0

A1

[1 mark]

(b) (i) setting
$$\ln(2x-9) = 2\ln x - \ln d$$
 M1

attempt to use power rule (M1)
$$2\ln x = \ln x^2$$
 (seen anywhere)

$$\ln(2x-9) = \ln\frac{x^2}{d} \text{ OR } \ln\frac{x^2}{2x-9} = \ln d \text{ OR } \ln(2x-9)d = \ln x^2$$

$$\frac{x^2}{d} = 2x - 9 \text{ OR } \frac{x^2}{2x - 9} = d \text{ OR } (2x - 9)d = x^2$$
A1

$$x^2 - 2dx + 9d = 0$$
 AG

- (ii) discriminant = $(-2d)^2 4 \times 1 \times 9d$ (A1) recognizing discriminant > 0 (M1) $(-2d)^2 - 4 \times 1 \times 9d > 0$ OR $(2d)^2 - 4 \times 9d > 0$ OR $4d^2 - 36d > 0$ A1 $d^2 - 9d > 0$ AG
- (iii) setting d(d-9) > 0 OR d(d-9) = 0 OR sketch graph OR sign test OR $d^2 > 9d$ (M1) d < 0 or d > 9, but $d \in \mathbb{R}^+$ d > 9 (or]9, ∞ [) A1

[9 marks]

Question 8 continued

(c)
$$x^2 - 20x + 90 (= 0)$$
 A1

 attempting to solve their 3 term quadratic equation
 (M1)

 $((x-10)^2 - 10 = 0)$ or $(x=) \frac{20 \pm \sqrt{(-20)^2 - 4 \times 1 \times 90}}{2})$
 (M1)

 $x = 10 - \sqrt{10} (= p)$ or $x = 10 + \sqrt{10} (= q)$
 (A1)

 subtracting their values of x
 (M1)

 $(a = 2, b = 10)$
 A1

 Note: Accept $1\sqrt{40}$ OR $\sqrt{40}$.

[5 marks] Total [15 marks]

(M1)

9. (a) attempt to use either the quotient or product rule

$$\frac{8(x^{2}+1)^{3}-8x\times3\times2x(x^{2}+1)^{2}}{(x^{2}+1)^{6}} \quad \text{OR} \quad 8(x^{2}+1)^{-3}+8x\times(-3)\times2x(x^{2}+1)^{-4} \qquad \textbf{A1A1}$$

Note: Award **A1** for correctly applying chain rule to $(x^2 + 1)^3$ and **A1** for everything else correct.

$$=\frac{8(x^{2}+1-6x^{2})}{(x^{2}+1)^{4}} \text{ OR } \frac{8(x^{2}+1)^{2}(x^{2}+1-6x^{2})}{(x^{2}+1)^{6}} \text{ OR } \frac{8-40x^{2}}{(x^{2}+1)^{4}} \text{ OR } \frac{8x^{2}+8-48x^{2}}{(x^{2}+1)^{4}}$$

 $=\frac{8(1-5x^{2})}{(x^{2}+1)^{4}}$

AG

[4 marks]

Question 9 continued

(b) **EITHER**

attempts to integrate by substitution using $u = x^2 + 1$ or $u = x^2$ (M1)

$$u = x^{2} + 1 \Longrightarrow \frac{\mathrm{d}u}{\mathrm{d}x} = 2x$$
 OR $u = x^{2} \Longrightarrow \frac{\mathrm{d}u}{\mathrm{d}x} = 2x$

Note: If candidate simply states $u = x^2 + 1$ or $u = x^2$, but does not attempt to substitute into their integral, do not award the *(M1)*.

$$\int \frac{8x}{(x^2+1)^3} dx = \int \frac{4}{u^3} du \quad \text{OR} \quad \int \frac{8x}{(x^2+1)^3} dx = \int \frac{4}{(u+1)^3} du$$
 (A1)

$$= -2u^{-2}(+c) \quad \text{OR} \quad -2(u+1)^{-2}(+c) \tag{A1}$$

OR

attempts to apply integration by inspection

$$4\int \frac{2x}{(x^{2}+1)^{3}} dx$$

= $4 \times \left(-\frac{1}{2}\right) (x^{2}+1)^{-2} (+c)$ (A1)(A1)

Note: Award **A1** for correct power of $(x^2 + 1)$ and **A1** for -

THEN

$$-2\left(x^{2}+1\right)^{-2}+c \text{ OR } -\frac{2}{\left(x^{2}+1\right)^{2}}+c \text{ (final answer must include } +c \text{)}$$

 $\frac{1}{2}$.

[4 marks]

(M1)

Question 9 continued

(c) recognizing
$$g'(x) = f'(x) \Rightarrow g(x) = f(x) + k$$
 (may be seen in diagram/drawing) **A1**

area of *R* is given by subtracting functions *f* and *g* in integral(s) (M1)

$$\pm \int_{0}^{3} k dx \quad OR = \int_{0}^{3} |g - f| dx \quad OR \quad \int_{0}^{3} f(x) + k - f(x) dx \quad OR \quad \int_{0}^{3} f(x) dx - \int_{0}^{3} g(x) dx$$

$$=\pm \left[kx\right]_{0}^{3} \text{ OR } \left[-\frac{2}{\left(x^{2}+1\right)^{2}}+kx\right]_{0}^{3}-\left[-\frac{2}{\left(x^{2}+1\right)^{2}}\right]_{0}^{3} \text{ OR } \left[-\frac{2}{\left(x^{2}+1\right)^{2}}\right]_{0}^{3}-\left[-\frac{2}{\left(x^{2}+1\right)^{2}}+kx\right]_{0}^{3} (A1)$$

$$\pm 3k = \frac{27}{2}$$

$$k = \pm \frac{27}{6} \left(= \pm \frac{9}{2} = \pm 4.5 \right)$$
(A1)

$$g(x) = \frac{8x}{(x^2+1)^3} - \frac{9}{2} \text{ AND } g(x) = \frac{8x}{(x^2+1)^3} + \frac{9}{2} \text{ (accept } f(x) + \frac{9}{2} \text{ AND } f(x) - \frac{9}{2} \text{)} \text{ A1}$$

[5 marks] Total [13 marks]



Markscheme

November 2023

Mathematics: analysis and approaches

Standard level

Paper 1 TZ1

19 pages



© International Baccalaureate Organization 2023

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2023

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2023

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.
Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *M0* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies A3, M2 etc., do not split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this
 working is incorrect and/or suggests a misunderstanding of the question. This will encourage a
 uniform approach to marking, with less examiner discretion. Although some candidates may be
 advantaged for that specific question item, it is likely that these candidates will lose marks
 elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used **in a subsequent part**. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	8√2	5.65685 (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)
2.	$\frac{35}{72}$	0.468111… (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)

3 Implied marks

Implied marks appear in **brackets e.g.** (*M1*), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is *(M1)A1*, it is possible to award full marks for *their* correct answer, **without working being seen.** For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (MR). A candidate should be penalized only once for a particular misread. Use the MR stamp to indicate that this has been a misread and do not award the first mark, even if this is an M mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** ... OR.

7 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, **M** marks and intermediate **A** marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "*from the use of 3 sf values*".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an *A* mark to be awarded, arithmetic should be completed, and

any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$. An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or

written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate A marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".

Section A

1. (a)
$$a = 7$$

(b) (i) period = π
(ii) $b = \frac{2\pi}{\pi} \text{ OR } \pi = \frac{2\pi}{b}$
(A1)
 $= 2$
(A1)
 $= 2$
(A1)
 $f(\pi) = 2 \text{ for } \pi$
(A1)

Total [7 marks]

2. (a) attempt to form
$$(g \circ f)(x)$$
 (M1)

$$((g \circ f)(x)) = (x+2)^2 - k^2 \qquad (=x^2 + 4x + 4 - k^2)$$
 A1

[2 marks]

(b) substituting x = 4 into their $(g \circ f)(x)$ and setting their expression = 11 (M1) $(4+2)^2 - k^2 = 11$ OR $4^2 + 4(4) + 4 - k^2 = 11$ $k^2 = 25$ OR $-k^2 = -25$ (A1) $k = \pm 5$ A1 [3 marks]

Total [5 marks]

3. (a)
$$(P(A \cup B) =)0.7 + 0.75 - 0.55$$
 (A1)
= 0.9 A1

[2 marks]

(b) recognition that $A' \cap B' = (A \cup B)'$ OR $A' \cap B' = 1 - A \cup B$ (region/value may be seen in a correctly shaded/labeled Venn diagram) (M1) (=1-0.9)= 0.1 A1 Note: For the final mark, 0.1 must be stated as the candidate's answer, or

labeled as $P(A' \cap B')$ in their Venn diagram. Just seeing an unlabeled 0.1 in the correct region of their diagram earns *M1A0*.

[2 marks] Total [4 marks]

(M1)

METHOD 1 4. (a)

attem	pt to form a	at least one eo	uation, using	either S_{ϵ}	or S_c	((M1)
auom				j onanon D_{z}	0.06			,

$$65 = 25p - 5q$$
 $(13 = 5p - q)$ and $96 = 36p - 6q$ $(16 = 6p - q)$ (A1)

valid attempt to solve simultaneous linear equations in p and q by substituting or eliminating one of the variables.

p = 3, q = 2A1A1

Note: If candidate does not explicitly state their values of p and q, but gives $S_n = 3n^2 - 2n$, award final two marks as **A1A0**.

METHOD 2

attempt to form at least one equation, using either $\,S_{\!_{5}}\,$ or $\,S_{\!_{6}}\,$ (M1)

$$65 = \frac{5}{2}(2u_1 + 4d) \quad (26 = 2u_1 + 4d) \text{ and } 96 = 3(2u_1 + 5d) \quad (32 = 2u_1 + 5d)$$
(A1)

valid attempt to solve simultaneous linear equations in u_1 and d by substituting or eliminating one of the variables.

$$u_1 = 1, d = 6$$
 A1

$$S_n = \frac{n}{2} \left(2 + 6 \left(n - 1 \right) \right) = 3n^2 - 2n$$

p = 3 and q = 2

Note: If candidate does not explicitly state their values of p and q, do not award the final mark.

[5 marks]

(M1)

A1

(b)
$$u_6 = S_6 - S_5$$
 OR substituting their values of u_1 and d into $u_6 = u_1 + 5d$
OR substituting their value of u_1 into $96 = \frac{6}{2}(u_1 + u_6)$ (M1)
 $(u_6 =)96 - 65$ OR $(u_6 =)1 + 5 \times 6$ OR $96 = 3(1 + u_6)$
 $= 31$

A1

[2 marks]

Total [7 marks]

5. METHOD 1

EITHER

- 44	- 1 1 - 1 - 1 - 1	l	the short state of the second	I a al Autor a alla	/8.4	1.4
attemi	nt to lise PVt	nadoras ineorem	in a right-and	lied triandle	(1/1	11
aucing			in a nynt ang	neu mangie.	(10)	
				,	•	

$$\left(\sqrt{4^2 - 1^2}\right) = \sqrt{15} \tag{A1}$$

OR

attempt to use the Pythagorean identity $\cos^2 \alpha + \sin^2 \alpha = 1$ (M1)

$$\sin^2 \mathbf{B} \hat{\mathbf{A}} \mathbf{C} = 1 - \left(\frac{1}{4}\right)^2 \tag{A1}$$

THEN

 $\sin B\hat{A}C = \frac{\sqrt{15}}{4}$ (may be seen in area formula) A1 attempt to use 'Area = $\frac{1}{2}ab\sin C$ ' (must include their calculated value of $\sin B\hat{A}C$) (M1)

$$= \frac{1}{2} \times 16 \times \sqrt{15} \times \frac{\sqrt{15}}{4}$$
= 30 (cm²)

[6 marks]

continued

Question 5 continued

METHOD 2

attempt to find perpendicular height of triangle BAC (M1)

EITHER

height = $\sqrt{15} \times \sin B\hat{A}C$ attempt to use the Pythagorean identity $\cos^2 \alpha + \sin^2 \alpha = 1$ (M1)

height =
$$\sqrt{15} \times \sqrt{1 - \left(\frac{1}{4}\right)^2}$$
 (A1)

$$=\sqrt{15} \times \frac{\sqrt{15}}{4} \left(=\frac{15}{4}\right)$$
 (may be seen in area formula) **A1**

OR

$$adjacent = \frac{\sqrt{15}}{4}$$
 (A1)

attempt to use Pythagoras' theorem in a right-angled triangle. (M1) $\sqrt{15}(15)$

height =
$$\sqrt{15 - \frac{15}{16}} \left(= \frac{15}{4} \right)$$
 (may be seen in area formula) A1

THEN

attempt to use 'Area = $\frac{1}{2}$ base×height' (must include their calculated value for height) (M1)

$$=\frac{1}{2} \times 16 \times \frac{15}{4}$$
$$= 30 \text{ (cm}^2)$$
 A1
[6 marks]

(M1)

6. attempt to apply binomial expansion

$$(1+kx)^{n} = 1 + {}^{n}C_{1}kx + {}^{n}C_{2}k^{2}x^{2} + \dots \text{ OR } {}^{n}C_{1}k = \frac{9}{2} \text{ OR } {}^{n}C_{2} = 15$$

$$nk = \frac{9}{2}$$
(A1)

$$\frac{n(n-1)}{2} = 15 \text{ OR } \frac{n!}{(n-2)!2!} = 15$$
 (A1)

$$(n^2 - n - 30 = 0)$$
 OR $n(n-1) = 30$

valid attempt to solve

(**M1)**

A1

A1

$$(n-6)(n+5) = 0$$
 OR $6(6-1) = 30$ OR finding correct value in Pascal's triangle

$$\Rightarrow n = 6$$

$$\Rightarrow k = \frac{3}{4}$$

Note: If candidate finds n = 6 with no working shown, award **M1A0A0M1A1A0**. If candidate finds n = 6 and $k = \frac{3}{4}$ with no working shown, award **M1A0A0M1A1A1**.

[6 marks]

Section B

(a)	(i)	<i>p</i> = 9	A
	(ii)	$600 < n \le 800$	A
Note	e: Av	vard A0 if candidate answers 700.	
			[2 marks
(b)	(i)	median = 600	A
	(ii)	80% of 800 = 640	(A1)
		40 (performances less than 80% of tickets sold)	(A1)
		20 (performances)	A
			[4 marks]
(c)	(i)	any reasonable answer which suggests a biased sample (must include do not accept reasons such as "sample size is too small", or answers th say "not representative of entire audience" without a valid reason)	reason, nat simply A 2
		eg likely to come from the same part of the theatre OR be part of same OR be from priority seating OR it is convenience sampling	group
	(ii)	every 20 th person	A1A1
Note	e: Aw "ev	vard A1 for recognizing that sampling occurs at regular intervals eg very".	
	Av	vard A1 for interval length is 20.	

(iii) quota (sampling method)

A1 [4 marks] continued...

Question 7 continued

(d)	(i)	75% (of 36000 spent between \$3 and \$25)	(M1)
		= 27000	A1
	(ii)	a = 7	Δ1

METHOD 1 (e) (i) old mean is 600 (tickets) (A1) recognising new mean is old mean + 18 (M1) 600 + 18= 618 (tickets) A1 **METHOD 2** new total number of tickets = $36000 + 60 \times 18 (= 37080)$ (A1) new mean = $\frac{36000 + 60 \times 18}{60} \left(= \frac{37080}{60} \right)$ (M1) A1 =618 (tickets) no effect on the variance A1 (ii) [4 marks] Total [17 marks]

8. (a) x = 0

A1

[1 mark]

(b) (i) setting
$$\ln(2x-7) = 2\ln x - \ln d$$
 M1

 $2\ln x = \ln x^2$ (seen anywhere)

attempt to use product/quotient rule for logs (M1)

$$\ln(2x-7) = \ln\frac{x^2}{d} \text{ OR } \ln\frac{x^2}{2x-7} = \ln d \text{ OR } \ln(2x-7)d = \ln x^2$$

$$\frac{x^2}{d} = 2x - 7 \text{ OR } \frac{x^2}{2x - 7} = d \text{ OR } (2x - 7)d = x^2$$
A1

$$x^2 - 2dx + 7d = 0$$
 AG

(ii) discriminant =
$$(-2d)^2 - 4 \times 7d$$
 (A1)

 recognizing discriminant > 0
 (M1)

 $(2d)^2 - 4 \times 7d > 0$ OR $4d^2 - 28d > 0$
 A1

 $d^2 - 7d > 0$
 AG

(iii) setting d(d-7) > 0 OR d(d-7) = 0 OR sketch graph OR sign test OR $d^2 > 7d$ (M1) d < 0 or d > 7, but $d \in \mathbb{R}^+$ d > 7 (or]7,∞[) A1

[9 marks]

continued...

(M1)

(A1)

Question 8 continued

(c)
$$x^2 - 20x + 70 \ (=0)$$
 A1

attempting to solve their 3 term quadratic equation

$$\left((x-10)^2 - 30 = 0 \right) \text{ or } \left((x=) \frac{20 \pm \sqrt{(-20)^2 - 4 \times 1 \times 70}}{2} \right)$$
$$x = 10 - \sqrt{30} (=p) \text{ or } x = 10 + \sqrt{30} (=q)$$

distance =
$$2\sqrt{30}$$
 (or $\sqrt{120}$) A1

$$(a=2, b=30)$$
 (or $a=1, b=120$)

[5 marks] Total [15 marks]



(M1)

9. (a) attempt to use either the quotient or product rule

$$\frac{12(x^{2}+2)^{3}-12x\times3\times2x(x^{2}+2)^{2}}{(x^{2}+2)^{6}} \text{ OR } 12(x^{2}+2)^{-3}+12x\times(-3)\times2x(x^{2}+2)^{-4} \text{ A1A1}$$

Note: Award **A1** for correctly applying chain rule to $(x^2 + 2)^3$ and **A1** for everything else correct.

$$=\frac{12(x^{2}+2-6x^{2})}{(x^{2}+2)^{4}} \text{ OR } \frac{12(x^{2}+2)^{2}(x^{2}+2-6x^{2})}{(x^{2}+2)^{6}} \text{ OR } \frac{24-60x^{2}}{(x^{2}+2)^{4}} \text{ OR } \frac{12x^{2}+24-72x^{2}}{(x^{2}+2)^{4}} \text{ A1}$$

$$=\frac{12(2-5x^{2})}{(x^{2}+2)^{4}} \text{ AG}$$
[4 marks] continued...

Question 9 continued

(b) **EITHER**

attempts to integrate by substitution using $u = x^2 + 2$ or $u = x^2$ (M1)

$$u = x^{2} + 2 \Rightarrow \frac{du}{dx} = 2x \text{ OR } u = x^{2} \Rightarrow \frac{du}{dx} = 2x$$

Note: If candidate simply states $u = x^2 + 2$ or $u = x^2$, but does not attempt to substitute into their integral, do not award the *(M1)*.

$$\int \frac{12x}{\left(x^{2}+2\right)^{3}} dx = \int \frac{6}{u^{3}} du \quad \text{OR} \quad \int \frac{12x}{\left(x^{2}+2\right)^{3}} dx = \int \frac{6}{\left(u+2\right)^{3}} du \tag{A1}$$

$$= -3u^{-2}(+c) \text{ OR } -3(u+1)^{-2}(+c)$$
(A1)

OR

attempts to apply integration by inspection

$$6\int \frac{2x}{(x^{2}+2)^{3}} dx$$

= $6 \times \left(-\frac{1}{2}\right) (x^{2}+2)^{-2} (+c)$ (A1)(A1)

Note: Award **A1** for correct power of $(x^2 + 2)$ and **A1** for $-\frac{1}{2}$.

THEN

$$-3(x^{2}+2)^{-2}+c \quad \text{OR} \quad -\frac{3}{(x^{2}+2)^{2}}+c \quad \text{(final answer must include } +c\text{)} \qquad \qquad \textbf{A1}$$

[4 marks]

(M1)

continued...

Question 9 continued

(c) recognizing
$$g'(x) = f'(x) \Rightarrow g(x) = f(x) + k$$
 (may be seen in diagram/drawing) **A1**

area of *R* is given by subtracting functions *f* and *g* in integral(s) (M1)

$$\pm \int_{0}^{3} k dx \quad OR = \int_{0}^{3} |g - f| dx \quad OR \quad \int_{0}^{3} f(x) + k - f(x) dx \quad OR \quad \int_{0}^{3} f(x) dx - \int_{0}^{3} g(x) dx$$

$$=\pm [kx]_{0}^{3} \text{ OR } \left[-\frac{2}{\left(x^{2}+1\right)^{2}}+kx\right]_{0}^{3} - \left[-\frac{2}{\left(x^{2}+1\right)^{2}}\right]_{0}^{3} \text{ OR } \left[-\frac{2}{\left(x^{2}+1\right)^{2}}\right]_{0}^{3} - \left[-\frac{2}{\left(x^{2}+1\right)^{2}}+kx\right]_{0}^{3} (A1)$$
$$\pm 3k = \frac{21}{2}$$
(A1)

$$k = \pm \frac{21}{6} \left(= \pm \frac{7}{2} = \pm 3.5 \right)$$

$$g(x) = \frac{12x}{\left(x^2 + 2\right)^3} - \frac{7}{2} \text{ AND } g(x) = \frac{12x}{\left(x^2 + 2\right)^3} + \frac{7}{2} \text{ (accept } f(x) + \frac{7}{2} \text{ AND } f(x) - \frac{7}{2} \text{)}$$

[5 marks] Total [13 marks]



Markscheme

May 2023

Mathematics: analysis and approaches

Standard level

Paper 1

22 pages



© International Baccalaureate Organization 2023

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2023

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2023

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *M0* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies A3, M2 etc., do not split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this
 working is incorrect and/or suggests a misunderstanding of the question. This will encourage a
 uniform approach to marking, with less examiner discretion. Although some candidates may be
 advantaged for that specific question item, it is likely that these candidates will lose marks
 elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used **in a subsequent part**. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	8√2	5.65685 (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)
2.	$\frac{35}{72}$	0.468111 (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)

3 Implied marks

Implied marks appear in **brackets e.g.** (*M1*), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is *(M1)A1*, it is possible to award full marks for *their* correct answer, **without working being seen.** For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (*MR*). A candidate should be penalized only once for a particular misread. Use the *MR* stamp to indicate that this has been a misread and do not award the first mark, even if this is an *M* mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** ... OR.

7 Alternative forms

Unless the question specifies otherwise, accept equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, **M** marks and intermediate **A** marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "*from the use of 3 sf values*".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an *A* mark to be awarded, arithmetic should be completed, and

any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$. An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or 5 10

written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate A marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".

$\operatorname{arc}+2 \times \operatorname{radius} OR 10 + 4 + 4$	
=18 (cm)	A1
	[2 marks]
$10 = 4\theta$	(A1)
$\theta = \frac{10}{4} \left(= \frac{5}{2}, 2.5 \right)$	A1
	[2 marks]
area $=\frac{1}{2}\left(\frac{10}{4}\right)(4^2)$ (=1.25×16)	(A1)

A1

(M1)

[2 marks] Total [6 marks]

Section A

1.

(a)

(b)

(c)

 $=20 (cm^2)$

attempts to find perimeter

2. (a) $u_1 + 3d = u_4$ (M1) 0.6 + 3d = 0.15d = -0.15 A1

(M1)

(b) METHOD 1

$u_2 = 0.45$ or $u_3 = 0.3$ (may be seen in their equation)	(A1)
---	------

summing their probabilities to 1 (seen anywhere)

$$\frac{0.6}{k} + \frac{u_2}{k} + \frac{u_3}{k} + \frac{0.15}{k} = 1$$

$$\frac{0.6}{k} + \frac{0.45}{k} + \frac{0.3}{k} + \frac{0.15}{k} = 1 \text{ (or equivalent)}$$

$$\frac{1.5}{k} = 1$$

$$k = 1.5$$
A1

METHOD 2 (using S_n formula)

$$S_{4} = \frac{4}{2} \Big(2 \Big(0.6 \Big) + \Big(4 - 1 \Big) \Big(-0.15 \Big) \Big) \quad \text{OR} \quad S_{4} = \frac{4}{2} \Big(2 \Big(\frac{0.6}{k} \Big) + \Big(4 - 1 \Big) \Big(\frac{-0.15}{k} \Big) \Big)$$

OR
$$S_{4} = \frac{4}{2} \Big(0.6 + 0.15 \Big) \quad \text{OR} \quad S_{4} = \frac{4}{2} \Big(\frac{0.6}{k} + \frac{0.15}{k} \Big) \quad \text{(or equivalent)}$$
(A1)

summing their probabilities to 1 (seen anywhere)

$$\frac{u_1}{k} + \frac{u_2}{k} + \frac{u_3}{k} + \frac{u_4}{k} = 1 \quad \text{OR} \quad u_1 + u_2 + u_3 + u_4 = k \quad \text{OR} \quad S_4 = 1 \quad \text{OR} \quad S_4 = k$$

$$\frac{4}{2}(2(0.6)+(4-1)(-0.15)) = k \text{ (or equivalent)}$$
(A1)

$$k = 1.5$$

A1

(M1)

[4 marks] Total [6 marks]

3. (a) (i)
$$x = 2$$
 A1

 (ii) $y = 1$
 A1

-9-

[2 marks]

(b) (i)
$$\left(0, \frac{3}{2}\right)$$

(ii) (3,0) A1

[2 marks]





Question 3 continued



two correct branches with correct asymptotic behaviour and intercepts clearly shown A1

[1 mark] Total [5 marks] 4. substitutes into $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ to form

$$0.55 = 0.4 + P(B) - P(A \cap B)$$
 (or equivalent) (A1)

substitutes into
$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$
 to form $0.25 = \frac{P(A \cap B)}{P(B)}$ (or equivalent) (A1)

attempts to combine their two probability equations to form an equation in P(B) (M1) Note: The above two **A** marks are awarded independently.

correct equation in P(B)

$$0.55 = 0.4 + P(B) - 0.25P(B)$$
 OR $\frac{P(B) - 0.15}{P(B)} = 0.25$ OR $P(B) - 0.15 = 0.25P(B)$

(or equivalent)

$$P(B) = \frac{15}{75} \left(= \frac{1}{5} = 0.2 \right)$$

A1

A1

Total [5 marks]

$$\mathbf{5.} \qquad A = \int_0^c \frac{x}{x^2 + 2} \mathrm{d}x$$

EITHER

attempts to integrate by inspection or substitution using $u = x^2 + 2$ or $u = x^2$

– 12 –

Note: If candidate simply states $u = x^2 + 2$ or $u = x^2$, but does not attempt to integrate, do not award the **(M1)**.

Note: If candidate does not explicitly state the *u*-substitution, award the *(M1)* only for expressions of the form $k \ln u$ or $k \ln (u+2)$.

$$\left[\frac{1}{2}\ln u\right]_{2}^{c^{2}+2} \text{ OR } \left[\frac{1}{2}\ln(u+2)\right]_{0}^{c^{2}} \text{ OR } \left[\frac{1}{2}\ln(x^{2}+2)\right]_{0}^{c}$$

Note: Limits may be seen in the substitution step.

OR

attempts to integrate by inspection

Note: Award the *(M1)* only for expressions of the form $k \ln(x^2 + 2)$.

$$\left\lceil \frac{1}{2} \ln \left(x^2 + 2 \right) \right\rceil_0^c$$

Note: Limits may be seen in the substitution step.

THEN

correctly substitutes their limits into their integrated expression

$$\frac{1}{2} \left(\ln \left(c^2 + 2 \right) - \ln 2 \right) \left(= \ln 3 \right) \text{ OR } \frac{1}{2} \ln \left(c^2 + 2 \right) - \frac{1}{2} \ln 2 \left(= \ln 3 \right)$$

continued...

(M1)

(M1)

A1

(M1)

Question 5 continued

correctly applies at least one log law to their expression (M1)

$$\frac{1}{2}\ln\left(\frac{c^2+2}{2}\right)(=\ln 3) \text{ OR } \ln\sqrt{c^2+2} - \ln\sqrt{2}(=\ln 3) \text{ OR } \ln\left(\frac{c^2+2}{2}\right) = \ln 9$$
OR $\ln(c^2+2) - \ln 2 = \ln 9$ OR $\ln\sqrt{\frac{c^2+2}{2}}(=\ln 3)$ OR $\ln\frac{\sqrt{c^2+2}}{\sqrt{2}}(=\ln 3)$
Note: Condone the absence of $\ln 3$ up to this stage.

$$\frac{c^2+2}{2} = 9 \text{ OR } \sqrt{\frac{c^2+2}{2}} = 3$$
A1
 $c^2 = 16$
 $c = 4$
A1
Note: Award A0 for $c = \pm 4$ as a final answer.
Total [6 marks]

– 13 –

6. attempts to form $(g \circ f)(x)$ (M1)

$$[f(x)]^{2} + f(x) + 3 \quad \text{OR} \quad (ax+b)^{2} + ax+b+3$$

$$a^{2}x^{2} + 2abx + b^{2} + ax+b+3(=4x^{2} - 14x + 15)$$
(A1)

equates their corresponding terms to form at least one equation

$$a^{2}x^{2} = 4x^{2}$$
 OR $a^{2} = 4$ OR $2abx + ax = -14x$ OR $2ab + a = -14$ OR $b^{2} + b + 3 = 15$

$$a = \pm 2$$
 (seen anywhere) A1

attempt to use 2ab + a = -14 to pair the correct values (seen anywhere) (M1)

$$f(x) = 2x - 4$$
 (accept $a = 2$ with $b = -4$), $f(x) = -2x + 3$ (accept $a = -2$ with $b = 3$) A1A1

[7 marks]

(M1)

Section B

(a)	x = -2 (must be an equation)	A1
		[1 mark]
(b)	h = -2, $k = -5$	A1A1
		[2 marks]
(c)	substituting $x = 0$ into $f(x)$	(M1)
	$y = \frac{1}{4}(0+2)^2 - 5$	
	y = -4 (accept P(0, -4))	A1
		[2 marks]
(d)	$f'(x) = \frac{1}{2}(x+2)\left(=\frac{1}{2}x+1\right)$	(A1)
	substituting $x = 0$ into their derivative	(M1)
	f'(0) = 1	
	gradient of normal is -1 (may be seen in their equation)	A1
N. 4	y = -x - 4 (accept $a = -1$, $b = -4$)	A1
Not	e: Award A0 for $L = -x - 4$ (without the $y =$).	
		[4 marks]
		continued

Question 7 continued

(e) equating their f(x) to their L (M1)

$$\frac{1}{4}(x+2)^2 - 5 = -x - 4$$

$$\frac{1}{4}x^2 + 2x = 0 \quad \text{(or equivalent)}$$
(A1)

valid attempt to solve their quadratic

$$\frac{1}{4}x(x+8) = 0 \quad \text{OR} \quad x(x+8) = 0$$

$$x = -8$$
A1

Note: Accept both solutions x = -8 and x = 0 here, x = -8 may be seen in working to find coordinates of Q or distance.

substituting their value of	f x (not $x = 0$) into the	r $f(x)$ or their L	(M1)
-----------------------------	-----------------------------	---------------------	------

$$y = -(-8) - 4$$
 or $y = \frac{1}{4}(-8+2)^2 - 5$
Q(-8, 4) A1

correct substitution into distance formula

$$\sqrt{(-8-0)^2+(4-(-4))^2}$$

distance = $\sqrt{128}$ (= $8\sqrt{2}$)

A1

(A1)

(M1)

[8 marks] Total [17 marks] 8.

(a) (i) attempt to use Pythagoras (M1)

$$\sin\theta = \frac{\sqrt{5}}{3}$$

(ii) attempt to substitute into double-angle identity using their value of $\sin\theta$ (M1)

$$\sin 2\theta = 2\left(\frac{\sqrt{5}}{3}\right)\left(\frac{2}{3}\right)$$

$$\sin 2\theta = \frac{4\sqrt{5}}{9}$$
[4 marks]

(b) METHOD 1 (using values from part (a))

$$\frac{b}{\sin\theta} = \frac{a}{\sin 2\theta}$$

attempt to use sine rule with their values from part (a) (M1)

$$\frac{b}{\left(\frac{\sqrt{5}}{3}\right)} = \frac{a}{\left(\frac{4\sqrt{5}}{9}\right)} \quad \text{OR} \quad \frac{\left(\frac{\sqrt{5}}{3}\right)}{b} = \frac{\left(\frac{4\sqrt{5}}{9}\right)}{a}$$

correct working that leads to AG

$$\frac{\sqrt{5}}{3}a = \frac{4\sqrt{5}}{9}b \quad \text{OR} \quad \frac{3b}{\sqrt{5}} = \frac{9a}{4\sqrt{5}} \quad \text{OR} \quad \frac{a}{3} = \frac{4b}{9} \text{ (or equivalent)}$$
$$b = \frac{3a}{4} \qquad \qquad \textbf{AG}$$

continued...

A1

Question 8 continued

METHOD 2 (double-angle identity)

 $\frac{b}{\sin\theta} = \frac{a}{\sin 2\theta}$

using double-angle identity

 $\frac{b}{\sin\theta} = \frac{a}{2\sin\theta\cos\theta} \quad \text{OR} \quad b = \frac{a\sin\theta}{2\sin\theta\cos\theta} \quad \text{OR} \quad b = \frac{a}{2\cos\theta}$

correct working (involving substituting $\cos \theta = \frac{2}{3}$) that leads to **AG A1**

$$b = \frac{a \sin \theta}{2 \sin \theta \left(\frac{2}{3}\right)} \quad \text{OR} \quad b = \frac{a \left(\frac{\sqrt{5}}{3}\right)}{2 \left(\frac{\sqrt{5}}{3}\right) \left(\frac{2}{3}\right)} \quad \text{OR} \quad b = \frac{a}{2 \left(\frac{2}{3}\right)} \quad \text{(or equivalent)}$$
$$b = \frac{3a}{4} \qquad \qquad \text{AG}$$

(c)	METHOD 1 (using supplementary angles)	
	recognizing CÂD and BÂC are supplementary	(M1)
	recognizing supplementary angles have the same sine value	(A1)
	$\sin \hat{CAD} = \sin 2\theta$	
	$\sin \hat{CAD} = \frac{4\sqrt{5}}{9}$	A1

continued...

(A1)

Question 8 continued

METHOD 2 (using sine rule)(M1) $a = \frac{b}{\sin C A D} = \frac{b}{\sin \theta}$

correct substitution of $\sin \theta = \frac{\sqrt{5}}{3}$ and $b = \frac{3a}{4}$ into sine rule (A1)

$$\frac{a}{\sin \hat{CAD}} = \frac{\left(\frac{3a}{4}\right)}{\left(\frac{\sqrt{5}}{3}\right)} \quad \text{OR} \quad \sin \hat{CAD} = \frac{a\left(\frac{\sqrt{5}}{3}\right)}{\left(\frac{3a}{4}\right)} \quad \text{(or equivalent)}$$

 $\sin \hat{CAD} = \frac{4\sqrt{5}}{9}$

[3 marks]

(M1)

(A1)

(d) METHOD 1 (using \hat{CAD} in area formula)

recognizing $\hat{DCA} = \theta$ (A1)

recognizing
$$AD = b\left(=\frac{3a}{4}\right)$$
 (A1)

correct substitution into area formula (must substitute expressions for two sides and name/expression/value for $\sin C\hat{A}D$)

area
$$=\frac{1}{2}(b)(b)\left(\frac{4\sqrt{5}}{9}\right)$$
 OR area $=\frac{1}{2}(b)(b)\sin 2\theta$ OR area $=\frac{1}{2}(b)(b)\sin C\hat{A}D$

correct substitution in terms of a

$$\operatorname{area} = \frac{1}{2} \left(\frac{3a}{4} \right) \left(\frac{3a}{4} \right) \left(\frac{4\sqrt{5}}{9} \right)$$
$$\operatorname{area} = \frac{\sqrt{5} a^2}{8}$$

continued...

– 19 –
Question 8 continued

METHOD 2 (using \hat{ACD} or \hat{ADC} in area formula)

recognizing
$$CD = a$$
 (A1)

recognizing
$$AD = b\left(=\frac{3a}{4}\right)$$
 and/or $D\hat{C}A = \theta$ (A1)

correct substitution into area formula (must substitute expressions for two sides and name/expression/value for $\sin A\hat{D}C$ or $\sin A\hat{C}D$)

area
$$=\frac{1}{2}(a)(b)\left(\frac{\sqrt{5}}{3}\right)$$
 OR area $=\frac{1}{2}(a)(b)\sin\theta$ OR area $=\frac{1}{2}(a)(b)\sin A\hat{D}C$
OR area $=\frac{1}{2}(a)(b)\sin A\hat{C}D$

correct substitution in terms of a

area
$$=\frac{1}{2}(a)\left(\frac{3a}{4}\right)\left(\frac{\sqrt{5}}{3}\right)$$

area $=\frac{\sqrt{5}a^2}{8}$

A1

(A1)

(M1)

[5 marks] Total [14 marks]

9. (a)
$$y^2 = 9 - x^2$$
 OR $y = \pm \sqrt{9 - x^2}$ A1
(since $y > 0$) $\Rightarrow y = \sqrt{9 - x^2}$ AG

[1 mark]

(b)
$$b = 2y \left(= 2\sqrt{9 - x^2}\right)$$
 or $h = x + 3$ (A1)

attempts to substitute their base expression and height expression into $A = \frac{1}{2}bh$ (M1)

$$A = \sqrt{9 - x^2} (x + 3) \text{ (or equivalent)} \left(= \frac{2(x + 3)\sqrt{9 - x^2}}{2} = x\sqrt{9 - x^2} + 3\sqrt{9 - x^2} \right)$$
 A1

[3 marks]

(c) attempts to use the product rule to find
$$\frac{dA}{dx}$$
 (M1)

attempts to use the chain rule to find
$$\frac{d}{dx}\sqrt{9-x^2}$$
 (M1)

$$\left(\frac{\mathrm{d}A}{\mathrm{d}x}\right)\sqrt{9-x^2} + (3+x)\left(\frac{1}{2}\right)(9-x^2)^{-\frac{1}{2}}(-2x)\left(=\sqrt{9-x^2}-\frac{x^2+3x}{\sqrt{9-x^2}}\right)$$
A1

$$\left(\frac{\mathrm{d}A}{\mathrm{d}x}\right) = \frac{9-x^2}{\sqrt{9-x^2}} - \frac{x^2+3x}{\sqrt{9-x^2}} \left(=\frac{9-x^2-(x^2+3x)}{\sqrt{9-x^2}}\right)$$

$$\frac{\mathrm{d}A}{\mathrm{d}x} = \frac{9 - 3x - 2x^2}{\sqrt{9 - x^2}}$$

[4 marks]

continued...

Question 9 continued

(d)
$$\frac{dA}{dx} = 0 \left(\frac{9 - 3x - 2x^2}{\sqrt{9 - x^2}} = 0 \right)$$
 (M1)

attempts to solve $9-3x-2x^2 = 0$ (or equivalent)

$$-(2x-3)(x+3)(=0)$$
 OR $x = \frac{3 \pm \sqrt{(-3)^2 - 4(-2)(9)}}{2(-2)}$ (or equivalent) (A1)

$$x = \frac{3}{2}$$

Note: Award the above **A1** if x = -3 is also given.

substitutes their value of x into either
$$y = \sqrt{9 - x^2}$$
 or $y = -\sqrt{9 - x^2}$ (M1)

Note: Do not award the above **(M1)** if $x \le 0$.

$$y = -\sqrt{9 - \left(\frac{3}{2}\right)^2}$$
$$= -\frac{\sqrt{27}}{2} \left(= -\frac{3\sqrt{3}}{2}, = -\sqrt{\frac{27}{4}}, = -\sqrt{6.75} \right)$$

A1

(M1)

A1

[6 marks] Total [14 marks]



Markscheme

May 2023

Mathematics: analysis and approaches

Standard level

Paper 1

18 pages



© International Baccalaureate Organization 2023

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2023

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2023

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *M0* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies A3, M2 etc., do not split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this
 working is incorrect and/or suggests a misunderstanding of the question. This will encourage a
 uniform approach to marking, with less examiner discretion. Although some candidates may be
 advantaged for that specific question item, it is likely that these candidates will lose marks elsewhere
 too.
- An exception to the previous rule is when an incorrect answer from further working is used **in a subsequent part**. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.

Examples:

	Correct	Further	Any FT issues?	Action	
	answer seen	working seen		Action	
1.		5.65685	No.	Award A1 for the final mark	
	$8\sqrt{2}$	(incorrect	Last part in question.	(condone the incorrect further	
		decimal value)		working)	
2.	35	0.468111	Yes.	Award A0 for the final mark	
	<u> </u>	(incorrect	Value is used in	(and full FT is available in	
	72	decimal value)	subsequent parts.	subsequent parts)	

3 Implied marks

Implied marks appear in **brackets e.g.** (*M1*), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is *(M1)A1*, it is possible to award full marks for *their* correct answer, **without working being seen**. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (*MR*). A candidate should be penalized only once for a particular misread. Use the *MR* stamp to indicate that this has been a misread and do not award the first mark, even if this is an *M* mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- **MR** can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** ... OR.

7 Alternative forms

Unless the question specifies otherwise, accept equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, **M** marks and intermediate **A** marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.

• In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "*from the use of 3 sf values*".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an *A* mark to be awarded, arithmetic should be completed, and any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$.

An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or

written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate A marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".

Section A

1. (a) M (6,-3) A1A1
[2 marks]
(b) gradient of [PQ] =
$$-\frac{5}{9}$$
 (A1)
gradient of $L = \frac{9}{5}$ A1
[2 marks]
(c) $y+3=\frac{9}{5}(x-6)$ OR $y=\frac{9}{5}x-\frac{69}{5}$ (or equivalent) A1

Note: Do not accept
$$L = \frac{9}{5}x - \frac{69}{5}$$
.

[1 mark] Total [5 marks]

2.	(a)	recognizing $f(x) = 0$	(M1)
		x = -1	A1
			[2 marks]

(b)	(i)	x = 2 (must be an equation with x)	A1
	(ii)	$y = \frac{7}{2}$ (must be an equation with y)	A1

[2 marks]

(c) **EITHER**

interchanging x and y	(M1)
2xy - 4x = 7y + 7	
correct working with y terms on the same side: $2xy - 7y = 4x + 7$	(A1)

OR

2yx - 4y = 7x + 7	
correct working with x terms on the same side: $2yx - 7x = 4y + 7$	(A1)

interchanging x and y OR making x the subject $x = \frac{4y+7}{2y-7}$ (M1)

THEN

$$f^{-1}(x) = \frac{4x+7}{2x-7}$$
 (or equivalent) $\left(x \neq \frac{7}{2}\right)$ A1

[3 marks] Total [7 marks]

probability =
$$\frac{34}{40}$$
 A1

(ii) attempt to find expected value (M1) $\frac{16}{40} + \left(2 \times \frac{13}{40}\right) + \left(3 \times \frac{2}{40}\right) + \left(4 \times \frac{3}{40}\right)$ $\frac{60}{40} (=1.5)$ A1

[4 marks]

(M1)

(b) evidence of **their** rides/visitor $\times 1000$ or $\div 10$ 1500 OR 0.15 150 (times)

A1 [2 marks] Total [6 marks] 4. (a) $1-2\sin^2 x = \sin x$ $2\sin^2 x + \sin x - 1 = 0$ AG

[1 mark]

(M1)

(b) valid attempt to solve quadratic (M1)

$$(2\sin x - 1)(\sin x + 1) \text{ OR } \frac{-1 \pm \sqrt{1 - 4(2)(-1)}}{2}$$

$$2(2)$$
 recognition to solve for sin *x*

 $\sin x = \frac{1}{2} \text{ OR } \sin x = -1$ any correct solution from $\sin x = -1$ A1
any correct solution from $\sin x = \frac{1}{2}$ Note: The previous two marks may be awarded for degree or radian values,
irrespective of domain. $x = -\frac{\pi}{2}, \frac{\pi}{6}, \frac{5\pi}{6}$ A1
Note: If no working shown, award no marks for a final value(s).
Award A0 for $-\frac{\pi}{2}, \frac{\pi}{6}, \frac{5\pi}{6}$ if additional values also given.

[5 marks] Total [6 marks]

5.	recognition of quadratic in e^x	(M1)
	$(e^{x})^{2} - 3e^{x} + \ln k (= 0) \text{ OR } A^{2} - 3A + \ln k (= 0)$	
	recognizing discriminant ≥ 0 (seen anywhere)	(M1)
	$(-3)^2 - 4(1)(\ln k)$ OR $9 - 4\ln k$	(A1)
	$\ln k \le \frac{9}{4}$	(A1)
	e ^{9/4} (seen anywhere)	A1

$$0 < k \le \mathrm{e}^{9/4}$$

[6 marks]

A1



6. (a) recognition that period is 4m OR substitution of a point on f (except the origin)

$$4m = \frac{2\pi}{q} \text{ OR } 1 = \sin qm$$

$$m = \frac{\pi}{2q}$$
A1

(M1)

(b) horizontal scale factor is
$$\frac{3}{2}$$
 (seen anywhere)

(A1)

Note: This **(A1)** may be earned by seeing a period of 6m, half period of 3m or the correct *x*-coordinate of the maximum/minimum point.



Note: Curve must be an approximate sinusoidal shape (sine or cosine). Only in this case, award the following:
A1 for correct amplitude.
A1 for correct domain.
A1 for correct max and min points and correct *x*-intercepts.

> [4 marks] Total [6 marks]

A1A1A1

Section B

– 13 –

7.	(a)	substitution of $x = 0$	(M1)
		(y =) 3 (accept $(0, 3)$)	A1
			[2 marks]
	(b)	evidence of using the product rule	(M1)
		$h'(x) = 2e^x + 2xe^x$	A1
			[2 marks]
	(c)	setting their derivative equal to zero	(M1)
		correct working	(A1)
		$2e^{x}(1+x) (=0) \text{ OR } -2x = 2$	
		x = -1 (seen anywhere, and must follow on from their derivative)	A1
		substituting their value of x into $h(x)$	(M1)
		$y = -\frac{2}{e} + 3 (= -2e^{-1} + 3)$	A1
		$A\left(-1,-\frac{2}{e}+3\right)$	
			[5 marks]
			continued

Question 7 continued

(d) (i)
$$h''(x) = 2e^x + 2e^x + 2xe^x \text{ OR } 2e^x + 2e^x(1+x)$$
 A1A1

– 14 –

Note: Award **A1** for
$$(2e^x)' = 2e^x$$
, **A1** for $2e^x + 2xe^x$ or $(2x+2)e^x$

$$h''(x) = (2x+4)e^x \qquad \qquad \mathbf{AG}$$

(ii) recognition that h'' > 0 OR attempt to find point of inflexion (M1) since $e^x > 0$, 2x + 4 > 0 OR 2x + 4 = 0 ($\Rightarrow x = -2$)

[4 marks] Total [13 marks]



8.	(a)	(i)	recognition that $n = 5$	(M1)
			$S_5 = 45$	A1

(ii) METHOD 1

recognition that $S_5 + u_6 = S_6$ (M1)

METHOD 2

recognition that $60 = \frac{6}{2}(S_1 + u_6)$	(M1)

$$b0 = 3(5 + u_6)$$

 $u_6 = 15$ A1

METHOD 3

substituting their u_1 and d values into $u_1 + (n-1)d$	(M1)
$u_6 = 15$	A1

[4 marks]

(b) recognition that $u_1 = S_1$ (may be seen in (a)) OR substituting their u_6 into S_6 (M1) OR equations for S_5 and S_6 in terms of u_1 and d

1+4 OR
$$60 = \frac{6}{2}(u_1 + 15)$$

 $u_1 = 5$

A1

[2 marks] continued...

– 15 –

Question 8 continued

(c) **EITHER**

valid attempt to find d (may be seen in (a) or (b))	(M1)
d = 2	(A1)

OR

valid attempt to find $S_n - S_{n-1}$ (M1)

$$n^{2} + 4n - (n^{2} - 2n + 1 + 4n - 4)$$
 (A1)

OR

equating $n^2 + 4n = \frac{n}{2}(5 + u_n)$ (M1)

 $2n+8=5+u_n$ (or equivalent) (A1)

THEN

$$u_n = 5 + 2(n-1)$$
 OR $u_n = 2n+3$

[3 marks]

A1

(d) recognition that
$$v_2 r^2 = v_4 OR (v_3)^2 = v_2 \times v_4$$
 (M1)
 $r^2 = 3 OR v_3 = (\pm)5\sqrt{3}$ (A1)
 $r = \pm\sqrt{3}$

Note: If no working shown, award *M1A1A0* for $\sqrt{3}$.

[3 marks]

(e) recognition that *r* is negative (M1) $v_5 = -15\sqrt{3} \quad \left(=-\frac{45}{\sqrt{3}}\right)$ A1

> [2 marks] Total [14 marks]

(a)	attempt to integrate v (integration of at least one term)	
	$(s(t) =) -\frac{1}{4}t^4 + \frac{7}{6}t^3 - t^2 + 6t \ (+C)$	A2
Note	e: Award A1 for at least two correct terms.	
	substitution of $t = 1$ into their integrated expression	(M1)
	displacement = $5\frac{11}{12}\left(=\frac{71}{12}\right)$ (m)	A1
		[5 marks]
(b)	attempt to differentiate v (differentiation of at least one term)	(M1)
	$a(t) = -3t^2 + 7t - 2$	A1
		[2 marks]
(c)	setting their $v'(t) = 0$	(M1)
	$-3t^2 + 7t - 2 = 0$	
	valid attempt to solve quadratic	(M1)
	$(3t-1)(t-2) = 0$ OR $\frac{-7 \pm \sqrt{49 - 4(-3)(-2)}}{-6}$	
	$t = \frac{1}{3}, 2$ ($t = \frac{1}{3}$ may be omitted)	(A1)
	substitute their largest positive <i>t</i> -value into $v(t)$	(M1)
	greatest speed is 8 (ms ⁻¹)	A1

– 17 –

9.

[5 marks]

continued...

Question 9 continued

(d) attempt to check other boundary value at t = 4(M1) $v(4) = -64 + 56 - 8 + 6 \quad (= -10)$ greatest speed is 10 ms⁻¹A1

(e) identifying correct intervals where speed increases (may be seen in integral) (A1)(A1) $t = \frac{1}{3}$ to t = 2 and t = k to t = 4 $\int_{\frac{1}{3}}^{2} v(t) dt + \int_{k}^{4} |v(t)| dt$ OR $\int_{\frac{1}{3}}^{2} v dt + \left| \int_{k}^{4} v dt \right|$ OR $\int_{\frac{1}{3}}^{2} v(t) dt - \int_{k}^{4} v(t) dt$ A1

Note: Condone missing dt.

[3 marks] Total [17 marks]



Markscheme

November 2022

Mathematics: analysis and approaches

Standard level

Paper 1

24 pages



© International Baccalaureate Organization 2022

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2022

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2022

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *M0* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies A3, M2 etc., do not split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this
 working is incorrect and/or suggests a misunderstanding of the question. This will encourage a
 uniform approach to marking, with less examiner discretion. Although some candidates may be
 advantaged for that specific question item, it is likely that these candidates will lose marks
 elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used in a **subsequent part**. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	8√2	5.65685 (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)
2.	$\frac{35}{72}$	0.468111… (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)

3 Implied marks

Implied marks appear in **brackets e.g.** (*M1*), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is *(M1)A1*, it is possible to award full marks for *their* correct answer, **without working being seen.** For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (*MR*). A candidate should be penalized only once for a particular misread. Use the *MR* stamp to indicate that this has been a misread and do not award the first mark, even if this is an *M* mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** ... OR.

7 Alternative forms

Unless the question specifies otherwise, accept equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, **M** marks and intermediate **A** marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come *"from the use of 3 sf values"*.

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an **A** mark to be awarded, arithmetic should be completed, and any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$. An exception to this is simplifying fractions, where lowest form is not required (although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Completing any operations such as addition and

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate A marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".

Section A

1.	(a)	gradient of g is -2 (may be seen in function, do not accept $-2x+3$)	(A1)
		g(x) = -2x	A1
			[2 marks]

(b) gradient is
$$\frac{1}{2}$$
 (may be seen in function) (A1)

attempt to substitute **their** gradient and (-1,2) into any form of equation for straight line *(M1)*

$$y-2 = \frac{1}{2}(x+1)$$
 OR $2 = \frac{1}{2} \cdot (-1) + c$

$$h(x) = \frac{1}{2}(x+1) + 2\left(=\frac{1}{2}x + \frac{5}{2}\right)$$
A1

[3 marks]

(c)
$$(g \circ h)(x) = -2\left(\frac{1}{2}x + \frac{5}{2}\right) \text{ OR } h(0) = \frac{5}{2} \text{ OR } g\left(\frac{5}{2}\right)$$
 (A1)

$$(g \circ h)(0) = -5 \tag{A1}$$

[2 marks]

Total [7 marks]

2.
$$g'(x) = 2x e^{x^2 + 1}$$
 (A2)

substitute x = -1 into **their** derivative

$$g'(-1) = -2e^2$$

Note: Award **A0M0A0** in cases where candidate's incorrect derivative is $g'(x) = e^{x^2+1}$.

[4 marks]

(M1)

A1



3. (a) (i) attempt to find midpoint of A and B (M1) centre (-1,3,-2) (accept vector notation and/or missing brackets) A1 (ii) attempt to find AB or half of AB or distance between the centre and A (or B) (M1) $\sqrt{4^2+2^2+4^2}$ $\sqrt{2^2+1^2+2^2}$

$$\frac{1}{2} \text{ or } \sqrt{2^2 + 1^2 + 2^2}$$
= 3
[4 marks]

(b) attempt to find the distance between their centre and V
(the perpendicular height of the cone) (M1)

$$\sqrt{0^2 + 4^2 + 2^2}$$
 OR $\sqrt{(\text{their slant height})^2 - (\text{their radius})^2}$
 $= \sqrt{20} (= 2\sqrt{5})$ (A1)
Volume $= \frac{1}{3}\pi 3^2 \sqrt{20}$
 $= 3\pi \sqrt{20} (= 6\pi \sqrt{5})$ A1
[3 marks]
Total [7 marks]

4. (a)

Note: Award a maximum of M1A0A0 if the candidate manipulates both sides of the equation (such as moving terms from one side to the other).

METHOD 1 (working with LHS)

attempting to expand
$$(a^2-1)^2$$
 (do not accept a^4+1 or a^4-1) (M1)

LHS =
$$a^2 + \frac{a^4 - 2a^2 + 1}{4}$$
 or $\frac{4a^2 + a^4 - 2a^2 + 1}{4}$ A1

$$=\frac{a^{4}+2a^{2}+1}{4}$$

$$=\left(\frac{a^{2}+1}{2}\right)^{2} (= \text{RHS})$$
A1

METHOD 2 (working with RHS)

attempting to expand $(a^2 + 1)^2$

$$RHS = \frac{a^4 + 2a^2 + a^2}{4}$$

$$=\frac{4a^2+a^4-2a^2+1}{4}$$

$$=a^{2}+\frac{a^{4}-2a^{2}+1}{4}$$
 A1

$$=a^{2}+\left(\frac{a^{2}-1}{2}\right)^{2}$$
 (=LHS) **AG**

Note: Do not award the final A1 if further working contradicts the AG.

[3 marks]

AG

(M1)

A1

continued...

Question 4 continued

(b) recognise base and height as a and $\left(\frac{a^2-1}{2}\right)$ (may be seen in diagram) (M1)

correct substitution into triangle area formula

Area
$$=\frac{a}{2}\left(\frac{a^2-1}{2}\right)$$
 (or equivalent) $\left(=\frac{a(a^2-1)}{4}=\frac{a^3-a}{4}\right)$

[2 marks]

A1

Total [5 marks]



– 11 –

5. recognizing need to integrate

$$\int \frac{6x}{x^{2}+1} dx \quad \text{OR} \quad u = x^{2}+1 \quad \text{OR} \quad \frac{du}{dx} = 2x$$

$$\int \frac{3}{u} du \quad \text{OR} \quad 3\int \frac{2x}{x^{2}+1} dx \tag{A1}$$

$$= 3\ln(x^{2}+1)(+c) \text{ or } 3\ln u(+c)$$
 A1

correct substitution of x = 1 and f(x) = 5 or x = 1 and u = 2 into equation

using **their** integrated expression (must involve *c*)

$$5 = 3 \ln 2 + c$$

$$f(x) = 3\ln(x^{2}+1) + 5 - 3\ln 2 \left(= 3\ln(x^{2}+1) + 5 - \ln 8 = 3\ln\left(\frac{x^{2}+1}{2}\right) + 5\right)$$

(or equivalent)

Note: Accept the use of the modulus sign in working and the final answer.

[5 marks]

(M1)

A1

(**M1**)

6. (a)
$$P(A \cap B) = 0.24$$
 A1

[1 mark]

(b)
$$P(A \cup B) = 1.1 - P(A \cap B)$$
 (A1)

$$(0 \le) \mathbf{P}(A \cup B) \le 1 \tag{M1}$$

Note: This may be conveyed in a clearly labelled diagram or written explanation where $P(A \cup B) = 1$

the minimum value of $P(A \cap B)$ is 0.1

(c) A is a subset of B (so $P(A \cap B) = P(A)$).

Note: This may be conveyed in a clearly labelled diagram where *A* is completely inside *B*, or in a written explanation indicating that $P(A \cap B) = P(A)$

so the maximum value of $P(A \cap B)$ is 0.3

Note: Do not award R0A1.

[2 marks]

Total [6 marks]

A1

R1

A1

[3 marks]

Section B

7. (a) correct substitution of
$$h = 3$$
 and $k = 2$ into $f(x)$

$$f(x) = a(x-3)^2 + 2$$

correct substitution of (5,0)

$$0 = a(5-3)^2 + 2 \left(a = -\frac{1}{2}\right)$$

Note: The first two A marks are independent.

$$f(x) = -\frac{1}{2}(x-3)^2 + 2$$

A1

(A1)

(A1)

(A1)

[3 marks]

(b)	(i)	METHOD 1		
		correct substitution of $(1, 4)$		
		p + (t-1) - p = 4		

substituting their value of t into 9p-3(t-1)-p=4 (M1)

$$8p-12=4$$

 $p=2$ **A1**

METHOD 2

correct substitution of ONE of the coordinates	(-3,4) or $(1,4)$	(A1)
--	-------------------	------

$$9p-3(t-1)-p=4$$
 OR $p+(t-1)-p=4$

valid attempt to solve their two equations

$$p = 2, t = 5$$
 A1A1

$$\left(g\left(x\right)=2x^{2}+4x-2\right)$$

(ii) attempt to find the *x*-coordinate of the vertex (M1)

$$x = \frac{-3+1}{2}(=-1)$$
 OR $\frac{-4}{2\times 2}$ OR $4x + 4 = 0$ OR $2(x+1)^2 - 4$

y-coordinate of the vertex
$$= -4$$
 (A1)

correct range

$$\begin{bmatrix} -4, +\infty \begin{bmatrix} \text{OR} & y \ge -4 & \text{OR} & g \ge -4 & \text{OR} & [-4, \infty) \end{bmatrix}$$

[7 marks]

A1

(M1)
(c) equating the two functions or equations (M1)

$$g(x) = j(x) \text{ OR } px^2 + (t-1)x - p = -x+3p$$

 $px^2 + tx - 4p = 0$ (A1)
attempt to find discriminant (do not accept only in quadratic formula) (M1)
 $\Delta = t^2 + 16p^2$ A1
 $\Delta = t^2 + 16p^2 > 0$, because $t^2 \ge 0$ and $p^2 > 0$, therefore the sum will be positive R1R1
Note: Award R1 for recognising that Δ is positive and R1 for the reason.
There are two distinct points of intersection between the graphs of g and j. AG
[6 marks]
Total [16 marks]

– 16 –

(M1)

(M1)

A1

8.

(a)

(i) valid approach to find the required logarithm

$$2^{x} = \frac{1}{16}$$
 OR $2^{x} = 2^{-4}$ OR $\frac{1}{16} = 2^{-4}$ OR $\log_{2} 1 - \log_{2} 16$
 $\log_{2} \frac{1}{16} = -4$ A1

(ii) valid approach to find the required logarithm

$$9^{x} = 3$$
 OR $3^{2x} = 3$ OR $3 = 9^{\frac{1}{2}}$ OR $\frac{\log_{3} 3}{\log_{3} 9}$

 $\log_9 3 = \frac{1}{2}$

(iii)
$$(\sqrt{3})^{x} = 81 \text{ OR } \frac{\log_{3} 81}{\log_{3} \sqrt{3}}$$
 (A1)
 $(3)^{\frac{x}{2}} = 3^{4} \text{ OR } \frac{x}{2} = 4 \text{ OR } \frac{4}{\frac{1}{2}}$ (A1)
 $x = 8$ [7 marks]

continued...

– 17 –

(b) (i)

Note: There are many valid approaches to the question, and the steps may be seen in different ways. Some possible methods are given here, but candidates may use a combination of one or more of these methods.

In all methods, the final *A* mark is awarded for working which leads directly to the *AG*.

METHOD 1

$$\left(ab\right)^{3} = a \tag{A1}$$

attempt to isolate b or a power of b

correct working

$$b = \frac{a}{a^3b^2}$$
 OR $b^3 = a^{-2}$ OR $b^{-1} = (ab)^2$ OR $b^3 = \frac{1}{a^2}$

$$b = \frac{1}{a^2 b^2}$$
 OR $b = (ab)^{-2}$ OR $3\log_{ab} b = -2\log_{ab} a$ OR $-\log_{ab} b = 2\log_{ab} ab$ A1

 $\log_{ab} b = -2$

AG

(M1)

(A1)

METHOD 2

$$\left(ab\right)^{3} = a \tag{A1}$$

taking logarithm to base ab on both sides

$$\log_{ab}(ab)^3 = \log_{ab}a$$
 OR $\log_{ab}a^3b^3 = \log_{ab}a$

correct application of log rules leading to equation in terms of \log_{ab} (A1)

$$3\log_{ab} a + 3\log_{ab} b = \log_{ab} a$$
 OR $3\log_{ab} b = -2\log_{ab} a$ OR $\log_{ab} b^3 = \log_{ab} a^{-2}$

$$\log_{ab} b = \log_{ab} a^{-\frac{2}{3}}$$
 OR $\log_{ab} b = -\frac{2}{3} \log_{ab} a$ OR $\log_{ab} b = -\frac{2}{3} (3)$ A1

$$\log_{ab} b = -2$$

Note: Candidates may substitute $\log_{ab} a = 3$ at any point in their working.

continued...

(M1)



METHOD 3

$$\log_{ab} a = 3$$

writing in terms of base a

$$\frac{\log_a a}{\log_a ab} (=3)$$

correct application of log rules

$$\frac{\log_a a}{\log_a a + \log_a b} (=3) \quad \text{OR} \quad \frac{1}{1 + \log_a b} (=3) \quad \text{OR} \quad 3\log_a b = -2 \quad \text{OR}$$
$$\log_a b = -\frac{2}{3}$$

writing $\log_{ab} b$ in terms of base a

$$\log_{ab} b = \frac{\log_a b}{\log_a a + \log_a b}$$

correct working

$$\log_{ab} b = \frac{-\frac{2}{3}}{1-\frac{2}{3}} \text{ OR } \frac{\left(-\frac{2}{3}\right)}{\left(\frac{1}{3}\right)}$$

 $\log_{ab} b = -2$

AG

(M1)

(A1)

(A1)

A1

continued...

METHOD 4

$$\log_{ab} ab = 1$$

$$\log_{ab} a + \log_{ab} b = 1 \tag{A1}$$

$$3 + \log_{ab} b = 1$$

$$\log_{ab} b = -2$$

(ii) applying the quotient rule or product rule for logs

$$\log_{ab}\frac{\sqrt[3]{a}}{\sqrt{b}} = \log_{ab}\sqrt[3]{a} - \log_{ab}\sqrt{b} \quad \text{OR} \quad \log_{ab}\frac{\sqrt[3]{a}}{\sqrt{b}} = \log_{ab}\sqrt[3]{a} + \log_{ab}\frac{1}{\sqrt{b}}$$
(A1)

correct working (A1)
$$\frac{1}{1} = \frac{1}{1} = \frac{1$$

$$= \frac{1}{3} \log_{ab} a - \frac{1}{2} \log_{ab} b \text{ OR } \log_{ab} ab - \log_{ab} \sqrt{b}$$

$$= \frac{1}{3} \cdot 3 - \frac{1}{2} (-2)$$
(A1)
$$= 2$$

Note: Award A1A0A0A1 for a correct answer with no working.

[8 marks]

Total [15 marks]

(a) $\cos^2 x - 3\sin^2 x = 0$ 9.

> valid attempt to reduce equation to one involving one trigonometric function (M1)

$$\frac{\sin^2 x}{\cos^2 x} = \frac{1}{3} \quad \text{OR} \quad 1 - \sin^2 x - 3\sin^2 x = 0 \quad \text{OR} \quad \cos^2 x - 3(1 - \cos^2 x) = 0$$

OR $\cos 2x - 1 + \cos 2x = 0$

correct equation

$$\tan^{2} x = \frac{1}{3} \quad \text{OR} \quad \cos^{2} x = \frac{3}{4} \quad \text{OR} \quad \sin^{2} x = \frac{1}{4} \quad \text{OR} \quad \cos 2x = \frac{1}{2}$$
$$\tan x = \pm \frac{1}{\sqrt{3}} \quad \text{OR} \quad \cos x = \pm \frac{\sqrt{3}}{2} \quad \text{OR} \quad \sin x = (\pm)\frac{1}{2} \quad \text{OR} \quad 2x = \frac{\pi}{3} \left(, \frac{5\pi}{3}\right)$$
$$(A1)$$
$$x = \frac{\pi}{6}, \ x = \frac{5\pi}{6} \quad \text{A1A1}$$

Note: Award ***M1A1A0A1A0*** for candidates who omit the
$$\pm$$
 (for tan or cos) and give only $x = \frac{\pi}{6}$.

Award **M1A1A0A0A0** for candidates who omit the \pm (for tan or cos) and give only $x = 30^{\circ}$.

Award M1A1A1A1A0 for candidates who give both answers in degrees.

Award M1A1A1A1A0 for candidates who give both correct answers in radians, but who include additional solutions outside the domain.

Award a maximum of *M1A0A0A1A1* for correct answers with no working.

[5 marks]

(A1)

A1A1

continued...

(b) (i) attempt to use the chain rule (may be evidenced by at least one $\cos x \sin x$ term) (*M1*)

- 23 -

$$f'(x) = -2\cos x \sin x - 6\sin x \cos x (= -8\sin x \cos x = -4\sin 2x)$$
 A1

(ii) valid attempt to solve their f'(x) = 0

At least 2 correct *x*-coordinates (may be seen in coordinates) (A1)

$$x = 0$$
, $x = \frac{\pi}{2}$, $x = \pi$

Note: Accept additional correct solutions outside the domain.

Award **A0** if any additional incorrect solutions are given.

correct coordinates (may be seen in graph for part (c))

A1A1A1

(M1)

$$(0,1), (\pi,1), (\frac{\pi}{2},-3)$$

Note: Award a maximum of *M1A1A1A1A0* if any additional solutions are given.

Note: If candidates do not find at least two correct x-coordinates, it is possible to award the appropriate final marks for their correct coordinates, such as *M1A0A0A1A0.*

[7 marks]

continued...



Note: The final two marks may be awarded independently of each other.

[3 marks]

Total [15 marks]

– 24 –

M22/5/MATHX/SP1/ENG/TZ2/XX/M



Markscheme

May 2022

Mathematics: analysis and approaches

Standard level

Paper 1





© International Baccalaureate Organization 2022

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2022

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2022

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *MO* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies *A3*, *M2 etc.*, do **not** split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this working is incorrect and/or suggests a misunderstanding of the question. This will encourage a uniform approach to marking, with less examiner discretion. Although some candidates may be advantaged for that specific question item, it is likely that these candidates will lose marks elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used in a subsequent part. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.

- 3 -

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	8√2	5.65685 (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)
2.	$\frac{35}{72}$	0.468111 (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)

3 Implied marks

Implied marks appear in **brackets e.g.** (M1), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is (*M1*)*A1*, it is possible to award full marks for *their* correct answer, without working being seen. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.

- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (MR). A candidate should be penalized only once for a particular misread. Use the MR stamp to indicate that this has been a misread and do not award the first mark, even if this is an M mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** ... **OR**.

7 Alternative forms

Unless the question specifies otherwise, accept equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, *M* marks and intermediate *A* marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "from the use of 3 sf values".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an *A* mark to be awarded, arithmetic should be completed, and

any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$.

An exception to this is simplifying fractions, where lowest form is not required (although the

numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or

written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate A marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".



Section A

1.	(a)	g(0) = -2	A1
			[1 mark]
	(b)	evidence of using composite function $f(g(0))$ OR $f(-2)$	(M1)
		$(f \circ g)(0) = 8$	A1
			[2 marks]
	(c)	<i>x</i> = 3	A2
			[2 marks]
			Total [5 marks]



- 8 -

2.	(a)	$u_1 = 12$	A1
			[1 mark]
	(b)	15 - 3n = -33	(A1)
	()	<i>n</i> =16	A1
			[2 marks]

(c)	valid approach to find d	(M1)
	$u_2 - u_1 = 9 - 12$ OR recognize gradient is -3 OR attempts to solve $-33 = 12 + 15d$	
	d = -3	A1
		[2 marks]
		Total [5 marks]

3. (a)
$$(n-1)+n+(n+1)$$
 (A1)

=3n

which is always divisible by 3

A1

AG

[2 marks]

(b)
$$(n-1)^2 + n^2 + (n+1)^2$$
 (= $n^2 - 2n + 1 + n^2 + n^2 + 2n + 1$) **A1**

attempts to expand either
$$(n-1)^2$$
 or $(n+1)^2$ (do not accept n^2-1 or n^2+1) (M1)

$$=3n^2+2$$
 A1

demonstrating recognition that 2 is not divisible by 3 or $\frac{2}{3}$ seen after correct expression divided by 3

 $3n^2$ is divisible by 3 and so $3n^2 + 2$ is never divisible by 3 OR the first term is divisible by 3, the second is not

OR
$$3\left(n^2 + \frac{2}{3}\right)$$
 OR $\frac{3n^2 + 2}{3} = n^2 + \frac{2}{3}$

hence the sum of the squares is never divisible by 3

AG [4 marks] Total [6 marks]





A1

A1

[2 marks]



continued...

(c) $x > \frac{1}{2}$

rational function shape with two branches in opposite quadrants, with two correctly positioned asymptotes and asymptotic behaviour shown

Note: The equations of the asymptotes are not required on the graph provided there is a clear indication of asymptotic behaviour at x = -1 and y = 2 (or at their FT asymptotes from part (a)).

axes intercepts clearly shown at
$$x = \frac{1}{2}$$
 and $y = -1$ A1A1

A1

A1

Note: Accept correct alternative correct notation, such as	$\left(\frac{1}{2},\infty\right)$	and	$\left \frac{1}{2},\infty\right $	[.
--	-----------------------------------	-----	-----------------------------------	----

[1 mark] Total [6 marks]

determines
$$\frac{\pi}{4}$$
 (or 45°) as the first quadrant (reference) angle (A1)
attempts to solve $\frac{x}{2} + \frac{\pi}{3} = \frac{\pi}{4}$ (M1)
Note: Award M1 for attempting to solve $\frac{x}{2} + \frac{\pi}{3} = \frac{\pi}{4}, \frac{7\pi}{4}, \dots$
 $\frac{x}{2} + \frac{\pi}{3} = \frac{\pi}{4} \Rightarrow x < 0$ and so $\frac{\pi}{4}$ is rejected (R1)
 $\frac{x}{2} + \frac{\pi}{3} = 2\pi - \frac{\pi}{4} \left(= \frac{7\pi}{4} \right)$ A1
 $x = \frac{17\pi}{6}$ (must be in radians) A1
[5 marks]

– 13 –

5.



(M1)

A1

6. (a) EITHER

recognises the required term (or coefficient) in the expansion

$$bx^{5} = {}^{7}C_{2} x^{5} 1^{2} \quad \text{OR} \quad b = {}^{7}C_{2} \quad \text{OR} \quad {}^{7}C_{5}$$
$$b = \frac{7!}{2!5!} \left(= \frac{7!}{2!(7-2)!} \right)$$

correct working

 $\frac{7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1}{2 \times 1 \times 5 \times 4 \times 3 \times 2 \times 1} \quad \text{OR} \quad \frac{7 \times 6}{2!} \quad \text{OR} \quad \frac{42}{2}$

OR

lists terms from row 7 of Pascal's triangle	(M1)
1,7,21,	A1

THEN

AG	b = 21
[2 marks]	

(b)	<i>a</i> = 7	(A1)
	correct equation	A1

$$21x^5 = \frac{ax^6 + 35x^4}{2} \quad \text{OR} \quad 21x^5 = \frac{7x^6 + 35x^4}{2}$$

correct quadratic equation A1 $7x^2 - 42x + 35 = 0$ OR $x^2 - 6x + 5 = 0$ (or equivalent)

valid attempt to solve their quadratic

$$(x-1)(x-5)=0$$
 OR $x=\frac{6\pm\sqrt{(-6)^2-4(1)(5)}}{2(1)}$

$$x = 1, x = 5$$

Note: Award final **A0** for obtaining x = 0, x = 1, x = 5.

[5 marks] Total [7 marks]

(M1)

A1

Section B

(a) *x*=3 7. **Note:** Must be an equation in the form "x = ". Do not accept 3 or $\frac{-b}{2a} = 3$. [1 mark]

(b) (i)
$$h=3, k=4$$
 (accept $a(x-3)^2+4$) A1A1

(ii) attempt to substitute coordinates of Q (M1)

$$12 = a(5-3)^2 + 4$$
, $4a+4=12$
 $a=2$ A1
[4 marks]

(c)	recognize need to find derivative of f	(M1)
	f'(x) = 4(x-3) or $f'(x) = 4x-12$	A1
	f'(5) = 8 (may be seen as gradient in their equation)	(A1)
	y-12=8(x-5) or $y=8x-28$	A1
Not	e: Award A0 for $L = 8x - 28$.	
		[4 marks] continued

A1

(d)	METHOD 1	
	Recognizing that for g to be increasing, $f(x) - d > 0$, or $g' > 0$	(M1)
	The vertex must be above the <i>x</i> -axis, $4 - d > 0$, $d - 4 < 0$	(R1)
	d < 4	A1
		[3 marks]
	METHOD 2	
	attempting to find discriminant of g'	(M1)
	$(-12)^2 - 4(2)(22 - d)$	
	recognizing discriminant must be negative	(R1)
	$-32+8d<0$ OR $\Delta<0$	
	d < 4	A1
		[3 marks]
(e)	recognizing that for g to be concave up, $g'' > 0$	(M1)
	g'' > 0 when $f' > 0$, $4x - 12 > 0$, $x - 3 > 0$	(R1)
	<i>x</i> >3	A1
		[3 marks]
		Total [15 marks]

8. (a)
$$\frac{1}{x-4} + 1 = x-3$$
 (M1)

$$x^{2}-8x+15=0$$
 OR $(x-4)^{2}=1$ (A1)

valid attempt to solve their quadratic

$$(x-3)(x-5)=0$$
 OR $x=\frac{8\pm\sqrt{8^2-4(1)(15)}}{2(1)}$ OR $(x-4)=\pm 1$

x=5 (x=3, x=5) (may be seen in answer)

$$B(5, 2)$$
 (accept $x = 5, y = 2$)

A1

(M1)

A1

[5 marks]

continued...



(b) recognizing two correct regions from x=3 to x=5 and from x=5 to x=k (R1) triangle $+\int_{5}^{k} f(x)dx \quad OR \quad \int_{3}^{5} g(x)dx + \int_{5}^{k} f(x)dx \quad OR \quad \int_{3}^{5} (x-3)dx + \int_{5}^{k} \left(\frac{1}{x-4}+1\right)dx$ area of triangle is $2 \quad OR \quad \frac{2 \cdot 2}{2} \quad OR \quad \left(\frac{5^{2}}{2}-3(5)\right) - \left(\frac{3^{2}}{2}-3(3)\right)$ (A1)

correct integration

$$\int \left(\frac{1}{x-4} + 1\right) dx = \ln(x-4) + x (+C)$$

Note: Award **A1** for $\ln(x-4)$ and **A1** for *x*. **Note:** The first three **A** marks may be awarded independently of the **R** mark.

substitution of their limits (for <i>x</i>) into their integrated function (in terms of <i>x</i>)	(M1)
$\ln(k-4)+k-(\ln 1+5)$	
$\left[\ln(x-4)+x\right]_{5}^{k}=\ln(k-4)+k-5$	A1
adding their two areas (in terms of k) and equating to $\ln p + 8$	(M1)
$2 + \ln(k-4) + k - 5 = \ln p + 8$	
equating their non-log terms to 8 (equation must be in terms of k)	(M1)
k-3=8	
k=11	A1
11 - 4 = p	
p = 7	A1

[10 marks] Total [15 marks]

(A1)(A1)

9.	(a)	uses $\sum P(X = x) = 1$ to form a linear equation in p and q	(M1)
		correct equation in terms of p and q from summing to 1	A1
		p + 0.3 + q + 0.1 = 1 OR $p + q = 0.6$ (or equivalent)	
		uses $E(X) = 2$ to form a linear equation in p and q	(M1)
		correct equation in terms of p and q from $E(X) = 2$	A1
		p + 0.6 + 3q + 0.4 = 2 OR $p + 3q = 1$ (or equivalent)	

– 19 –

Note: The marks for using $\sum P(X = x) = 1$ and the marks for using E(X) = 2 may be awarded independently of each other.

evidence of correctly solving these equations simultaneously	A1
for example, $2q = 0.4 \Rightarrow q = 0.2$ or $p + 3 \times (0.6 - p) = 1 \Rightarrow p = 0.4$	
so $p = 0.4$ and $q = 0.2$	AG
	[5 marks]

(M1)	valid approach	(b)
	P(X > 2) = P(X = 3) + P(X = 4) OR $P(X > 2) = 1 - P(X = 1) - P(X = 2)$	
A1	= 0.3	
[2 marks]		
continued		

(c) recognises at least one of the valid scores (6, 7, or 8) required to win the game (M1)

Note: Award *M0* if candidate also considers scores other than 6, 7, or 8 (such as 5).

I	et T represent the score on the last two rolls				
ä					
	$P(T=6) = 2(0.3)(0.1) + (0.2)^2 (=0.1)$				
ä	a score of 7 is obtained by rolling $(3,4)$ or $(4,3)$				
	P(T=7) = 2(0.2)(0.1) (=0.04)				
a score of 8 is obtained by rolling $(4,4)$					
	$P(T=8) = (0.1)^2 (=0.01)$	A1			
Note:	Note: The above 3 A1 marks are independent of each other.				
	P(Nickywins) = 0.1 + 0.04 + 0.01				
	=0.15	A1 [5 marks]			
		[o markoj			
(d)	3+ <i>b</i> =8	(M1)			
	<i>b</i> =5	A1			
		[2 marks]			
		continued			

(e) **METHOD 1** EITHER $P(S=5) = \frac{4}{16}$ $P(S=a+2) = \frac{4}{16}$ A1 $\Rightarrow a + 2 = 5$ OR $P(S=6) = \frac{3}{16}$ $P(S = a+3) = \frac{2}{16}$ and $P(S = 5+1) = \frac{1}{16}$ A1 $\Rightarrow a+3=6$ OR $P(S=4) = \frac{3}{16}$ $P(S = a+1) = \frac{2}{16}$ and $P(S = 1+3) = \frac{1}{16}$ A1 $\Rightarrow a+1=4$ THEN $\Rightarrow a = 3$ **A1**

Note: Award **A0A0** for a = 3 obtained without working/reasoning/justification.

[2 marks] continued...

METHOD 2

EITHER

correctly lists a relevant part of the sample space

for example, $\{S = 4\} = \{(3,1), (1,a), (1,a)\}$ or $\{S = 5\} = \{(2,a), (2,a), (2,a), (2,a)\}$

or
$$\{S = 6\} = \{(3, a), (3, a), (1, 5)\}$$

 $a + 3 = 6$

OR

eliminates possibilities (exhaustion) for a < 5convincingly shows that $a \ne 2,4$ $a \ne 4$, for example, $P(S = 7) = \frac{2}{16}$ from (2,5), (2,5) and so

 $(3,a),(3,a) \Rightarrow a+3 \neq 7$

THEN

 $\Rightarrow a = 3$

A1 [2 marks] Total [16 marks]

A1

A1



Markscheme

May 2022

Mathematics: analysis and approaches

Standard level

Paper 1

25 pages



© International Baccalaureate Organization 2022

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2022

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2022

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *MO* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies *A3*, *M2 etc.*, do **not** split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this working is incorrect and/or suggests a misunderstanding of the question. This will encourage a uniform approach to marking, with less examiner discretion. Although some candidates may be advantaged for that specific question item, it is likely that these candidates will lose marks elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used in a subsequent part. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.

- 3 -

Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action
1.	8√2	5.65685 (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)
2.	$\frac{35}{72}$	0.468111 (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)

3 Implied marks

Implied marks appear in **brackets e.g.** (M1), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is (*M1*)*A1*, it is possible to award full marks for *their* correct answer, without working being seen. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.

- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (MR). A candidate should be penalized only once for a particular misread. Use the MR stamp to indicate that this has been a misread and do not award the first mark, even if this is an M mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** ... **OR**.
7 Alternative forms

Unless the question specifies otherwise, accept equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, *M* marks and intermediate *A* marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "from the use of 3 sf values".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an *A* mark to be awarded, arithmetic should be completed, and

any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$.

An exception to this is simplifying fractions, where lowest form is not required (although the

numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or

written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate A marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".



[3 marks]

- 8 -

1. (a)
$$m_{BC} = \frac{12-6}{-14-4} \left(= -\frac{1}{3} \right)$$
 (A1)
finding $m_L = \frac{-1}{m_{BC}}$ using their m_{BC} (M1)
 $m_L = 3$
 $y - 20 = 3(x+2), y = 3x + 26$ A1
Note: Do not accept $L = 3x + 26$.

(b) substituting (k, 2) into their *L* (M1) 2-20=3(k+2) OR 2=3k+26 k=-8 A1 [2 marks] Total [5 marks]

2. (a)
$$\frac{3\sqrt{x}-5}{\sqrt{x}} = 3-5x^{-\frac{1}{2}}$$

 $p = -\frac{1}{2}$ A1

- 9 -

[1 mark]

(b)
$$\int \frac{3\sqrt{x}-5}{\sqrt{x}} dx = 3x - 10x^{\frac{1}{2}} (+c)$$
 A1A1

substituting limits into their integrated function and subtracting (M1)

$$3(9) - 10(9)^{\frac{1}{2}} - \left(3(1) - 10(1)^{\frac{1}{2}}\right) \text{ OR } 27 - 10 \times 3 - (3 - 10)$$

=4

A1 [4 marks] Total [5 marks]

3.	(a)	IQR = 10 - 6 (= 4)	(A1)
		attempt to find $Q_3 + 1.5 \times IQR$	(M1)
		10+6	
		16	A1

(b) (i) choosing
$$c = \frac{1}{2}a - 9$$
 (M1)
 $\frac{1}{2} \times 42 - 9$
 $= 12$ (years old) A1
(ii) attempt to solve system by substitution or elimination (M1)
34 (years old) A1
[4 marks]
Total [7 marks]

4. (a)
$$(f \circ g)(x) = f(2x)$$
 (A1)
 $f(2x) = \sqrt{3} \sin 2x + \cos 2x$ A1

(b)
$$\sqrt{3} \sin 2x + \cos 2x = 2\cos 2x$$

 $\sqrt{3} \sin 2x = \cos 2x$
recognising to use tan or cot
 $\tan 2x = \frac{1}{\sqrt{3}}$ OR $\cot 2x = \sqrt{3}$ (values may be seen in right triangle) (A1)
 $\left(\arctan\left(\frac{1}{\sqrt{3}}\right) = \right) \frac{\pi}{6}$ (seen anywhere) (accept degrees) (A1)
 $2x = \frac{\pi}{6}, \frac{7\pi}{6}$
 $x = \frac{\pi}{12}, \frac{7\pi}{12}$ A1A1

Note: Do not award the final **A1** if any additional solutions are seen. Award **A1A0** for correct answers in degrees. Award **A0A0** for correct answers in degrees with additional values.

> [5 marks] Total [7 marks]

5.	evidence of using product rule	(M1)
	$\frac{\mathrm{d}y}{\mathrm{d}x} = (2x-1) \times \left(k\mathrm{e}^{kx}\right) + 2 \times \mathrm{e}^{kx} \left(=\mathrm{e}^{kx}(2kx-k+2)\right)$	A1

correct working for one of (seen anywhere)

OR

slope of tangent is $5e^k$

their $\frac{dy}{dx}$ at x = 1 equals the *slope* of $y = 5e^k x$ (= $5e^k$) (seen anywhere) (M1) $ke^k + 2e^k = 5e^k$ k = 3A1



A1



6. (a) translation (shift) by $\frac{3\pi}{2}$ to the right/positive horizontal direction A1 translation (shift) by q upwards/positive vertical direction A1 Note: accept translation by (

	$\left(\begin{array}{c} \frac{3\pi}{2} \\ q \end{array}\right)$
Do not accept 'move	e' for translation/shift.

[2 marks]

(b) METHOD 1

minimum of $4\sin\left(x-\frac{3\pi}{2}\right)$ is -4 (may be seen in sketch)	(M1)
$-4 + 2.5 + q \ge 7$	
$q \ge 8.5$ (accept $q = 8.5$)	A1
substituting $x=0$ and their q (=8.5) to find r	(M1)
$(r=) 4\sin\left(\frac{-3\pi}{2}\right) + 2.5 + 8.5$	
4+2.5+8.5	(A1)
smallest value of r is 15	A1
	continued

Question 6 continued

METHOD 2

substituting	x = 0 to fin	nd an expressio	n (for r) in terms of a	a l	M1	Ì
Substituting	$\lambda = 0$ to m	iu ali explessio			9	1111)	,

$$\left(g(0)=r=\right) \quad 4\sin\left(\frac{-3\pi}{2}\right)+2.5+q$$

$$(r=)$$
 6.5+q A1

minimum of
$$4\sin\left(x-\frac{3\pi}{2}\right)$$
 is -4 (M1)

$$-4 + 2.5 + q \ge 7$$

$$-4+2.5+(r-6.5) \ge 7$$
 (accept =) (A1)

smallest value of r is 15

METHOD 3

$$4\sin\left(x - \frac{3\pi}{2}\right) + 2.5 + q = 4\cos x + 2.5 + q$$
y-intercept of $4\cos x + 2.5 + q$ is a maximum
(M1)

y-intercept of $4\cos x + 2.5 + q$ is a maximum

amplitude of g(x) is 4

attempt to find least maximum

$$r = 2 \times 4 + 7$$

smallest value of r is 15

A1

(A1)

(M1)

A1

[5 marks] Total [7 marks]

Section B

7. (a) EITHER

attempt to use $x = -\frac{b}{2a}$	(M1)
$q = -\frac{-12}{2 \times 3}$	

attempt to complete the square (M1) $3(x-2)^2-12+p$

OR

attempt to differentiate and equate to 0	(M1)
f''(x) = 6x - 12 = 0	

THEN

q = 2

A1 [2 marks] continued...

Question 7 continued

(b)	(i)	discriminant $= 0$	A1
	(ii)	EITHER	
		attempt to substitute into $b^2 - 4ac$	(M1)
		$(-12)^2 - 4 \times 3 \times p = 0$	A1
		OR	
		f'(2) = 0	(M1)
		-12 + p = 0	A1
		THEN	
		<i>p</i> =12	A1
			[4 marks]
(c)	f"(.	x) = 6x - 12	A1
()	atter	mpt to find $f''(0)$	(M1)
	=62	×0-12	()
	grac	lient = -12	A1
	-		[3 marks]
			continued

– 16 –

Question 7 continued

(d)



A1A1

Note	e: A	ward A1 for line with positive gradient, A1 for correct intercepts.	
			[2 marks]
(e)	(i)	<i>a</i> = 2	A1
	(ii)	x < 2 f''(x) < 0 (for $x < 2$) OR the f'' is below the x-axis (for $x < 2$)	A1
		$OR \xleftarrow{-} + \\ 2 \qquad \qquad f'' \text{ (sign diagram must be labelled } f'')$	R1

[3 marks] Total [14 marks]

8. (a) (i) EITHER

attempt to use a ratio from consecutive terms

$$\frac{p\ln x}{\ln x} = \frac{\frac{1}{3}\ln x}{p\ln x} \quad \text{OR} \quad \frac{1}{3}\ln x = (\ln x)r^2 \quad \text{OR} \quad p\ln x = \ln x \left(\frac{1}{3p}\right)$$

Note: Candidates may use $\ln x^1 + \ln x^p + \ln x^{\frac{1}{3}} + \dots$ and consider the powers of x in geometric sequence.

Award **M1** for $\frac{p}{1} = \frac{\overline{3}}{p}$.

OR

$$r = p$$
 and $r^2 = \frac{1}{3}$

М1

М1

THEN

$$p^{2} = \frac{1}{3} \text{ OR } r = \pm \frac{1}{\sqrt{3}}$$
 A1
 $p = \pm \frac{1}{\sqrt{3}}$ AG

Note: Award *MOA0* for $r^2 = \frac{1}{3}$ or $p^2 = \frac{1}{3}$ with no other working seen.

(ii)
$$\frac{\ln x}{1 - \frac{1}{\sqrt{3}}} (= 3 + \sqrt{3})$$
 (A1)

$$\ln x = 3 - \frac{3}{\sqrt{3}} + \sqrt{3} - \frac{\sqrt{3}}{\sqrt{3}} \quad \text{OR} \quad \ln x = 3 - \sqrt{3} + \sqrt{3} - 1 \quad (\Rightarrow \ln x = 2)$$

$$x = e^2$$

[5 marks]

continued...

М1

A1

Question 8 continued

(b) (i) **METHOD 1**

attempt to find a difference from consecutive terms or from u_2

correct equation

$$p \ln x - \ln x = \frac{1}{3} \ln x - p \ln x$$
 OR $\frac{1}{3} \ln x = \ln x + 2(p \ln x - \ln x)$

Note: Candidates may use $\ln x^1 + \ln x^p + \ln x^{\frac{1}{3}} + \dots$ and consider the powers of x in arithmetic sequence.

Award ***M1A1*** for
$$p - 1 = \frac{1}{3} - p$$

METHOD 2

attempt to use arithmetic mean
$$u_2 = \frac{u_1 + u_3}{2}$$
 M1

$$p\ln x = \frac{\ln x + \frac{1}{3}\ln x}{2}$$

$$2p\ln x = \frac{4}{3}\ln x \quad \left(\Rightarrow 2p = \frac{4}{3}\right)$$

$$p = \frac{2}{3}$$
 AG

continued...

– 19 –

Question 8 continued

METHOD 3

attempt to find difference using u_3

$$\frac{1}{3}\ln x = \ln x + 2d \quad \left(\Rightarrow d = -\frac{1}{3}\ln x\right)$$

$$u_2 = \ln x + \frac{1}{2} \left(\frac{1}{3} \ln x - \ln x \right)$$
 OR $p \ln x - \ln x = -\frac{1}{3} \ln x$ A1

$$p\ln x = \frac{2}{3}\ln x$$

$$p = \frac{2}{3}$$
 AG

(ii)
$$d = -\frac{1}{3} \ln x$$

A1

М1

continued...

Question 8 continued

(iii) **METHOD 1**

$$S_n = \frac{n}{2} \left[2\ln x + (n-1) \times \left(-\frac{1}{3}\ln x \right) \right]$$

attempt to substitute into S_{μ}	$_{n}$ and equate to $-3\ln x$	(M1)
--------------------------------------	--------------------------------	------

 $\frac{n}{2}\left[2\ln x + (n-1)\times\left(-\frac{1}{3}\ln x\right)\right] = -3\ln x$

correct working with $S_{\scriptscriptstyle n}$ (seen anywhere)

$$\frac{n}{2} \left[2\ln x - \frac{n}{3} \ln x + \frac{1}{3} \ln x \right] \quad \text{OR} \quad n\ln x - \frac{n(n-1)}{6} \ln x \quad \text{OR}$$
$$\frac{n}{2} \left(\ln x + \left(\frac{4-n}{3}\right) \ln x \right)$$

correct equation without $\ln x$

$$\frac{n}{2}\left(\frac{7}{3}-\frac{n}{3}\right) = -3 \quad \text{OR} \quad n - \frac{n(n-1)}{6} = -3 \quad \text{or equivalent}$$

Note: Award as above if the series $1 + p + \frac{1}{3} + \dots$ is considered leading to $\frac{n}{2}\left(\frac{7}{3}-\frac{n}{3}\right)=-3.$

attempt to form a quadratic = 0

$$n^2 - 7n - 18 = 0$$

attempt to solve their quadratic (M1)

attempt to solve their quadratic

n = 9

$$(n-9)(n+2) = 0$$

continued...

A1

(M1)

(A1)

Question 8 continued

METHOD 2

listing the first 7 terms of the sequence	(A1)
$\ln x + \frac{2}{3}\ln x + \frac{1}{3}\ln x + 0 - \frac{1}{3}\ln x - \frac{2}{3}\ln x - \ln x + \dots$	
recognizing first 7 terms sum to 0	М1
8 th term is $-\frac{4}{3}\ln x$	(A1)
9 th term is $-\frac{5}{3}\ln x$	(A1)
sum of 8 th and 9 th terms $= -3\ln x$	(A1)
<i>n</i> = 9	A1
	[10 marks]
	Total [15 marks]

9.	(a)	(i)	EITHER	
			attempt to use binomial expansion	(M1)
			$1 + {}^{3}C_{1} \times 1 \times (-a) + {}^{3}C_{2} \times 1 \times (-a)^{2} + 1 \times (-a)^{3}$	
			OR	
			(1-a)(1-a)(1-a)	
			$=(1-a)(1-2a+a^2)$	(M1)
			THEN	
			$=1-3a+3a^2-a^3$	A1
		(ii)	$a = \cos 2x$	(A1)
			So, $1 - 3\cos 2x + 3\cos^2 2x - \cos^3 2x =$	
			$(1-\cos 2x)^3$	A1
			attempt to substitute any double angle rule for $\cos 2x$ into $(1 - \cos 2x)^3$	(M1)
			$=(2\sin^2 x)^3$	A1
			$=8\sin^6 x$	AG
[Note	e: All	ow working RHS to LHS.	
				[6 marks] continued

– 23 –

Question 9 continued

(b) (i) recognizing to integrate
$$\int (4\cos x \times 8\sin^6 x) dx$$
 (M1)

EITHER

applies integration by inspection (M1)

$$32\int \left(\cos x \times (\sin x)^6\right) \mathrm{d}x$$

$$=\frac{32}{7}\sin^7 x (+c)$$
 A1

$$\left[\frac{32}{7}\sin^7 x\right]_0^m \quad \left(=\frac{32}{7}\sin^7 m - \frac{32}{7}\sin^7 0\right)$$

OR

$$u = \sin x \Longrightarrow \frac{du}{dx} = \cos x$$

$$\int 32\cos x (\sin^6 x) dx = \int 32u^6 du$$
(M1)

$$=\frac{32}{7}u^{7}(+c)$$

$$\left[\frac{32}{7}\sin^7 x\right]_0^m \text{ OR } \left[\frac{32}{7}u^7\right]_0^{\sin m} \left(=\frac{32}{7}\sin^7 m - \frac{32}{7}\sin^7 0\right)$$
 A1

THEN

$$=\frac{32}{7}\sin^7 m$$

continued...

Question 9 continued

EITHER (ii)

$$\frac{32}{7}\sin^7\frac{\pi}{2} - \frac{32}{7}\sin^7 m = \frac{127}{28} \quad \text{OR} \quad \frac{32}{7}\left(1 - \sin^7 m\right) = \frac{127}{28} \tag{M1}$$

$$\int_{0}^{\frac{\pi}{2}} f(x) dx = \int_{0}^{m} f(x) dx + \int_{m}^{\frac{\pi}{2}} f(x) dx$$

$$\frac{32}{7} = \frac{32}{7} \sin^{7} m + \frac{127}{28}$$
(M1)
THEN

THEN

 $m = \frac{\pi}{6}$

2

$$\sin^{7} m = \frac{1}{128} \left(= \frac{1}{2^{7}} \right)$$
(A1)
$$\sin m = \frac{1}{2}$$
(A1)

A1

[9 marks] Total [15 marks]



Markscheme

November 2021

Mathematics: analysis and approaches

Standard level

Paper 1

23 pages



© International Baccalaureate Organization 2021

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2021

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2021

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- A Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *M0* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies *A3*, *M2 etc.*, do **not** split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this working is incorrect and/or suggests a misunderstanding of the question. This will encourage a uniform approach to marking, with less examiner discretion. Although some candidates may be advantaged for that specific question item, it is likely that these candidates will lose marks elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used in a subsequent part. For example, when a correct exact value is followed by an incorrect decimal approximation in the first part and this approximation is then used in the second part. In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part. Examples:

	Correct answer seen	Further working seen	Any FT issues?	Action	
1.	8√2	5.65685 (incorrect decimal value)	No. Last part in question.	Award A1 for the final mark (condone the incorrect further working)	
2.	$\frac{35}{72}$	0.468111 (incorrect decimal value)	Yes. Value is used in subsequent parts.	Award A0 for the final mark (and full FT is available in subsequent parts)	

3 Implied marks

Implied marks appear in **brackets e.g.** (*M1*), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is (*M1*)*A1*, it is possible to award full marks for *their* correct answer, without working being seen. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.

- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (MR). A candidate should be penalized only once for a particular misread. Use the MR stamp to indicate that this has been a misread and do not award the first mark, even if this is an M mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does not constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by **EITHER** ... **OR**.

7 Alternative forms

Unless the question specifies otherwise, accept equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, *M* marks and intermediate *A* marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "*from the use of 3 sf values*".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an **A** mark to be awarded, arithmetic should be completed, and any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as $\frac{5}{2}$. An exception to this is simplifying fractions, where lowest form is not required (although the

numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left in this form or

written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate *A* marks do NOT need to be simplified.

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".



Section A

1.	(a)	(i)	setting $f(x) = 0$	(M1)
			x = 1, $x = -3$ (accept $(1,0), (-3,0)$)	A1
		(ii)	METHOD 1	
			x = -1	A1
			substituting their x-coordinate into f	(M1)
			<i>y</i> = 8	A1
			(-1,8)	
			METHOD 2	
			attempt to complete the square	(M1)
			$-2\left(\left(x+1\right)^2-4\right)$	
			x = -1, y = 8	A1A1
			(-1,8)	
				[5 marks]
	(b)	h = -	4	A1
		<i>k</i> = 8		A1
				[2 marks]
				Total [7 marks]

2. recognition that
$$y = \int \cos\left(x - \frac{\pi}{4}\right) dx$$
 (M1)

$$y = \sin\left(x - \frac{\pi}{4}\right) \ (+c) \tag{A1}$$

substitute both x and y values into their integrated expression including c (M1)

$$2 = \sin\frac{\pi}{2} + c$$

$$c = 1$$

$$y = \sin\left(x - \frac{\pi}{4}\right) + 1$$
[4 marks]



3. (a) (i)
$$x=3$$

(ii) $y=-2$ **A1**

– 10 –

[2 marks]

(b) (i)
$$(-2,0)$$
 (accept $x = -2$) A1

(ii)
$$\left(0,\frac{4}{3}\right)$$
 (accept $y = \frac{4}{3}$ and $f(0) = \frac{4}{3}$) **A1**

[2 marks] continued...

GAT PRESS

Question 3 continued.

(C)



– 11 –

4. (a) valid approach to find P(R) (M1) tree diagram (must include probability of picking box) with correct required probabilities $OR P(R \cap B_1) + P(R \cap B_2) OR P(R|B_1)P(B_1) + P(R|B_2)P(B_2)$

$$\frac{5}{7} \cdot \frac{1}{2} + \frac{4}{7} \cdot \frac{1}{2}$$
(A1)

$$P(R) = \frac{9}{14}$$
[3 marks]

(b) events *A* and *R* are not independent, since $\frac{9}{14} \cdot \frac{1}{2} \neq \frac{5}{14}$ OR $\frac{5}{7} \neq \frac{9}{14}$ OR $\frac{5}{9} \neq \frac{1}{2}$ OR an explanation e.g. different number of red balls in each box

Note: Both conclusion and reasoning are required. Do not split the **A2**.

[2 marks] Total [5 marks]

A2



5. (a) f'(4) = 6(b) $f(4) = 6 \times 4 = 1 = 23$ A1 [1 mark]

(b)
$$f(4) = 6 \times 4 - 1 = 25$$
 [1 mark]

(c)
$$h(4) = f(g(4))$$

 $h(4) = f(4^2 - 3 \times 4) = f(4)$
(M1)

$$h(4) = 23$$
 A1

(d) attempt to use chain rule to find
$$h'$$
 (M1)
 $f'(g(x)) \times g'(x) \quad \text{OR} \quad (x^2 - 3x)' \times f'(x^2 - 3x)$

$$h'(4) = (2 \times 4 - 3) f'(4^2 - 3 \times 4)$$

= 30

$$y-23=30(x-4)$$
 OR $y=30x-97$

A1 [3 marks] Total [7 marks]

A1

(a) METHOD 1 6.

attempt to write all LHS terms with a common denominator of $x-1$	(M1)
$2x-3-\frac{6}{x-1} = \frac{2x(x-1)-3(x-1)-6}{x-1} \text{OR} \frac{(2x-3)(x-1)}{x-1} - \frac{6}{x-1}$	
$=\frac{2x^2-2x-3x+3-6}{x-1} \text{OR} \frac{2x^2-5x+3}{x-1}-\frac{6}{x-1}$	A1
$2r^2 - 5r - 3$	

$$=\frac{2x-5x-5}{x-1}$$
 AG

METHOD 2

attempt to use algebraic division on RHS	(M1)
correctly obtains quotient of $2x-3$ and remainder -6	A1
$=2x-3-\frac{6}{3}$ as required.	AG

$$=2x-3-\frac{6}{x-1}$$
 as required.

[2 marks] continued...



stion 6	S continued.	
(b)	consider the equation $\frac{2\sin^2 2\theta - 5\sin 2\theta - 3}{\sin 2\theta - 1} = 0$	(M1)
	$\Rightarrow 2\sin^2 2\theta - 5\sin 2\theta - 3 = 0$	
	EITHER	
	attempt to factorise in the form $(2\sin 2\theta + a)(\sin 2\theta + b)$	(M1)
	Note: Accept any variable in place of $\sin 2\theta$.	
	$(2\sin 2\theta + 1)(\sin 2\theta - 3) = 0$	
	OR	
	attempt to substitute into quadratic formula	(M1)
	$\sin 2\theta = \frac{5 \pm \sqrt{49}}{4}$	
	THEN	
	$\sin 2\theta = -\frac{1}{2} \text{ or } \sin 2\theta = 3$	(A1)
	Note: Award A1 for $\sin 2\theta = -\frac{1}{2}$ only.	
	one of $\frac{7\pi}{6}$ OR $\frac{11\pi}{6}$ (accept 210 or 330)	(A1)
	$\theta = \frac{7\pi}{12}, \frac{11\pi}{12}$ (must be in radians)	A1

Note: Award A0 if additional answers given.

[5 marks] Total [7 marks]

Questic

Section B

7. (a) (i) valid approach to find turning point (
$$v' = 0$$
, $-\frac{b}{2a}$, average of roots) (M1)

(ii) attempt to integrate v (M1)

$$\int v \, dt = \int (4 + 4t - 3t^2) \, dt = 4t + 2t^2 - t^3(+c)$$
A1A1
Note: Award A1 for $4t + 2t^2$, A1 for $-t^3$.
attempt to substitute their t into their solution for the integral (M1)
distance $= 4\left(\frac{2}{3}\right) + 2\left(\frac{2}{3}\right)^2 - \left(\frac{2}{3}\right)^3$
 $= \frac{8}{3} + \frac{8}{9} - \frac{8}{27}$ (or equivalent)
 $= \frac{88}{27}$ (m)
AG
[7 marks] continued...

[7 marks] continued...
Question 7 continued.

(b)



– 17 –

valid approach to solve $4 + 4t - 3t^2 = 0$ (may be seen in part (a))

$$(2-t)(2+3t)$$
 OR $\frac{-4\pm\sqrt{16+48}}{-6}$

correct *x*- intercept on the graph at t = 2

Note: The following two **A** marks may only be awarded if the shape is a concave down parabola. These two marks are independent of each other and the (*M1*).

correct domain from 0 to 3 starting at (0,4)

Note: The 3 must be clearly indicated.

vertex in approximately correct place for $t = \frac{2}{3}$ and v > 4

[4 marks]

continued...

(M1)

A1

A1

A1

Question 7 continued.

(c) recognising to integrate between 0 and 2, or 2 and 3 OR $\int_{0}^{3} |4+4t-3t^{2}| dt$ (M1)

$$\int_{0}^{2} (4+4t-3t^{2}) dt$$

= 8
$$\int_{2}^{3} (4+4t-3t^{2}) dt$$

valid approach to sum the two areas (seen anywhere)

$$\int_{0}^{2} v \, dt - \int_{2}^{3} v \, dt \quad OR \quad \int_{0}^{2} v \, dt + \left| \int_{2}^{3} v \, dt \right|$$

total distance travelled =13 (m)

A1

(M1)

[5 marks] Total [16 marks]

(a)
$$f\left(\frac{2}{3}\right) = 4 \text{ OR } a^{\frac{2}{3}} = 4$$
 (M1)

– 19 –

$$a = 4^{\frac{3}{2}}$$
 OR $a = (2^2)^{\frac{3}{2}}$ OR $a^2 = 64$ OR $\sqrt[3]{a} = 2$ A1
 $a = 8$ AG

[2 marks]

A1

(b)
$$f^{-1}(x) = \log_8 x$$

Note: Accept
$$f^{-1}(x) = \log_a x$$
.
Accept any equivalent expression for f^{-1} e.g. $f^{-1}(x) = \frac{\ln x}{\ln 8}$.

[1 mark]

(A1)

(A1)

5

(c) correct substitution

$$\log_8 \sqrt{32}$$
 OR $8^x = 32^{\frac{1}{2}}$

correct working involving log/index law

$$\frac{1}{2}\log_8 32 \text{ OR } \frac{5}{2}\log_8 2 \text{ OR } \log_8 2 = \frac{1}{3}\text{ OR } \log_2 2^{\frac{5}{2}} \text{ OR } \log_2 8 = 3\text{ OR } \frac{\ln 2^{\frac{7}{2}}}{\ln 2^3}\text{ OR } 2^{3x} = 2^{\frac{5}{2}}$$

$$f^{-1}(\sqrt{32}) = \frac{5}{6}$$
A1

[3 marks] continued...

Question 8 continued.

(d)

(i) **METHOD 1** equating a pair of differences (M1) $u_2 - u_1 = u_4 - u_3 (= u_3 - u_2)$ $\log_8 p - \log_8 27 = \log_8 125 - \log_8 q$ $\log_8 125 - \log_8 q = \log_8 q - \log_8 p$

$$\log_8\left(\frac{p}{27}\right) = \log_8\left(\frac{125}{q}\right), \ \log_8\left(\frac{125}{q}\right) = \log_8\left(\frac{q}{p}\right)$$

$$p \quad 125 \quad + 125 \quad q \quad A1A1$$

$$\frac{p}{27} = \frac{125}{q}$$
 and $\frac{125}{q} = \frac{q}{p}$ A1

27, p, q and 125 are in geometric sequence

Note: If candidate assumes the sequence is geometric, award no marks for part (i). If $r = \frac{5}{3}$ has been found, this will be awarded marks in part (ii).

METHOD 2

expressing a pair of consecutive terms, in terms of d	(M1)
$p = 8^{d} \times 27$ and $q = 8^{2d} \times 27$ OR $q = 8^{2d} \times 27$ and $125 = 8^{3d} \times 27$	

two correct pairs of consecutive terms, in terms of d A1

$$\frac{8^{d} \times 27}{27} = \frac{8^{2d} \times 27}{8^{d} \times 27} = \frac{8^{3d} \times 27}{8^{2d} \times 27}$$
 (must include 3 ratios) **A1**

all simplify to 8^d A1

27, p, q and 125 are in geometric sequence

continued...

AG

AG

Question 8 continued.

(ii) METHOD 1 (geometric, finding *r*)

$$u_4 = u_1 r^3 \text{ OR } 125 = 27(r)^3$$
 (M1)

$$r = \frac{5}{3}$$
 (seen anywhere) A1

$$p = 27r \text{ OR } \frac{125}{q} = \frac{5}{3}$$
 (M1)

$$p = 45, q = 75$$
 A1A1

METHOD 2 (arithmetic)

$$u_4 = u_1 + 3d \quad \text{OR} \quad \log_8 125 = \log_8 27 + 3d \tag{M1}$$

$$d = \log_8\left(\frac{5}{3}\right) \text{ (seen anywhere)}$$
 A1

$$\log_8 p = \log_8 27 + \log_8 \left(\frac{5}{3}\right) \text{ OR } \log_8 q = \log_8 27 + 2\log_8 \left(\frac{5}{3}\right)$$

$$p = 45, q = 75$$
(M1)
A1A1

METHOD 3 (geometric using proportion)

recognizing proportion $pq = 125 \times 27$ OR $q^2 = 125p$ OR $p^2 = 27q$	(M1)
two correct proportion equations	A1
attempt to eliminate either p or q	(M1)

 $q^2 = 125 \times \frac{125 \times 27}{q} \text{ OR } p^2 = 27 \times \frac{125 \times 27}{p}$

[9 marks] Total [15 marks]

Special note: In this question if candidates use the word 'gradient' in their reasoning. e.g. gradient is positive, it must be clear whether this is the gradient of f or the gradient of f' to earn the **R** mark.

9.	(a)	f increases when $p < x < 0$	A1
		f increases when $f'(x) > 0$ OR f' is above the <i>x</i> -axis	R1

Note: Do not award AOR1.

(b) x = 0

A1

[2 marks]

[1 mark]

A1

R1

A2

R1

(c) (i) f is minimum when x = pbecause f'(p) = 0, f'(x) < 0 when x < p and f'(x) > 0 when x > p(may be seen in a sign diagram clearly labelled as f') OR because f' changes from negative to positive at x = pOR f'(p) = 0 and slope of f' is positive at x = p

Note: Do not award A0 R1

(ii) f has points of inflexion when x = q, x = r and x = t

OR f''(q) = 0, f''(r) = 0 and f''(t) = 0 and f' changes from increasing to decreasing or vice versa at each of these *x*-values (may be seen in a sign diagram clearly labelled as f'' and f')

Note: Award A0 if any incorrect answers are given. Do not award A0R1.

f' has turning points at x = q, x = r and x = t

[5 marks] continued...

Question 9 continued.

(d) recognizing area from p to t (seen anywhere) M1

$$\int_{p}^{t} |f'(x)| dx$$

recognizing to negate integral for area below x-axis

$$\int_{p}^{0} f'(x) dx - \int_{0}^{t} f'(x) dx \quad OR \quad \int_{p}^{0} f'(x) dx + \int_{t}^{0} f'(x) dx$$

$$\int_{m}^{\infty} f'(x) dx = f(n) - f(m)$$
 (for any integral) (M1)

$$f(0) - f(p) - [f(t) - f(0)] \quad \text{OR} \quad f(0) - f(p) + f(0) - f(t)$$
 (A1)

$$2f(0) - [f(t) + f(p)] = 20, 2f(0) - 4 = 20$$
 (A1)

f(0) = 12

A1

(M1)

[6 marks] Total [14 marks]



Markscheme

May 2021

Mathematics: analysis and approaches

Standard level

Paper 1

20 pages



© International Baccalaureate Organization 2021

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2021

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2021

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award **M0** followed by **A1**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **AOA1A1**.
- Where the markscheme specifies A3, M2 etc., do not split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this working is incorrect and/or suggests a misunderstanding of the question. This will encourage a uniform approach to marking, with less examiner discretion. Although some candidates may be advantaged for that specific question item, it is likely that these candidates will lose marks elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used in
 a subsequent part. For example, when a correct exact value is followed by an incorrect
 decimal approximation in the first part and this approximation is then used in the second part.
 In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.
 Examples:

Correct	Further	Any FT issues?	Action
answer seen	working seen		
8./2	5.65685 (incorrect	No. Last part in question.	Award A1 for the final mark (condone the incorrect further
072	decimal value)		working)
35	0.468111	Yes.	Award A0 for the final mark
$\overline{72}$	(incorrect decimal value)	Value is used in subsequent parts	(and full FI is available in subsequent parts)
	Correct answer seen $8\sqrt{2}$ $\frac{35}{72}$	Correct answer seenFurther working seen $8\sqrt{2}$ 5.65685 (incorrect decimal value) $\frac{35}{72}$ 0.468111 (incorrect decimal value)	Correct answer seenFurther working seenAny FT issues? $8\sqrt{2}$ 5.65685 (incorrect decimal value)No. Last part in question. $\frac{35}{72}$ 0.468111 (incorrect (incorrect decimal value)Yes. Value is used in subsequent parts.

3 Implied marks

Implied marks appear in **brackets e.g.** (M1), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is (*M1*)*A1*, it is possible to award full marks for *their* correct answer, without working being seen. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (e.g. probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (MR). A candidate should be penalized only once for a particular misread. Use the MR stamp to indicate that this has been a misread and do not award the first mark, even if this is an M mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by EITHER ... OR.

7 Alternative forms

Unless the question specifies otherwise, accept equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, **M** marks and intermediate **A** marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "from the use of 3 sf values".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an *A* mark to be awarded, arithmetic should be completed,

and any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as

 $\frac{5}{2}$. An exception to this is simplifying fractions, where lowest form is not required

(although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left

in this form or written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate *A* marks do NOT need to be simplified.

- 6 -

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".



Section A

1.	(a)	minor arc AB has length <i>r</i>	(A1)
		recognition that perimeter of shaded sector is $3r$	(A1)
		3r = 12	
		r = 4	A1
			[3 marks]

(b) **EITHER**

$\theta = 2\pi - A\hat{O}B(=2\pi - 1)$	(M1)

Area of non-shaded region =
$$\frac{1}{2}(2\pi-1)(4^2)$$
 (A1)

OR

area of circle - area of shaded sector	(M1)
$16\pi - \left(\frac{1}{2} \times 1 \times 4^2\right)$	(A1)
THEN	

area = $16\pi - 8(2\pi - 1)$)

[3 marks] Total [6 marks]

A1

(M1)

A1A1

A1

AG

2. attempt to subtract squares of integers

$$(n+1)^2 - n^2$$

EITHER

correct order of subtraction and correct expansion of $(n+1)^2$, seen anywhere A1	1A1
---	-----

-9-

$$= n^{2} + 2n + 1 - n^{2} (= 2n + 1)$$

OR

correct order of subtraction and correct factorization of difference of squares

$$=(n+1-n)(n+1+n)(=2n+1)$$

THEN

 $= n + n + 1 = \mathsf{RHS}$

Note: Do not award final A1 unless all previous working is correct.

which is the sum of n and n+1

Note: If expansion and order of subtraction are correct, award full marks for candidates who find the sum of the integers as 2n+1 and then show that the difference of the squares (subtracted in the correct order) is 2n+1.

[4 marks]

[1 mark]

(a) METHOD 1 correct substitution of $\cos^2 x = 1 - \sin^2 x$ A1 $2(1 - \sin^2 x) + 5\sin x = 4$ $2\sin^2 x - 5\sin x + 2 = 0$ AG

METHOD 2

3.

correct substitution using double-angle identities	A1
$\left(2\cos^2 x - 1\right) + 5\sin x = 3$	
$1 - 2\sin^2 x + 5\sin x = 3$	
$2\sin^2 x - 5\sin x + 2 = 0$	AG

continued...

Satprep.

Question 3 continued

(b) **EITHER**

attempting to factorise	М1
$(2\sin x - 1)(\sin x - 2)$	A1

attempting to use the quadratic formula M1 $\sin x = \frac{5 \pm \sqrt{5^2 - 4 \times 2 \times 2}}{4} \left(= \frac{5 \pm 3}{4}\right)$ A1 THEN $\sin x = \frac{1}{2}$ (A1) $x = \frac{\pi}{6}, \frac{5\pi}{6}$ [5 marks] Total [6 marks]

4. EITHER

attempt to use the binomial expansion of
$$(x+k)^7$$
 (M1)
 ${}^7C_0x^7k^0 + {}^7C_1x^6k^1 + {}^7C_2x^5k^2 + ...$ (or ${}^7C_0k^7x^0 + {}^7C_1k^5x^1 + {}^7C_2k^5x^2 + ...$)
identifying the correct term ${}^7C_2x^5k^2$ (or ${}^7C_5k^2x^5$) (A1)

– 12 –

OR

attempt to use the general term ${}^7C_r x^r k^{7-r}$ (or ${}^7C_r k^r x^{7-r}$)	(M1)
r = 2 (or $r = 5$)	(A1)

THEN

${}^{7}C_{2} = 21$ (or ${}^{7}C_{5} = 21$) (seen anywhere)	(A1)
$21x^5k^2 = 63x^5 \ (21k^2 = 63, \ k^2 = 3)$	A1
$k = \pm \sqrt{3}$	A1

Note: If working shown, award *M1A1A1A1A0* for $k = \sqrt{3}$.

[5 marks]

5.

(a)
$$\ln(x^2 - 16) = 0$$
 (M1)
 $e^0 = x^2 - 16(=1)$

$$x^2 = 17 \text{ OR } x = \pm \sqrt{17}$$
 (A1)

$$a = \sqrt{17}$$

М1

(b) attempt to differentiate (must include
$$2x$$
 and/or $\frac{1}{x^2 - 16}$) (M1)

$$f'(x) = \frac{2x}{x^2 - 16}$$
 A1

setting their derivative $=\frac{1}{3}$

$$\frac{2x}{x^2 - 16} = \frac{1}{3}$$

$$x^2 - 16 = 6x \text{ OR } x^2 - 6x - 16 = 0 \text{ (or equivalent)}$$
valid attempt to solve their quadratic
$$(M1)$$

$$x = 8$$
A1

Note: Award *A0* if the candidate's final answer includes additional solutions (such as x = -2, 8).

[6 marks] Total [9 marks]

(M1)

(M1)

(M1)

6. METHOD 1

attempt to use the cosine rule to find the value of x

- 14 -

$$100 = x^{2} + 4x^{2} - 2(x)(2x)\left(\frac{3}{4}\right)$$
 A1

$$2x^2 = 100$$

 $x^2 = 50 \text{ OR } x = \sqrt{50} (= 5\sqrt{2})$ A1

attempt to find $\sin \hat{C}$ (seen anywhere)

$$\sin^2 \hat{C} + \left(\frac{3}{4}\right)^2 = 1$$
 OR $x^2 + 3^2 = 4^2$ or right triangle with side 3 and hypotenuse 4
 $\sin \hat{C} = \frac{\sqrt{7}}{4}$ (A1)

Note: The marks for finding $\sin \hat{C}$ may be awarded independently of the first three marks for finding *x*.

correct substitution into the area formula using their value of x (or x^2) and their value of $\sin \hat{C}$

$$A = \frac{1}{2} \times 5\sqrt{2} \times 10\sqrt{2} \times \frac{\sqrt{7}}{4} \text{ or } A = \frac{1}{2} \times \sqrt{50} \times 2\sqrt{50} \times \frac{\sqrt{7}}{4}$$

$$A = \frac{25\sqrt{7}}{2}$$
A1

continued...

Question 6 continued

METHOD 2

attempt to find the height, h, of the triangle in terms of x

$$h^{2} + \left(\frac{3}{4}x\right)^{2} = x^{2} \text{ OR } h^{2} + \left(\frac{5}{4}x\right)^{2} = 10^{2} \text{ OR } h = \frac{\sqrt{7}}{4}x$$
 A1

equating their expressions for either h^2 or h

$$x^{2} - \left(\frac{3}{4}x\right)^{2} = 10^{2} - \left(\frac{5}{4}x\right)^{2}$$
 OR $\sqrt{100 - \frac{25}{16}x^{2}} = \frac{\sqrt{7}}{4}x$ (or equivalent) **A1**

$$x^2 = 50 \text{ OR } x = \sqrt{50} (= 5\sqrt{2})$$
 A1

correct substitution into the area formula using their value of x (or x^2) (M1)

$$A = \frac{1}{2} \times 2\sqrt{50} \times \frac{\sqrt{7}}{4} \sqrt{50} \text{ OR } A = \frac{1}{2} \left(2 \times 5\sqrt{2} \right) \left(\frac{\sqrt{7}}{4} 5\sqrt{2} \right)$$

$$A = \frac{1}{2}$$

A1

(M1)

(M1)

Total [7 marks]

		Section B	
7.	(a)	evidence of median position	(M1)
		40 students	
		median $= 14$ (hours)	A1
			[2 marks]
	(h)	recognizing there are 8 students in the top 10%	<i>(M</i> 1)
	(0)	72 students spent less than k hours	(<i>M</i> 1)
		k = 18 (hours)	A1
			[3 marks]
			[0
	(c)	15 hours is 60 students OR $p = 60 - 4$	(M1)
		<i>p</i> = 56	A1
		21 hours is 76 students OR $q = 80 - 76$ OR $q = 80 - 4 - 56 - 16$	(A1)
		q = 4	A1
			[4 marks]
	(d)	20 of the 80 students $OR^{\frac{1}{2}}$ spend more than 15 hours doing homework	(A1)
	(9)	4 A A A A A A A A A A A A A A A A A A A	(717)
		$\frac{20}{20} = \frac{x}{200} \text{ OR } \frac{1}{4} \times 320 \text{ OR } 4 \times 20$	(A1)
		80 (students)	A1
			[3 marks]
	(e)	(i) only year 12 students surveyed OR amount of homework might be of	lifferent for
	(-)	different year levels	R1
			_
		(ii) stratified sampling OR survey students in all years	R1
			[2 marks]

8. (a) $6+6\cos x=0$ (or setting their f'(x)=0) (M1) $\cos x=-1$ (or $\sin x=0$)

$$x = \pi, x = 3\pi$$

A1A1

(b) attempt to integrate
$$\int_{\pi}^{3\pi} (6 + 6\cos x) dx$$
 (M1)

$$= \left[6x + 6\sin x \right]_{\pi}^{3\pi}$$
 A1A1

substitute their limits into their integrated expression and subtract	(M1)
$= (18\pi + 6\sin 3\pi) - (6\pi + 6\sin \pi)$	
$=(6(3\pi)+0)-(6\pi+0)(=18\pi-6\pi)$	A1
area = 12π	AG
	[5 marks]

continued...

Question 8 continued

(c)	attempt to substitute into formula for surface area (including base)	(M1)
	$\pi(2^2) + \pi(2)(l) = 12\pi$	(A1)
	$4\pi + 2\pi l = 12\pi$	
	$2\pi l = 8\pi$	
	l = 4	A1
		[3 marks]

(d) valid attempt to find the height of the cone (M1) e.g. $2^2 + h^2 = (\text{their } l)^2$ $h = \sqrt{12} (= 2\sqrt{3})$ (A1) attempt to use $V = \frac{1}{3}\pi r^2 h$ with their values substituted M1 $\left(\frac{1}{3}\pi (2^2)(\sqrt{12})\right)$ $\text{volume} = \frac{4\pi\sqrt{12}}{3} \left(=\frac{8\pi\sqrt{3}}{3}=\frac{8\pi}{\sqrt{3}}\right)$ A1 [4 marks] Total [15 marks] 9. (a) setting s(t) = 0

$$8t - t^2 = 0$$
$$t(8-t) = 0$$

p = 8 (accept t = 8, (8,0))

Note: Award A0 if the candidate's final answer includes additional solutions (such as p = 0, 8).

[2 marks]

(M1)

A1

(M1)	recognition that when particle changes direction $v = 0$ OR local maximum on graph of s OR vertex of parabola	(i)	(b)
A1	q = 4 (accept $t = 4$)		
(M1)	substituting their value of q into $s(t)$ OR integrating $v(t)$ from $t = 0$ to $t = 4$	(ii)	
A1	displacement = 16 (m)		
narks]	[4 n		

(c)	s(10) = -20 OR distance $= s(t) $ OR integrating $v(t)$ from $t = 0$ to $t = 10$	(M1)
	distance=20 (m)	A1

[2 marks]

(d) 16 forward + 36 backward OR 16+16+20 OR
$$\int_{0}^{10} |v(t)| dt$$
 (M1)
 $d=52$ (m) A1

A1

[2 marks]

continued...

Question 9 continued

(e) METHOD 1

graphical method with triangles on v(t) graph

$$49 + \left(\frac{x(2x)}{2}\right) \tag{A1}$$

$$49 + x^2 = 52, \ x = \sqrt{3}$$
 (A1)

$$k = 7 + \sqrt{3}$$

М1

М1

METHOD 2

recognition that distance = $\int |v(t)| dt$

$$\int_{0}^{7} (14-2t) dt + \int_{7}^{k} (2t-14) dt$$

$$\begin{bmatrix} 14t-t^{2} \end{bmatrix}_{0}^{7} + \begin{bmatrix} t^{2}-14t \end{bmatrix}_{7}^{k}$$
(A1)
$$14(7)-7^{2} + ((k^{2}-14k)-(7^{2}-14(7))) = 52$$
(A1)
$$k = 7 + \sqrt{3}$$
[4 marks]
Total [14 marks]



Markscheme

May 2021

Mathematics: analysis and approaches

Standard level

Paper 1





© International Baccalaureate Organization 2021

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

© Organisation du Baccalauréat International 2021

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse https://ibo.org/become-an-ib-school/ ib-publishing/licensing/applying-for-a-license/.

© Organización del Bachillerato Internacional, 2021

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/ applying-for-a-license/.

Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- AG Answer given in the question and so no marks are awarded.
- *FT* Follow through. The practice of awarding marks, despite candidate errors in previous parts, for their correct methods/answers using incorrect results.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award **M0** followed by **A1**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **AOA1A1**.
- Where the markscheme specifies A3, M2 etc., do not split the marks, unless there is a note.
- The response to a "show that" question does not need to restate the *AG* line, unless a **Note** makes this explicit in the markscheme.
- Once a correct answer to a question or part question is seen, ignore further working even if this working is incorrect and/or suggests a misunderstanding of the question. This will encourage a uniform approach to marking, with less examiner discretion. Although some candidates may be advantaged for that specific question item, it is likely that these candidates will lose marks elsewhere too.
- An exception to the previous rule is when an incorrect answer from further working is used in
 a subsequent part. For example, when a correct exact value is followed by an incorrect
 decimal approximation in the first part and this approximation is then used in the second part.
 In this situation, award *FT* marks as appropriate but do not award the final *A1* in the first part.
 Examples:

	Correct	Further	Any FT issues?	Action	
	answer seen	working seen		Action	
1.	<u>د ا</u> م	5.65685 (incorrect	No. Last part in question.	Award A1 for the final mark (condone the incorrect further	
	072	decimal value)		working)	
2.	35	0.468111	Yes.	Award A0 for the final mark	
	72	(incorrect decimal value)	Value is used in subsequent parts.	(and full FI is available in subsequent parts)	

3 Implied marks

Implied marks appear in **brackets e.g.** (M1), and can only be awarded if **correct** work is seen or implied by subsequent working/answer.

4 Follow through marks (only applied after an error is made)

Follow through (*FT*) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) (e.g. incorrect value from part (a) used in part (d) or incorrect value from part (c)(i) used in part (c)(ii)). Usually, to award *FT* marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if all the marks awarded in a subsequent part are for the answer or are implied, then *FT* marks should be awarded for *their* correct answer, even when working is not present.

For example: following an incorrect answer to part (a) that is used in subsequent parts, where the markscheme for the subsequent part is (*M1*)*A1*, it is possible to award full marks for *their* correct answer, without working being seen. For longer questions where all but the answer marks are implied this rule applies but may be overwritten by a **Note** in the Markscheme.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks, by reflecting on what each mark is for and how that maps to the simplified version.
- If the error leads to an inappropriate value (e.g. probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- If the candidate's answer to the initial question clearly contradicts information given in the question, it is not appropriate to award any *FT* marks in the subsequent parts. This includes when candidates fail to complete a "show that" question correctly, and then in subsequent parts use their incorrect answer rather than the given value.
- Exceptions to these *FT* rules will be explicitly noted on the markscheme.
- If a candidate makes an error in one part but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the command term was "Hence".

5 Mis-read

If a candidate incorrectly copies values or information from the question, this is a mis-read (MR). A candidate should be penalized only once for a particular misread. Use the MR stamp to indicate that this has been a misread and do not award the first mark, even if this is an M mark, but award all others as appropriate.

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does **not** constitute a misread, it is an error.
- If a candidate uses a correct answer, to a "show that" question, to a higher degree of accuracy than given in the question, this is NOT a misread and full marks may be scored in the subsequent part.
- *MR* can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If the command term is 'Hence' and not 'Hence or otherwise' then alternative methods are not permitted unless covered by a note in the mark scheme.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for parts of questions are indicated by EITHER ... OR.

7 Alternative forms

Unless the question specifies otherwise, accept equivalent forms.

- As this is an international examination, accept all alternative forms of **notation** for example 1.9 and 1,9 or 1000 and 1,000 and 1.000.
- Do not accept final answers written using calculator notation. However, **M** marks and intermediate **A** marks can be scored, when presented using calculator notation, provided the evidence clearly reflects the demand of the mark.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, some **equivalent** answers will generally appear in brackets. Not all equivalent notations/answers/methods will be presented in the markscheme and examiners are asked to apply appropriate discretion to judge if the candidate work is equivalent.

8 Format and accuracy of answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. If the level of accuracy is not stated in the question, the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Where values are used in subsequent parts, the markscheme will generally use the exact value, however candidates may also use the correct answer to 3 sf in subsequent parts. The markscheme will often explicitly include the subsequent values that come "from the use of 3 sf values".

Simplification of final answers: Candidates are advised to give final answers using good mathematical form. In general, for an **A** mark to be awarded, arithmetic should be completed,

and any values that lead to integers should be simplified; for example, $\sqrt{\frac{25}{4}}$ should be written as

 $\frac{5}{2}$. An exception to this is simplifying fractions, where lowest form is not required

(although the numerator and the denominator must be integers); for example, $\frac{10}{4}$ may be left

in this form or written as $\frac{5}{2}$. However, $\frac{10}{5}$ should be written as 2, as it simplifies to an integer.

Algebraic expressions should be simplified by completing any operations such as addition and multiplication, e.g. $4e^{2x} \times e^{3x}$ should be simplified to $4e^{5x}$, and $4e^{2x} \times e^{3x} - e^{4x} \times e^{x}$ should be simplified to $3e^{5x}$. Unless specified in the question, expressions do not need to be factorized, nor do factorized expressions need to be expanded, so x(x+1) and $x^2 + x$ are both acceptable.

Please note: intermediate *A* marks do NOT need to be simplified.

- 6 -

9 Calculators

No calculator is allowed. The use of any calculator on this paper is malpractice and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice.

10. Presentation of candidate work

Crossed out work: If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work unless an explicit note from the candidate indicates that they would like the work to be marked.

More than one solution: Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise. If the layout of the responses makes it difficult to judge, examiners should apply appropriate discretion to judge which is "first".



A1

1. (a) (i)
$$f(2) = 6$$

(ii) $(f \circ f)(2) = -2$



(b)



[3 marks] Total [5 marks]

A1

2. (a) 3×10⁴ OR 30000 (km) (accept 3•10⁴)

[1 mark]

(b)
$$\frac{4}{3}\pi (3 \times 10^4)^3 \text{ OR } \frac{4}{3}\pi (30000)^3$$
 (A1)
= $\frac{4}{3}\pi \times 27 \times 10^{12} (=\pi (36 \times 10^{12})) \text{ OR } = \frac{4}{3}\pi \times 2700000000000$ (A1)
= $\pi (3.6 \times 10^{13}) \text{ (km}^3) \text{ OR } a = 3.6, k = 13$ A1

[3 marks] Total [4 marks]

у 8¹

6

3. METHOD 1 (finding u_1 first, from S₈)

$4(u_1+8)=8$	(A1)
$u_1 = -6$	A1
$u_1 + 7d = 8$ OR $4(2u_1 + 7d) = 8$ (may be seen with their value of u_1)	(A1)
attempt to substitute their u_1	
d = 2	A1

METHOD 2 (solving simultaneously)

$u_1 + 7d = 8$	(A1)
$4(u_1+8) = 8 \text{ OR } 4(2u_1+7d) = 8 \text{ OR } u_1 = -3d$	(A1)
attempt to solve linear or simultaneous equations	(M1)
$u_1 = -6, d = 2$	A1A1

[5 marks]

4.	(a)	attempt to use definition of outlier		
		$1.5 \times 20 + Q_3$	(M1)	
		$1.5 \times 20 + U \ge 75 \ (\Rightarrow U \ge 45, \text{ accept } U > 45) \ \text{OR} \ 1.5 \times 20 + Q_3 = 75$	A1	
		minimum value of $U = 45$	A1	
			[3	marks]
	(b)	attempt to use interquartile range $U - L = 20$ (may be seen in part (a)) OR $L \ge 25$ (accept $L \ge 25$)	(M1)	
		minimum value of $L=25$	A1	
			[2	markel

[2 marks] Total [5 marks]
5. (a)
$$f'(x) = -2(x-h)$$
 A1

[1 mark]

(b)
$$g'(x) = e^{x-2} \text{ OR } g'(3) = e^{3-2}$$
 (may be seen anywhere) A1
Note: The derivative of g must be explicitly seen, either in terms of x or 3.
recognizing $f'(3) = g'(3)$ (M1)
 $-2(3-h) = e^{3-2} (=e)$
 $-6+2h = e \text{ OR } 3-h = -\frac{e}{2}$ A1
Note: The final A1 is dependent on one of the previous marks being awarded.
 $h = \frac{e+6}{2}$ AG

(c) f(3) = g(3) (M1) $-(3-h)^2 + 2k = e^{3-2} + k$ correct equation in k EITHER $-\left(3 - \frac{e+6}{2}\right)^2 + 2k = e^{3-2} + k$ A1

$$k = e + \left(\frac{6 - e - 6}{2}\right)^2 \left(= e + \left(\frac{-e}{2}\right)^2\right)$$

OR

$$k = e + \left(3 - \frac{e+6}{2}\right)^{2}$$

$$k = e + 9 - 3e - 18 + \frac{e^{2} + 12e + 36}{4}$$
A1

THEN

$$k = e + \frac{e^2}{4}$$
 AG

[3 marks] Total [7 marks]

– 10 –

6.	(a)
-	(-)

tempt to use double angle formula for $\sin 2x$ or $\cos 2x$	М1
$HS = 2\sin x \cos x + \cos 2x - 1 \text{ OR}$	
$n 2x + 1 - 2 \sin^2 x - 1$ OR	
$\sin x \cos x + 1 - 2\sin^2 x - 1$	
$2\sin x \cos x - 2\sin^2 x$	A1
$n 2x + \cos 2x - 1 = 2\sin x(\cos x - \sin x) = RHS$	AG
ETHOD 2 (RHS to LHS)	
$HS = 2\sin x \cos x - 2\sin^2 x$	
attempt to use double angle formula for $\sin 2x$ or $\cos 2x$	M1
$=\sin 2x + 1 - 2\sin^2 x - 1$	A1
$=\sin 2x + \cos 2x - 1 = LHS$	AG
tempt to factorise	M1
$\cos x - \sin x(2\sin x + 1) = 0$	A1
cognition of $\cos x = \sin x \Rightarrow \frac{\sin x}{\cos x} = \tan x = 1$ OR $\sin x = -\frac{1}{2}$	(M1)
ne correct reference angle seen anywhere, accept degrees	(A1)
$\frac{\pi}{4}$ OR $\frac{\pi}{6}$ (accept $-\frac{\pi}{6}, \frac{7\pi}{6}$)	
ote: This (M1)(A1) is independent of the previous M1A1.	
$=\frac{7\pi}{11\pi},\frac{11\pi}{2},\frac{\pi}{2},\frac{5\pi}{2}$	A2
6 6 4 4	
ote: Award A1 for any two correct (radian) answers.	(radian) answara
Awaru A IAU II additional values given with the four correct	(iauiaii) answers.
Award A1A0 for four correct answers given in degrees.	

(a)	METHOD 1 (discriminant)	
	$mx^2 - 2mx = mx - 9$	(M1)
	$mx^2 - 3mx + 9 = 0$	
	recognizing $\Delta = 0$ (seen anywhere)	M1
	$\Delta = (-3m)^2 - 4(m)(9)$ (do not accept only in quadratic formula for x)	A1
	valid approach to solve quadratic for m	(M1)
	$9m(m-4) = 0 \text{ OR } m = \frac{36 \pm \sqrt{36^2 - 4 \times 9 \times 0}}{2 \times 9}$	
	both solutions $m = 0, 4$	A1
	$m \neq 0$ with a valid reason	R1
	the two graphs would not intersect OR $0 \neq -9$	
	m = 4	AG
	METHOD 2 (equating slopes)	
	$mx^2 - 2mx = mx - 9$ (seen anywhere)	(M1)
	f'(x) = 2mx - 2m	A1
	equating slopes, $f'(x) = m$ (seen anywhere)	М1
	2mx - 2m = m	
	3	
	$x = \frac{1}{2}$	AI
	substituting their x value	(M1)
	$\left(\frac{3}{2}\right)^2 m - 2m \times \frac{3}{2} = m \times \frac{3}{2} - 9$	
	$\frac{9}{4}m - \frac{12}{4}m = \frac{6}{4}m - 9$	A1
	$\frac{-9m}{-9} = -9$	
	4	
	m=4	AG

continued...

7.

Question 7(a) continued

	METHOD 3 (using $\frac{-b}{2a}$)		
	$mx^2 - 2mx = mx - 9$ $mx^2 - 3mx + 9 = 0$	(M1)	
	attempt to find x -coord of vertex using $\frac{-b}{2a}$	(M1)	
	$\frac{-(-3m)}{2m}$	A1	
	$x = \frac{3}{2}$	A1	
	substituting their x value	(M1)	
	$\left(\frac{3}{2}\right)^2 m - 3m \times \frac{3}{2} + 9 = 0$. ,	
	$\frac{9}{4}m - \frac{9}{2}m + 9 = 0$	A1	
	-9m = -36 $m = 4$	AG	
			[6 marks]
(b)	4x(x-2)	(A1)	
	p=0 and $q=2$ OR $p=2$ and $q=0$	A1	
			[2 marks]
(c)	attempt to use valid approach	(M1)	
	$\frac{0+2}{2}, \frac{-(-8)}{2\times4}, f(1), 8x-8=0 \text{ OR } 4(x^2-2x+1-1)(=4(x-1)^2-4)$		
	h=1, k=-4	A1A1	10
			[3 marks]
(d)	EITHER		
	recognition $x = h$ to 2 (may be seen on sketch)	(M1)	
	OR		
	recognition that $f(x) < 0$ and $f'(x) > 0$	(M1)	
	THEN		
	1 < <i>x</i> < 2	A1A1	
	Note: Award A1 for two correct values, A1 for correct inequality signs.		

[3 marks] Total [14 marks]

(M1)

– 14 –

8. (a) attempt to use quotient or product rule

 $P\!\left(e^{\frac{1}{4}},\frac{1}{4e}\right)$

$$\frac{dy}{dx} = \frac{x^4 \left(\frac{1}{x}\right) - (\ln x)(4x^3)}{\left(x^4\right)^2} \quad \text{OR} \quad (\ln x)(-4x^{-5}) + (x^{-4})\left(\frac{1}{x}\right)$$

correct working

$$= \frac{x^{3}(1-4\ln x)}{x^{8}} \quad \text{OR cancelling } x^{3} \quad \text{OR} \quad \frac{-4\ln x}{x^{5}} + \frac{1}{x^{5}}$$

$$= \frac{1-4\ln x}{x^{5}} \quad \text{AG}$$

(b)
$$f'(x) = \frac{dy}{dx} = 0$$
 (M1)

$$\frac{1-4\ln x}{x^5} = 0$$

$$\ln x = \frac{1}{4}$$
 (A1)

$$x = e^{\frac{1}{4}}$$
 A1
substitution of their x to find y (M1)

$$y = \frac{\ln e^{\frac{1}{4}}}{\left(e^{\frac{1}{4}}\right)^4}$$

$$= \frac{1}{4e} \left(= \frac{1}{4} e^{-1}\right)$$
 A1

[5 marks]

continued...

Question 8 continued





9.	(a)	recognising probabilities sum to 1	(M1)	
		$p + p + p + \frac{1}{2}p = 1$		
		$p = \frac{2}{2}$	A1	
		' 7		[2 marks]
	(h)	valid attempt to find $F(X)$	(M1)	
	(0)	$\frac{1}{2} \left(\frac{1}{2} \right) \left(1$	(1117)	
		$1 \times p + 2 \times p + 3 \times p + 4 \times \frac{2}{2} p (= 8p)$		
		$E(X) = \frac{16}{7}$	A1	
				[2 marks]
	(c)	(i) $0 \le r \le 1$	A1	
		(ii) attempt to find a value of q	(M1)	
		$0 \le 1 - 3q \le 1$ OR $r = 0 \Longrightarrow q = \frac{1}{3}$ OR $r = 1 \Longrightarrow q = 0$		
		$0 \le q \le \frac{1}{2}$	A1	
		1 3		[3 marks]
				[0
	(d)	$E(Y) = 1 \times q + 2 \times q + 3 \times q + 4 \times r (= 2 + 2r \text{ OR } 4 - 6q)$	(A1)	
		one correct boundary value	A1	
		$1 \times \frac{1}{3} + 2 \times \frac{1}{3} + 3 \times \frac{1}{3} + 4 \times 0 (= 2) \text{ OR}$		
		$1 \times 0 + 2 \times 0 + 3 \times 0 + 4 \times 1 (= 4)$ OR		
		2+2(0)(=2) OR		
		2+2(1)(=4) OR		
		$4-6(0)(=4) \text{ OR } 4-6\left(\frac{1}{3}\right)(=2)$		
		$2 \le \mathrm{E}(Y) \le 4$	A1	
				[3 marks]

continued...

М1

Question 9 continued

METHOD 1 (e)

evidence of choosing at least four correct outcomes from 1&2, 1&3, 1&4, 2&3, 2&4, 3&4 (M1)

$$\frac{6}{7}q + \frac{6}{7}r \text{ OR } 3pq + 3pr \text{ OR } pq + pq + p(1-3q) + pq + p(1-3q) + p(1-3q) \text{ (A1)}$$

solving for either q or r

$$\frac{6}{7}(q+1-3q) = \frac{1}{2} \text{ OR } \frac{6}{7}\left(\frac{1-r}{3}+r\right) = \frac{1}{2} \text{ OR } 3pq+3p(1-3q) = \frac{1}{2}$$

OR $3p\left(\frac{1-r}{3}\right) + 3pr = \frac{1}{2}$

EITHER two correct values

$$q = \frac{5}{24}$$
 and $r = \frac{3}{8}$ **A1A1**

OR one correct value

$q = \frac{5}{24}$ OR $r = \frac{3}{8}$	A1
substituting their value for q or r	A1
$4 - 6\left(\frac{5}{24}\right) \text{OR } 2 + 2\left(\frac{3}{8}\right)$	
THEN	

$$\mathrm{E}(Y) = \frac{11}{4}$$

A1

[6 marks]

continued...

М1

Question 9 continued

METHOD 2 (solving for E(Y))evidence of choosing at least four correct outcomes from1&2, 1&3, 1&4, 2&3, 2&4, 3&4(M1)

$$\frac{6}{7}q + \frac{6}{7}r \text{ OR } 3pq + 3pr \text{ OR } pq + pq + p(1-3q) + pq + p(1-3q) + p(1-3q) \text{ (A1)}$$

rearranging to make q the subject

$$q = \frac{4 - E(Y)}{6}$$

3pq + 3p(1-3q) = $\frac{1}{2}$ M1

$$\frac{6}{7} \times \left(\frac{4 - E(Y)}{6}\right) + \frac{6}{7} \left(1 - 3\left(\frac{4 - E(Y)}{6}\right)\right) = \frac{1}{2}$$

$$\frac{2(E(Y) - 1)}{7} = \frac{1}{2}$$

$$E(Y) = \frac{11}{4}$$

$$A1$$
[6 marks]
Total [16 marks]



Markscheme

Specimen paper

Mathematics: analysis and approaches

Standard level

Paper 1

11 pages



Instructions to Examiners

Abbreviations

- *M* Marks awarded for attempting to use a correct **Method**.
- **A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- *R* Marks awarded for clear **Reasoning**.
- **AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg M1, A2.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award *M0* followed by *A1*, as *A* mark(s) depend on the preceding *M* mark(s), if any.
- Where *M* and *A* marks are noted on the same line, *e.g. M1A1*, this usually means *M1* for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and *A1* for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies M2, A2, etc., do not split the marks, unless there is a note.
- Once a correct answer to a question or part-question is seen, ignore further correct working. However, if further working indicates a lack of mathematical understanding do not award the final *A1*. An exception to this may be in numerical answers, where a correct exact value is followed by an incorrect decimal. However, if the incorrect decimal is carried through to a subsequent part, and correct working shown, award *FT* marks as appropriate but do not award the final *A1* in that part.

	Correct answer seen	Further working seen	Action
1.	8\sqrt{2}	5.65685 (incorrect decimal value)	Award the final A1 (ignore the further working)
2.	$\frac{1}{4}\sin 4x$	sin x	Do not award the final A1
3.	$\log a - \log b$	$\log(a-b)$	Do not award the final A1

Examples

3 Implied marks

Implied marks appear in **brackets e.g. (M1)**, and can only be awarded if **correct** work is seen or if implied in subsequent working.

- Normally the correct work is seen or implied in the next line.
- Marks without brackets can only be awarded for work that is seen.

4 Follow through marks (only applied after an error is made)

Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) or subpart(s). Usually, to award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if the only marks awarded in a subpart are for the answer (i.e. there is no working expected), then **FT** marks should be awarded if appropriate.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer *FT* marks.
- If the error leads to an inappropriate value (*e.g.* probability greater than 1, use of r > 1 for the sum of an infinite GP, $\sin \theta = 1.5$, non integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word "their" in a description, to indicate that candidates may be using an incorrect value.
- Exceptions to this rule will be explicitly noted on the markscheme.
- If a candidate makes an error in one part, but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the question says hence. It is often possible to use a different approach in subsequent parts that does not depend on the answer to previous parts.

5 Mis-read

If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). Apply a **MR** penalty of 1 mark to that question

- If the question becomes much simpler because of the *MR*, then use discretion to award fewer marks.
- If the *MR* leads to an inappropriate value (*e.g.* probability greater than 1, $\sin \theta = 1.5$, non-integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates' own work does not constitute a misread, it is an error.
- The *MR* penalty can only be applied when work is seen. For calculator questions with no working and incorrect answers, examiners should **not** infer that values were read incorrectly.

- 3 -

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, *etc*.
- Alternative solutions for part-questions are indicated by **EITHER** . . . **OR**.

7 Alternative forms

Unless the question specifies otherwise, *accept* equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

8 Accuracy of Answers

If the level of accuracy is specified in the question, a mark will be linked to giving the answer to the required accuracy. There are two types of accuracy errors, and the final answer mark should not be awarded if these errors occur.

- Rounding errors: only applies to final answers not to intermediate steps.
- Level of accuracy: when this is not specified in the question the general rule applies to final answers: unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.

9 Calculators

No calculator is allowed. The use of any calculator on paper 1 is malpractice, and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice. Examples: finding an angle, given a trig ratio of 0.4235.

Section A

1.	(a)	valid approach using Pythagorean identity	(M1)	
		$\sin^2 A + \left(\frac{5}{6}\right)^2 = 1$ (or equivalent)	(A1)	
		$\sin A = \frac{\sqrt{11}}{6}$	A1	
				[3 marks]
	(b)	$\frac{1}{2} \times 8 \times 6 \times \frac{\sqrt{11}}{6}$ (or equivalent)	(A1)	
		area = $4\sqrt{11}$	A1	[2 marks]
			Total	[5 marks]
2.	atten	npt to substitute into $P(A \cup B) = P(A) + P(B) - P(A \cap B)$	(M1)	
Not	:e: Acc 0.6 =	cept use of Venn diagram or other valid method. = $0.5 + 0.4 - P(A \cap B)$	(A1)	
	P(A	$(\cap B) = 0.3$ (seen anywhere)	A1	
	atten	npt to substitute into $P(A B) = \frac{P(A \cap B)}{P(B)}$	(M1)	
	$=\frac{0.3}{0.4}$	$\frac{3}{4}$		
	P(A	$ B) = 0.75 \left(=\frac{3}{4}\right)$	A1	
		satprev.	Total	[5 marks]

SPEC/5/MATAA/SP1/ENG/TZ0/XX/M

3. (a) attempting to expand the LHS (M1) $LHS = (4n^2 - 4n + 1) + (4n^2 + 4n + 1)$ A1 $= 8n^2 + 2(= RHS)$ AG

[2 marks]

(b) METHOD 1

recognition that $2n-1$ and $2n+1$ represent two consecutive odd		
integers (for $n \in \mathbb{Z}$)	R1	
$8n^2 + 2 = 2(4n^2 + 1)$	A1	
valid reason <i>eg</i> divisible by 2 (2 is a factor)	R1	
so the sum of the squares of any two consecutive odd integers is even	AG	
	[3	} marks]

METHOD 2

recognition, <i>eg</i> that $_n$ and $n+2$ represent two consecutive odd integers	
(for $n \in \mathbb{Z}$)	R1
$n^{2} + (n+2)^{2} = 2(n^{2} + 2n + 2)$	A1
valid reason <i>eg</i> divisible by 2 (2 is a factor)	R1
so the sum of the squares of any two consecutive odd integers is even	AG [3 marks]
	Total [5 marks]
attempt to integrate $u = 2x^2 + 1 \Rightarrow \frac{du}{dx} = 4x$	(M1)
$\int \frac{8x}{\sqrt{2x^2 + 1}} \mathrm{d}x = \int \frac{2}{\sqrt{u}} \mathrm{d}u$	(A1)
EITHER	
$=4\sqrt{u}(+C)$	A1

OR

4.

 $=4\sqrt{2x^2+1}(+C)$ A1

THEN

correct substitution into their integrated function (must have C)	(M1)
$5 = 4 + C \Longrightarrow C = 1$	
$f(x) = 4\sqrt{2x^2 + 1} + 1$	A1
	Total [5 marks]

-6-

5.	(a)	attempt to form composition	М1
		correct substitution $g\left(\frac{x+3}{4}\right) = 8\left(\frac{x+3}{4}\right) + 5$	A1
		$(g \circ f)(x) = 2x + 11$	AG
			[2 marks]
	(b)	attempt to substitute 4 (seen anywhere) correct equation $a = 2 \times 4 + 11$ a = 19	(M1) (A1) A1 [3 marks]
			Total [5 marks]
6.	(a)	attempting to use the change of base rule	М1
		$\log_{9}(\cos 2x + 2) = \frac{\log_{3}(\cos 2x + 2)}{\log_{3} 9}$	A1
		$=\frac{1}{2}\log_3(\cos 2x+2)$	A1
		$=\log_3\sqrt{\cos 2x+2}$	AG
			[3 marks]
	(b)	$\log_3(2\sin x) = \log_3\sqrt{\cos 2x + 2}$	
		$2\sin x = \sqrt{\cos 2x + 2}$	M1
		$4\sin^2 x = \cos 2x + 2$ (or equivalent)	A1
		use of $\cos 2x = 1 - 2\sin^2 x$ $6\sin^2 x = 3$	(M1)
		$\sin x = (\pm) \frac{1}{\sqrt{2}}$	A1
		$x = \frac{\pi}{4}$	A1
	No	te: Award A0 if solutions other than $x = \frac{\pi}{4}$ are included.	

[5 marks]

Total [8 marks]

Section B

7.	(a)	evidence of median position	(M1)	
		40 minutes	A1	[2 marks]
	(b)	valid attempt to find interval (25–55) 18 (employees), 142 (employees) 124	(M1) A1 A1	
				[3 marks]
	(c)	recognising that there are 16 employees in the top 10% 144 employees travelled more than k minutes $k = 56$	(M1) (A1) A1	
				[3 marks]
	(d)	<i>b</i> = 70	A1	[1 mark]
	(e)	 (i) recognizing <i>a</i> is first quartile value 40 employees 	(M1)	
		<i>a</i> = 33	A1	
		(ii) $47-33$ IQR = 14	(M1) A1	
				[4 marks]
	(f)	attempt to find $1.5 \times$ their IQR $33-21$	(M1)	
		12	(A1)	[2 marks]
			[Total	15 marks]
8.	(a)	$f'(x) = x^2 + 2x - 15$	(M1)A1	[2 marks]
	(b)	correct reasoning that $f'(x) = 0$ (seen anywhere)	(M1)	
		$x^{2} + 2x - 15 = 0$ valid approach to solve quadratic (x - 3)(x + 5), quadratic formula correct values for x 3x - 5	М1	

correct values for a and b

a = -5 and b = 3

A1 [3 marks]

continued...

- 8 -

Question 8 continued



(a) attempt to use quotient rule (M1) correct substitution into quotient rule $f'(x) = \frac{5kx\left(\frac{1}{5x}\right) - k\ln 5x}{\left(kx\right)^2}$ (or equivalent) A1 $=\frac{k-k\ln 5x}{k^2x^2}, (k\in\mathbb{R}^+)$ A1 $=\frac{1-\ln 5x}{kx^2}$ AG [3 marks] (b) f'(x) = 0М1 $\frac{1-\ln 5x}{kx^2} = 0$ $\ln 5x = 1$ (A1) $x = \frac{e}{5}$ A1 [3 marks] (c) f''(x) = 0 $\frac{2 \ln 5x - 3}{kx^3} = 0$ М1 $\ln 5x = \frac{3}{2}$ A1 $5x = e^{\frac{3}{2}}$ A1 so the point of inflexion occurs at $x = \frac{1}{5}e^{\frac{3}{2}}$ AG [3 marks] continued...

– 10 –

9.

Question 9 continued

attempt to integrate (M1) (d) $u = \ln 5x \Longrightarrow \frac{\mathrm{d}u}{\mathrm{d}x} = \frac{1}{x}$ c ln 5 r 1 .

$$\int \frac{\ln 3x}{kx} dx = \frac{1}{k} \int u \, du \tag{A1}$$

– 11 –

EITHER

$$=\frac{u^2}{2k}$$

so
$$\frac{1}{k} \int_{1}^{\frac{3}{2}} u \, \mathrm{d}u = \left[\frac{u^2}{2k}\right]_{1}^{\frac{3}{2}}$$

OR

$$= \frac{(\ln 5x)^{2}}{2k}$$
so $\int_{\frac{e}{5}}^{\frac{1}{5}e^{\frac{3}{2}}} \frac{\ln 5x}{kx} dx = \left[\frac{(\ln 5x)^{2}}{2k}\right]_{\frac{e}{5}}^{\frac{1}{5}e^{\frac{3}{2}}}$
THEN

THEN

$$=\frac{1}{2k}\left(\frac{9}{4}-1\right)$$

$$=\frac{5}{8k}$$
Setting their expression for area equal to 3
$$\frac{5}{8k} = 3$$

$$k = \frac{5}{24}$$
A1

[7 marks]

Total [16 marks]