* 0000800000001 *

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

* 0 7 5 9 5 7 4 0 8 2 *

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **October/November 2024**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2024

[Turn over

BLANK PAGE

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

1 The polynomial $4x^3 + ax^2 + 5x + b$, where a and b are constants, is denoted by $\mathrm{p}(x)$. It is given that $(2x+1)$ is a factor of $\mathrm{p}(x)$. When $\mathrm{p}(x)$ is divided by $(x-4)$ the remainder is equal to 3 times the remainder when $\mathrm{p}(x)$ is divided by $(x-2)$.

Find the values of a and b. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

2 Find the exact value of $\int_1^3 x^2 \ln 3x \, dx$. Give your answer in the form $a \ln b + c$, where a and c are rational and b is an integer. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

3 The equation of a curve is $\ln(x+y) = 3x^2y$.

Find the gradient of the curve at the point $(1, 0)$. [4]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

4 **(a)** Show that $\sec^4\theta - \tan^4\theta \equiv 1 + 2\tan^2\theta$. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) Hence or otherwise solve the equation $\sec^4 2\alpha - \tan^4 2\alpha = 2\tan^2 2\alpha \sec^2 2\alpha$ for $0° < \alpha < 180°$. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

5 **(a)** By sketching a suitable pair of graphs, show that the equation $2+e^{-0.2x} = \ln(1+x)$ has only one root. [2]

(b) Show by calculation that this root lies between 7 and 9. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN

(c) Use the iterative formula

$$x_{n+1} = \exp\left(2 + \mathrm{e}^{-0.2x_n}\right) - 1$$

to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places.

[exp(x) is an alternative notation for e^x.] [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

6



The diagram shows the curve $y = \sin 2x(1 + \sin 2x)$, for $0 \leqslant x \leqslant \frac{3}{4}\pi$, and its minimum point M. The shaded region bounded by the curve that lies above the x-axis and the x-axis itself is denoted by R.

(a) Given that the x-coordinate of M lies in the interval $\frac{1}{2}\pi < x < \frac{3}{4}\pi$, find the exact coordinates of M. [4]

DO NOT WRITE IN THIS MARGIN

(b) Find the exact area of the region R. [4]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

7 Let $f(x) = \dfrac{5x^2 + 8x + 5}{(1 + 2x)(2 + x^2)}$.

(a) Express $f(x)$ in partial fractions. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) Hence find the coefficient of x^3 in the expansion of f(x). [4]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

8 **(a)** Given that $z = 1 + y\mathrm{i}$ and that y is a real number, express $\frac{1}{z}$ in the form $a + b\mathrm{i}$, where a and b are functions of y. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show that $\left(a - \frac{1}{2}\right)^2 + b^2 = \frac{1}{4}$, where a and b are the functions of y found in part **(a)**. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

.....

.....

.....

.....

.....

(c) On a single Argand diagram, sketch the loci given by the equations $\mathrm{Re}(z) = 1$ and $\left|z - \frac{1}{2}\right| = \frac{1}{2}$, where z is a complex number. [3]

(d) The complex number z is such that $\mathrm{Re}(z) = 1$. Use your answer to part **(b)** to give a geometrical description of the locus of $\frac{1}{z}$. [1]

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

9 The position vector of point A relative to the origin O is $\overrightarrow{OA} = 8\mathbf{i} - 5\mathbf{j} + 6\mathbf{k}$.
The line l passes through A and is parallel to the vector $2\mathbf{i} + \mathbf{j} + 4\mathbf{k}$.

(a) State a vector equation for l. [2]

.....

.....

.....

.....

.....

.....

.....

.....

(b) The position vector of point B relative to the origin O is $\overrightarrow{OB} = -t\mathbf{i} + 4t\mathbf{j} + 3t\mathbf{k}$, where t is a constant.
The line l also passes through B.

Find the value of t. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(c) The line m has vector equation $\mathbf{r} = 5\mathbf{i} - \mathbf{j} + 2\mathbf{k} + \mu(a\mathbf{i} - \mathbf{j} + 3\mathbf{k})$. The acute angle between the directions of l and m is θ, where $\cos\theta = \frac{1}{\sqrt{6}}$.

Find the possible values of a. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

10



A large cylindrical tank is used to store water. The base of the tank is a circle of radius 4 metres. At time t minutes, the depth of the water in the tank is h metres. There is a tap at the bottom of the tank. When the tap is open, water flows out of the tank at a rate proportional to the square root of the volume of water in the tank.

(a) Show that $\frac{\mathrm{d}h}{\mathrm{d}t} = -\lambda\sqrt{h}$, where λ is a positive constant. [4]

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) At time $t = 0$ the tap is opened. It is given that $h = 4$ when $t = 0$ and that $h = 2.25$ when $t = 20$.

Solve the differential equation to obtain an expression for t in terms of h, and hence find the time taken to empty the tank. [6]

SAT PREP SP www.satprep.co.in

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

* 0000800000001 *

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

* 3 4 4 9 8 5 6 2 9 5 *

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **October/November 2024**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2024

[Turn over

1 Expand $(9-3x)^{\frac{1}{2}}$ in ascending powers of x, up to and including the term in x^2, simplifying the coefficients. [4]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

2 **(a)** By sketching a suitable pair of graphs, show that the equation $\cot 2x = \sec x$ has exactly one root in the interval $0 < x < \frac{1}{2}\pi$. [2]

(b) Show that if a sequence of real values given by the iterative formula

$$x_{n+1} = \frac{1}{2}\tan^{-1}(\cos x_n)$$

converges, then it converges to the root in part **(a)**. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

3 The square roots of $6-8\mathrm{i}$ can be expressed in the Cartesian form $x+\mathrm{i}y$, where x and y are real and exact.

By first forming a quartic equation in x or y, find the square roots of $6-8\mathrm{i}$ in exact Cartesian form. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

4 Solve the equation $5^x = 5^{x+2} - 10$. Give your answer correct to 3 decimal places. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

5 **(a)** The complex number u is given by

$$u = \frac{(\cos\frac{1}{7}\pi + \mathrm{i}\sin\frac{1}{7}\pi)^4}{\cos\frac{1}{7}\pi - \mathrm{i}\sin\frac{1}{7}\pi}.$$

Find the exact value of arg u. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The complex numbers u and u^* are plotted on an Argand diagram.

Describe the single geometrical transformation that maps u onto u^* and state the exact value of arg u^*. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN

6



The variables x and y satisfy the equation $ay = b^x$, where a and b are constants. The graph of $\ln y$ against x is a straight line passing through the points (0.50, 2.24) and (3.40, 8.27), as shown in the diagram.

Find the values of a and b. Give each value correct to 1 significant figure. [4]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

7 **(a)** Show that the equation $\tan^3 x + 2\tan 2x - \tan x = 0$ may be expressed as

$$\tan^4 x - 2\tan^2 x - 3 = 0$$

for $\tan x \neq 0$. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) Hence solve the equation $\tan^3 2\theta + 2\tan 4\theta - \tan 2\theta = 0$ for $0 < \theta < \pi$. Give your answers in exact form. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

8 The parametric equations of a curve are

$$x = \tan^2 2t, \quad y = \cos 2t,$$

for $0 < t < \frac{1}{4}\pi$.

(a) Show that $\frac{\mathrm{d}y}{\mathrm{d}x} = -\frac{1}{2}\cos^3 2t$. [4]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) Hence find the equation of the normal to the curve at the point where $t = \frac{1}{8}\pi$. Give your answer in the form $y = mx + c$. [4]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

9 With respect to the origin O, the points A, B and C have position vectors given by

$$\overrightarrow{OA} = \begin{pmatrix} 2 \\ 1 \\ -3 \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 0 \\ 4 \\ 1 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OC} = \begin{pmatrix} -3 \\ -2 \\ 2 \end{pmatrix}.$$

(a) The point D is such that $ABCD$ is a trapezium with $\overrightarrow{DC} = 3\overrightarrow{AB}$.

Find the position vector of D. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The diagonals of the trapezium intersect at the point P.

Find the position vector of P. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Using a scalar product, calculate angle ABC. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10 A balloon in the shape of a sphere has volume V and radius r. Air is pumped into the balloon at a constant rate of 40π starting when time $t = 0$ and $r = 0$. At the same time, air begins to flow out of the balloon at a rate of $0.8\pi r$. The balloon remains a sphere at all times.

(a) Show that r and t satisfy the differential equation

$$\frac{\mathrm{d}r}{\mathrm{d}t} = \frac{50 - r}{5r^2}.$$ [3]

(b) Find the quotient and remainder when $5r^2$ is divided by $50 - r$. [3]

DO NOT WRITE IN THIS MARGIN

(c) Solve the differential equation in part **(a)**, obtaining an expression for t in terms of r. [6]

(d) Find the value of t when the radius of the balloon is 12. [1]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

11 Let $f(x) = \dfrac{2e^{2x}}{e^{2x} - 3e^{x} + 2}$.

(a) Find $f'(x)$ and hence find the exact coordinates of the stationary point of the curve with equation $y = f(x)$. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) Use the substitution $u = e^x$ and partial fractions to find the exact value of $\int_{\ln 3}^{\ln 5} f(x)\,dx$.

Give your answer in the form $\ln a$, where a is a rational number in its simplest form. [9]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

BLANK PAGE

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

BLANK PAGE

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

* 0000008000001 *

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

4045182736

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **October/November 2024**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages.

© UCLES 2024

[Turn over

1 The complex number z satisfies $|z| = 2$ and $0 \leqslant \arg z \leqslant \frac{1}{4}\pi$.

(a) On the Argand diagram below, sketch the locus of the points representing z. [2]

(b) On the **same diagram**, sketch the locus of the points representing z^2. [2]



DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

2 Let $f(x) = 2x^3 - 5x^2 + 4$.

(a) Show that if a sequence of values given by the iterative formula

$$x_{n+1} = \sqrt{\frac{4}{5 - 2x_n}}$$

converges, then it converges to a root of the equation $f(x) = 0$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The equation has a root close to 1.2 .

Use the iterative formula from part **(a)** and an initial value of 1.2 to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

3



The number of bacteria in a population, P, at time t hours is modelled by the equation $P = ae^{kt}$, where a and k are constants. The graph of $\ln P$ against t, shown in the diagram, has gradient $\frac{1}{20}$ and intersects the vertical axis at $(0, 3)$.

(a) State the value of k and find the value of a correct to 2 significant figures. [3]

.....

.....

.....

.....

.....

.....

.....

(b) Find the time taken for P to double. Give your answer correct to the nearest hour. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN

4 Find the complex number z satisfying the equation

$$\frac{z-3\mathrm{i}}{z+3\mathrm{i}} = \frac{2-9\mathrm{i}}{5}.$$

Give your answer in the form $x+\mathrm{i}y$, where x and y are real. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

5 **(a)** Show that $\cos^4\theta - \sin^4\theta - 4\sin^2\theta\cos^2\theta \equiv \cos^2 2\theta + \cos 2\theta - 1$. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) Solve the equation $\cos^4\alpha - \sin^4\alpha = 4\sin^2\alpha\cos^2\alpha$ for $0° \leqslant \alpha \leqslant 180°$. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

6 The lines l and m have vector equations

$$l: \quad \mathbf{r} = 2\mathbf{i} + \mathbf{j} - 3\mathbf{k} + \lambda(-\mathbf{i} + 2\mathbf{k}) \quad \text{and} \quad m: \quad \mathbf{r} = 2\mathbf{i} + \mathbf{j} - 3\mathbf{k} + \mu(2\mathbf{i} - \mathbf{j} + 5\mathbf{k}).$$

Lines l and m intersect at the point P.

(a) State the coordinates of P. [1]

.....

.....

.....

.....

.....

.....

.....

(b) Find the exact value of the cosine of the acute angle between l and m. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN

(c) The point A on line l has coordinates $(0,1,1)$. The point B on line m has coordinates $(0,2,-8)$.

Find the exact area of triangle APB. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

7 The parametric equations of a curve are

$$x = 3\sin 2t, \quad y = \tan t + \cot t,$$

for $0 < t < \frac{1}{2}\pi$.

(a) Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{-2}{3\sin^2 2t}$. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) Find the equation of the normal to the curve at the point where $t = \frac{1}{4}\pi$. Give your answer in the form $py + qx + r = 0$, where p, q and r are integers. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

8 Let $f(x) = \dfrac{7a^2}{(a-2x)(3a+x)}$, where a is a positive constant.

(a) Express $f(x)$ in partial fractions. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [4]

(c) State the set of values of x for which the expansion in part **(b)** is valid. [1]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

9 **(a)** Find the quotient and remainder when $x^4 + 16$ is divided by $x^2 + 4$. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) Hence show that $\int_2^{2\sqrt{3}} \frac{x^4+16}{x^2+4}\,\mathrm{d}x = \frac{4}{3}(\pi+4)$. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

10 A water tank is in the shape of a cuboid with base area $40\,000\,\text{cm}^2$. At time t minutes the depth of water in the tank is h cm. Water is pumped into the tank at a rate of $50\,000\,\text{cm}^3$ per minute. Water is leaking out of the tank through a hole in the bottom at a rate of $600h\,\text{cm}^3$ per minute.

(a) Show that $200\dfrac{\mathrm{d}h}{\mathrm{d}t} = 250 - 3h$. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) It is given that when $t = 0$, $h = 50$.

Find the time taken for the depth of water in the tank to reach 80 cm. Give your answer correct to 2 significant figures. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

11



The diagram shows the curve $y = 2\sin x\sqrt{2+\cos x}$, for $0 \leqslant x \leqslant 2\pi$, and its minimum point M, where $x = a$.

(a) Find the value of a correct to 2 decimal places. [5]

DO NOT WRITE IN THIS MARGIN

(b) Use the substitution $u = 2 + \cos x$ to find the exact area of the shaded region R. [6]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

SAT PREP SP www.satprep.co.in

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER | | | | | |

CANDIDATE NUMBER | | | | |

3353608262

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **May/June 2024**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2024

1 Expand $(3+x)(1-2x)^{\frac{1}{2}}$ in ascending powers of x, up to and including the term in x^2, simplifying the coefficients. [4]

SAT PREP SP www.satprep.co.in

2 Solve the equation $\ln(x-5) = 7 - \ln x$. Give your answer correct to 2 decimal places. [4]

SAT PREP SP www.satprep.co.uk

3



The variables x and y satisfy the equation $a^y = bx$, where a and b are constants. The graph of y against $\ln x$ is a straight line passing through the points $(0.336, 1.00)$ and $(1.31, 1.50)$, as shown in the diagram.

Find the values of a and b. Give each value correct to the nearest integer. [4]

SAT PREP SP www.satprep.co.in

4 The complex number u is given by $u = -1 - i\sqrt{3}$.

(a) Express u in the form $r(\cos\theta + i\sin\theta)$, where $r > 0$ and $-\pi < \theta \leqslant \pi$. Give the exact values of r and θ. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The complex number v is given by $v = 5\left(\cos\frac{1}{6}\pi + i\sin\frac{1}{6}\pi\right)$.

(b) Express the complex number $\frac{v}{u}$ in the form $re^{i\theta}$ where $r > 0$ and $-\pi < \theta \leqslant \pi$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 The equation of a curve is $y = \dfrac{e^{\sin x}}{\cos^2 x}$ for $0 \leqslant x \leqslant 2\pi$.

Find $\dfrac{dy}{dx}$ and hence find the x-coordinates of the stationary points of the curve. [7]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

6 **(a)** By sketching a suitable pair of graphs, show that the equation $\operatorname{cosec}\frac{1}{2}x = e^x - 3$ has exactly one root, denoted by α, in the interval $0 < x < \pi$. [2]

(b) Verify by calculation that α lies between 1 and 2. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(c) Show that if a sequence of values in the interval $0 < x < \pi$ given by the iterative formula

$$x_{n+1} = \ln(\operatorname{cosec} \tfrac{1}{2}x_n + 3)$$

converges, then it converges to α. [1]

.....

.....

.....

.....

.....

.....

.....

(d) Use this iterative formula with an initial value of 1.4 to determine α correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(e) State the minimum number of calculated iterations needed with this initial value to determine α correct to 2 decimal places. [1]

.....

.....

© UCLES 2024

BLANK PAGE

SAT PREP SP www.satprep.co.in

7 **(a)** On a single Argand diagram sketch the loci given by the equations $|z-3+2\mathrm{i}|=2$ and $|w-3+2\mathrm{i}|=|w+3-4\mathrm{i}|$ where z and w are complex numbers. [4]

(b) Hence find the least value of $|z-w|$ for points on these loci. Give your answer in an exact form. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2024

8 Use the substitution $u = 1 - \sin x$ to find the exact value of

$$\int_{\pi}^{\frac{3}{2}\pi} \frac{\sin 2x}{\sqrt{1-\sin x}} \mathrm{d}x.$$

Give your answer in the form $a + b\sqrt{2}$ where a and b are rational numbers to be determined. [7]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

9 The equations of two straight lines l_1 and l_2 are

$$l_1: \quad \mathbf{r} = \mathbf{i} - 2\mathbf{j} + 3\mathbf{k} + \lambda(2\mathbf{i} - \mathbf{j} + a\mathbf{k}) \quad \text{and} \quad l_2: \quad \mathbf{r} = -\mathbf{i} - \mathbf{j} - \mathbf{k} + \mu(3\mathbf{i} - 2\mathbf{j} - 2\mathbf{k}),$$

where a is a constant.

The lines l_1 and l_2 are perpendicular.

(a) Show that $a = 4$. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The lines l_1 and l_2 also intersect.

(b) Find the position vector of the point of intersection. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2024

The point A has position vector $-5\mathbf{i}+\mathbf{j}-9\mathbf{k}$.

(c) Show that A lies on l_1. [2]

The point B is the image of A after a reflection in the line l_2.

(d) Find the position vector of B. [2]

10 **(a)** Given that $2x = \tan y$, show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{2}{1+4x^2}$. [3]

(b) Hence find the exact value of $\displaystyle\int_{\frac{1}{2}}^{\frac{\sqrt{3}}{2}} x\tan^{-1}(2x)\,\mathrm{d}x$. [7]

© UCLES 2024

SAT PREP SP www.satprep.co.in

11 In a field there are 300 plants of a certain species, all of which can be infected by a particular disease. At time t after the first plant is infected there are x infected plants. The rate of change of x is proportional to the product of the number of plants infected and the number of plants that are **not** yet infected. The variables x and t are treated as continuous, and it is given that $\frac{dx}{dt} = 0.2$ and $x = 1$ when $t = 0$.

(a) Show that x and t satisfy the differential equation

$$1495 \frac{dx}{dt} = x(300 - x). \quad [2]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Using partial fractions, solve the differential equation and obtain an expression for t in terms of a single logarithm involving x. [9]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2024

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

8567013635

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **May/June 2024**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2024

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 **(a)** Sketch the graph of $y = |x - 2a|$, where a is a positive constant. [1]

(b) Solve the inequality $2x - 3a < |x - 2a|$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

2 Express $\frac{6x^2 - 9x - 16}{2x^2 - 5x - 12}$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

3 The variables x and y satisfy the equation $a^{2y-1} = b^{x-y}$, where a and b are constants.

(a) Show that the graph of y against x is a straight line. [3]

(b) Given that $a = b^3$, state the equation of the straight line in the form $y = px + q$, where p and q are rational numbers in their simplest form. [2]

4 The equation of a curve is $ye^{2x} + y^2e^x = 6$.

Find the gradient of the curve at the point where $y = 1$. [6]

© UCLES 2024

5 **(a)** It is given that the equation $e^{2x} = 5 + \cos 3x$ has only one root.

Show by calculation that this root lies in the interval $0.7 < x < 0.8$. [2]

.....

.....

.....

.....

.....

.....

(b) Show that if a sequence of values in the interval $0.7 < x < 0.8$ given by the iterative formula

$$x_{n+1} = \frac{1}{2}\ln\left(5 + \cos 3x_n\right)$$

converges then it converges to the root of the equation in part **(a)**. [1]

.....

.....

.....

.....

.....

.....

(c) Use this iterative formula to determine the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2024

6



The diagram shows the curve $y = xe^{-ax}$, where a is a positive constant, and its maximum point M.

(a) Find the exact coordinates of M. [4]

SAT PREP SP www.satprep.co.in

(b) Find the exact value of $\int_0^{\frac{2}{a}} x\mathrm{e}^{-ax}\,\mathrm{d}x$. [5]

SAT PREP SP www.satprep.co.in

7 **(a)** Show that $\cos^4\theta - \sin^4\theta \equiv \cos 2\theta$. [3]

SAT PREP SP www.satprep.co.uk

(b) Hence find the exact value of $\int_{-\frac{1}{8}\pi}^{\frac{1}{8}\pi} \left(\cos^4\theta - \sin^4\theta + 4\sin^2\theta\cos^2\theta\right) \mathrm{d}\theta$. [6]

SAT PREP SP www.satprep.co.in

8 The points A, B and C have position vectors $\overrightarrow{OA} = -2\mathbf{i} + \mathbf{j} + 4\mathbf{k}$, $\overrightarrow{OB} = 5\mathbf{i} + 2\mathbf{j}$ and $\overrightarrow{OC} = 8\mathbf{i} + 5\mathbf{j} - 3\mathbf{k}$, where O is the origin. The line l_1 passes through B and C.

(a) Find a vector equation for l_1. [3]

.....

.....

.....

.....

.....

.....

.....

.....

The line l_2 has equation $\mathbf{r} = -2\mathbf{i} + \mathbf{j} + 4\mathbf{k} + \mu(3\mathbf{i} + \mathbf{j} - 2\mathbf{k})$.

(b) Find the coordinates of the point of intersection of l_1 and l_2. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2024

(c) The point D on l_2 is such that $AB = BD$.

Find the position vector of D. [5]

SAT PREP SP www.satprep.co.uk

9 The complex numbers z and ω are defined by $z = 1 - \mathrm{i}$ and $\omega = -3 + 3\sqrt{3}\mathrm{i}$.

(a) Express $z\omega$ in the form $a + b\mathrm{i}$, where a and b are real and in exact surd form. [1]

.....

.....

.....

.....

.....

(b) Express z and ω in the form $re^{\mathrm{i}\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$. Give the exact values of r and θ in each case. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) On an Argand diagram, the points representing ω and $z\omega$ are A and B respectively.

Prove that OAB is an isosceles right-angled triangle, where O is the origin. [2]

© UCLES 2024

(d) Using your answers to part **(b)**, prove that $\tan \frac{5}{12}\pi = \frac{\sqrt{3}+1}{\sqrt{3}-1}$. [3]

SAT PREP SP www.satprep.co.in

10 (a) By writing $y = \sec^3\theta$ as $\dfrac{1}{\cos^3\theta}$, show that $\dfrac{\mathrm{d}y}{\mathrm{d}\theta} = 3\sin\theta\sec^4\theta$. [2]

(b) The variables x and θ satisfy the differential equation

$$\left(x^2+9\right)\sin\theta\frac{\mathrm{d}\theta}{\mathrm{d}x} = (x+3)\cos^4\theta.$$

It is given that $x = 3$ when $\theta = \frac{1}{3}\pi$.

Solve the differential equation to find the value of $\cos\theta$ when $x = 0$. Give your answer correct to 3 significant figures. [8]

© UCLES 2024

SAT PREP SP www.satprep.co.in

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2024

* 0019655419901 *

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

* 2 6 3 4 3 8 9 6 3 5 *

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **May/June 2024**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2024

[Turn over

1 Solve the equation $8^{3-6x} = 4 \times 5^{-2x}$. Give your answer correct to 3 decimal places. [4]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

2 Find the exact coordinates of the stationary point of the curve $y = e^{2x} \sin 2x$ for $0 \leqslant x \leqslant \frac{1}{2}\pi$. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

3 The square roots of $24-7\mathrm{i}$ can be expressed in the Cartesian form $x+\mathrm{i}y$, where x and y are real and exact.

By first forming a quartic equation in x or y, find the square roots of $24-7\mathrm{i}$ in exact Cartesian form. [5]

DO NOT WRITE IN THIS MARGIN

4



The variables x and y satisfy the equation $ky = e^{cx}$, where k and c are constants. The graph of $\ln y$ against x is a straight line passing through the points $(2.80, 0.372)$ and $(5.10, 2.21)$, as shown in the diagram.

Find the values of k and c. Give each value correct to 2 significant figures. [4]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

5 Express $\frac{6x^2-2x+2}{(x-1)(2x+1)}$ in partial fractions. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

6 **(a)** On an Argand diagram shade the region whose points represent complex numbers z which satisfy both the inequalities $|z-4-3\mathrm{i}| \leqslant 2$ and $\arg(z-2-\mathrm{i}) \geqslant \frac{1}{3}\pi$. [5]

(b) Calculate the greatest value of $\arg z$ for points in this region. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

7 Let $f(x) = 8x^3 + 54x^2 - 17x - 21$.

(a) Show that $x + 7$ is a factor of $f(x)$. [1]

(b) Find the quotient when $f(x)$ is divided by $x + 7$. [2]

DO NOT WRITE IN THIS MARGIN

(c) Hence solve the equation

$$8\cos^3\theta + 54\cos^2\theta - 17\cos\theta - 21 = 0,$$

for $0° \leqslant \theta \leqslant 360°$. [3]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

8 **(a)** Express $3\cos 2x - \sqrt{3}\sin 2x$ in the form $R\cos(2x+\alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. Give the exact values of R and α. [3]

DO NOT WRITE IN THIS MARGIN

(b) Hence find the exact value of $\int_0^{\frac{1}{12}\pi} \frac{3}{\left(3\cos 2x - \sqrt{3}\sin 2x\right)^2}\,\mathrm{d}x$, simplifying your answer. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

9



A container in the shape of a cuboid has a square base of side x and a height of $(10-x)$. It is given that x varies with time, t, where $t > 0$. The container decreases in volume at a rate which is inversely proportional to t.

When $t = \frac{1}{10}$, $x = \frac{1}{2}$ and the rate of decrease of x is $\frac{20}{37}$.

(a) Show that x and t satisfy the differential equation

$$\frac{\mathrm{d}x}{\mathrm{d}t} = \frac{-1}{2t\left(20x - 3x^2\right)}.$$ [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

(b) Solve the differential equation, obtaining an expression for t in terms of x. [6]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

10 The equations of two straight lines are

$$\mathbf{r} = \mathbf{i} + \mathbf{j} + 2a\mathbf{k} + \lambda(3\mathbf{i} + 4\mathbf{j} + a\mathbf{k}) \quad \text{and} \quad \mathbf{r} = -3\mathbf{i} - \mathbf{j} + 4\mathbf{k} + \mu(-\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}),$$

where a is a constant.

(a) Given that the acute angle between the directions of these lines is $\frac{1}{4}\pi$, find the possible values of a. [6]

DO NOT WRITE IN THIS MARGIN

SAT PREP SP www.satprep.co.in

(b) Given instead that the lines intersect, find the value of a and the position vector of the point of intersection. [5]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

11 Use the substitution $2x = \tan\theta$ to find the exact value of

$$\int_0^{\frac{1}{2}} \frac{12}{\left(1+4x^2\right)^2}\,\mathrm{d}x\ .$$

Give your answer in the form $a+b\pi$, where a and b are rational numbers. [9]

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

BLANK PAGE

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

BLANK PAGE

DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN
DO NOT WRITE IN THIS MARGIN

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

5552004552

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **February/March 2024**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2024

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Find the quotient and remainder when $x^4 - 3x^3 + 9x^2 - 12x + 27$ is divided by $x^2 + 5$. [3]

SAT PREP SP www.satprep.co.uk

2 **(a)** Find the coefficient of x^2 in the expansion of $(2x-5)\sqrt{4-x}$. [4]

(b) State the set of values of x for which the expansion in part **(a)** is valid. [1]

SAT PREP SP www.satprep.co.uk

3 It is given that $z = -\sqrt{3} + \mathrm{i}$.

(a) Express z^2 in the form $r\mathrm{e}^{\mathrm{i}\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$. [3]

(b) The complex number ω is such that $z^2\omega$ is real and $\left|\frac{z^2}{\omega}\right| = 12$.

Find the two possible values of ω, giving your answers in the form $R\mathrm{e}^{\mathrm{i}\alpha}$, where $R > 0$ and $-\pi < \alpha \leqslant \pi$. [3]

4 The positive numbers p and q are such that

$$\ln\left(\frac{p}{q}\right) = a \quad \text{and} \quad \ln\left(q^2 p\right) = b.$$

Express $\ln\left(p^7 q\right)$ in terms of a and b. [4]

SAT PREP SP www.satprep.co.in

5 **(a)** On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z-4-2\mathrm{i}| \leqslant 3$ and $|z| \geqslant |10-z|$. [4]

(b) Find the greatest value of $\arg z$ for points in this region. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2024

6 The equation of a curve is $2y^2 + 3xy + x = x^2$.

(a) Show that $\frac{dy}{dx} = \frac{2x - 3y - 1}{4y + 3x}$. [4]

SAT PREP www.satprep.co.in

(b) Hence show that the curve does **not** have a tangent that is parallel to the x-axis. [3]

SAT PREP SP www.satprep.co.in

7



The diagram shows the curve $y = xe^{2x} - 5x$ and its minimum point M, where $x = \alpha$.

(a) Show that α satisfies the equation $\alpha = \frac{1}{2}\ln\left(\frac{5}{1+2\alpha}\right)$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2024

(b) Verify by calculation that α lies between 0.4 and 0.5 . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in part **(a)** to determine α correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2024

8 **(a)** Express $3\sin x + 2\sqrt{2}\cos\left(x + \frac{1}{4}\pi\right)$ in the form $R\sin(x+\alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. State the exact value of R and give α correct to 3 decimal places. [4]

SAT PREP SP www.satprep.co.in

(b) Hence solve the equation

$$6\sin\tfrac{1}{2}\theta + 4\sqrt{2}\cos\left(\tfrac{1}{2}\theta + \tfrac{1}{4}\pi\right) = 3$$

for $-4\pi < \theta < 4\pi$. [5]

9 Relative to the origin O, the position vectors of the points A, B and C are given by

$$\overrightarrow{OA} = 5\mathbf{i} - 2\mathbf{j} + \mathbf{k}, \quad \overrightarrow{OB} = 8\mathbf{i} + 2\mathbf{j} - 6\mathbf{k} \quad \text{and} \quad \overrightarrow{OC} = 3\mathbf{i} + 4\mathbf{j} - 7\mathbf{k}.$$

(a) Show that $OABC$ is a rectangle. [4]

SAT PREP SP www.satprep.co.in

(b) Use a scalar product to find the acute angle between the diagonals of $OABC$. [4]

SAT PREP SP www.satprep.co.in

10 Let $f(x) = \dfrac{36a^2}{(2a+x)(2a-x)(5a-2x)}$, where a is a positive constant.

(a) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence find the exact value of $\int_{-a}^{a} \mathrm{f}(x)\,\mathrm{d}x$, giving your answer in the form $p\ln q + r\ln s$ where p and r are integers and q and s are prime numbers. [5]

SAT PREP SP www.satprep.co.uk

11 The variables y and θ satisfy the differential equation

$$(1+y)(1+\cos 2\theta)\frac{\mathrm{d}y}{\mathrm{d}\theta} = \mathrm{e}^{3y}.$$

It is given that $y = 0$ when $\theta = \frac{1}{4}\pi$.

Solve the differential equation and find the exact value of $\tan\theta$ when $y = 1$. [9]

SAT PREP SP www.satprep.co.uk

SAT PREP SP www.satprep.co.in

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2024

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

5051231203

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **October/November 2023**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2023

1 Find the exact coordinates of the points on the curve $y = \dfrac{x^2}{1 - 3x}$ at which the gradient of the tangent is equal to 8. [5]

SAT PREP SP www.satprep.co.in

2 On an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 2i| \leqslant |z + 2 - i|$ and $0 \leqslant \arg(z + 1) \leqslant \frac{1}{4}\pi$. [4]

SAT PREP SP www.satprep.co.in

3



The variables x and y are related by the equation $y = ab^x$, where a and b are constants. The diagram shows the result of plotting $\ln y$ against x for two pairs of values of x and y. The coordinates of these points are (1, 3.7) and (2.2, 6.46).

Use this information to find the values of a and b. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

4 The complex number u is defined by $u = \dfrac{3 + 2\mathrm{i}}{a - 5\mathrm{i}}$, where a is real.

(a) Express u in the Cartesian form $x + \mathrm{i}y$, where x and y are in terms of a. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Given that $\arg u = \frac{1}{4}\pi$, find the value of a. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

5 **(a)** Given that

$$\sin\left(x+\tfrac{1}{6}\pi\right)-\sin\left(x-\tfrac{1}{6}\pi\right)=\cos\left(x+\tfrac{1}{3}\pi\right)-\cos\left(x-\tfrac{1}{3}\pi\right),$$

find the exact value of $\tan x$. [4]

SAT PREP SP www.satprep.co.in

(b) Hence find the exact roots of the equation

$$\sin\left(x+\tfrac{1}{6}\pi\right)-\sin\left(x-\tfrac{1}{6}\pi\right)=\cos\left(x+\tfrac{1}{3}\pi\right)-\cos\left(x-\tfrac{1}{3}\pi\right)$$

for $0 \leqslant x \leqslant 2\pi$. [2]

SAT PREP SP www.satprep.co.in

6 The parametric equations of a curve are

$$x = \sqrt{t} + 3, \qquad y = \ln t,$$

for $t > 0$.

(a) Obtain a simplified expression for $\frac{\mathrm{d}y}{\mathrm{d}x}$ in terms of t. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence find the exact coordinates of the point on the curve at which the gradient of the normal is -2. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

7 The variables x and θ satisfy the differential equation

$$\frac{x}{\tan\theta}\frac{\mathrm{d}x}{\mathrm{d}\theta} = x^2 + 3.$$

It is given that $x = 1$ when $\theta = 0$.

Solve the differential equation, obtaining an expression for x^2 in terms of θ. [7]

SAT PREP SP www.satprep.co.in

8 **(a)** By sketching a suitable pair of graphs, show that the equation

$$\sqrt{x} = e^x - 3$$

has only one root. [2]

(b) Show by calculation that this root lies between 1 and 2. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(c) Show that, if a sequence of values given by the iterative formula

$$x_{n+1} = \ln\left(3 + \sqrt{x_n}\right)$$

converges, then it converges to the root of the equation in **(a)**. [1]

.....

.....

.....

.....

.....

.....

.....

(d) Use the iterative formula to calculate the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

[Turn over

9



The diagram shows the curve $y = xe^{-\frac{1}{4}x^2}$, for $x \geqslant 0$, and its maximum point M.

(a) Find the exact coordinates of M. [4]

SAT PREP SP www.satprep.co.uk

(b) Using the substitution $x = \sqrt{u}$, or otherwise, find by integration the exact area of the shaded region bounded by the curve, the x-axis and the line $x = 3$. [5]

10 Let $\mathrm{f}(x) = \dfrac{24x + 13}{(1 - 2x)(2 + x)^2}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) State the set of values of x for which the expansion in **(b)** is valid. [1]

.....

.....

.....

.....

© UCLES 2023

11



In the diagram, $OABCDEFG$ is a cuboid in which $OA = 3$ units, $OC = 2$ units and $OD = 2$ units. Unit vectors $\mathbf{i}$, $\mathbf{j}$ and $\mathbf{k}$ are parallel to OA, OD and OC respectively. M is the midpoint of EF.

(a) Find the position vector of M. [1]

.....

.....

.....

The position vector of P is $\mathbf{i} + \mathbf{j} + 2\mathbf{k}$.

(b) Calculate angle PAM. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

(c) Find the exact length of the perpendicular from P to the line passing through O and M. [5]

SAT PREP SP www.satprep.co.uk

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

9146949940

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **October/November 2023**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2023

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 **(a)** Sketch the graph of $y = |4x - 2|$. [1]

(b) Solve the inequality $1 + 3x < |4x - 2|$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

2 The parametric equations of a curve are

$$x = (\ln t)^2, \qquad y = \mathrm{e}^{2-t^2},$$

for $t > 0$.

Find the gradient of the curve at the point where $t = \mathrm{e}$, simplifying your answer. [4]

SAT PREP SP www.satprep.co.in

3 The polynomial $2x^3 + ax^2 - 11x + b$ is denoted by $p(x)$. It is given that $p(x)$ is divisible by $(2x - 1)$ and that when $p(x)$ is divided by $(x + 1)$ the remainder is 12.

Find the values of a and b. [5]

SAT PREP SP www.satprep.co.in

4 **(a)** On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 4 - 3i| \leqslant 2$ and $\mathrm{Re}\, z \leqslant 3$. [4]

(b) Find the greatest value of $\arg z$ for points in this region. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

5 Find the exact value of $\int_0^6 \frac{x(x+1)}{x^2+4}\,\mathrm{d}x$. [6]

SAT PREP SP www.satprep.co.in

6 **(a)** By sketching a suitable pair of graphs, show that the equation

$$\cot x = 2 - \cos x$$

has one root in the interval $0 < x \leqslant \frac{1}{2}\pi$. [2]

(b) Show by calculation that this root lies between 0.6 and 0.8. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(c) Use the iterative formula $x_{n+1} = \tan^{-1}\left(\dfrac{1}{2 - \cos x_n}\right)$ to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

SAT PREP SP www.satprep.co.uk

7 **(a)** By expressing 3θ as $2\theta + \theta$, prove the identity $\cos 3\theta \equiv 4\cos^3\theta - 3\cos\theta$. [3]

SAT PREP SP www.satprep.co.uk

(b) Hence solve the equation

$$\cos 3\theta + \cos \theta \cos 2\theta = \cos^2 \theta$$

for $0° \leqslant \theta \leqslant 180°$. [5]

8 It is given that $\dfrac{2+3a\mathrm{i}}{a+2\mathrm{i}} = \lambda(2-\mathrm{i})$, where a and λ are real constants.

(a) Show that $3a^2 + 4a - 4 = 0$. [4]

SAT PREP SP www.satprep.co.in

(b) Hence find the possible values of a and the corresponding values of λ. [3]

9



The diagram shows the curve $y = \sin x \cos 2x$, for $0 \leqslant x \leqslant \pi$, and a maximum point M, where $x = a$. The shaded region between the curve and the x-axis is denoted by R.

(a) Find the value of a correct to 2 decimal places. [5]

SAT PREP SP www.satprep.co.uk

(b) Find the exact area of the region R, giving your answer in simplified form. [4]

SAT PREP SP www.satprep.co.in

10 The equations of the lines l and m are given by

$$l: \ \mathbf{r} = \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} \qquad \text{and} \qquad m: \ \mathbf{r} = \begin{pmatrix} 6 \\ -3 \\ 6 \end{pmatrix} + \mu \begin{pmatrix} -2 \\ 4 \\ c \end{pmatrix},$$

where c is a positive constant. It is given that the angle between l and m is 60°.

(a) Find the value of c. [4]

SAT PREP SP www.satprep.co.in

(b) Show that the length of the perpendicular from $(6, -3, 6)$ to l is $\sqrt{11}$. [5]

SAT PREP SP www.satprep.co.in

11 The variables x and y satisfy the differential equation

$$x^2 \frac{\mathrm{d}y}{\mathrm{d}x} + y^2 + y = 0.$$

It is given that $x = 1$ when $y = 1$.

(a) Solve the differential equation to obtain an expression for y in terms of x. [8]

SAT PREP SP www.satprep.co.in

(b) State what happens to the value of y when x tends to infinity. Give your answer in an exact form. [1]

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2023

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

7354647755

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **October/November 2023**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2023

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Find the set of values of x satisfying the inequality $\left|2^{x+1} - 2\right| < 0.5$, giving your answer to 3 significant figures. [4]

SAT PREP SP www.satprep.co.in

2 On an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 1 + 2i| \leqslant |z|$ and $|z - 2| \leqslant 1$. [5]

SAT PREP SP www.satprep.co.in

3 The polynomial $2x^3 + ax^2 + bx + 6$, where a and b are constants, is denoted by $\mathrm{p}(x)$. When $\mathrm{p}(x)$ is divided by $(x + 2)$ the remainder is -38 and when $\mathrm{p}(x)$ is divided by $(2x - 1)$ the remainder is $\frac{19}{2}$.

Find the values of a and b. [5]

SAT PREP SP www.satprep.co.in

4 Solve the quadratic equation $(3 + \mathrm{i})w^2 - 2w + 3 - \mathrm{i} = 0$, giving your answers in the form $x + \mathrm{i}y$, where x and y are real. [5]

© UCLES 2023

5 Find the exact coordinates of the stationary points of the curve $y = \dfrac{e^{3x^2-1}}{1-x^2}$. [6]

SAT PREP SP www.satprep.co.in

6 **(a)** Show that the equation $\cot^2\theta + 2\cos 2\theta = 4$ can be written in the form

$$4\sin^4\theta + 3\sin^2\theta - 1 = 0.$$ [3]

SAT PREP SP www.satprep.co.uk

(b) Hence solve the equation $\cot^2\theta + 2\cos 2\theta = 4$, for $0° < \theta < 360°$. [3]

SAT PREP SP www.satprep.co.in

7 The equation of a curve is $x^3 + y^2 + 3x^2 + 3y = 4$.

(a) Show that $\frac{dy}{dx} = -\frac{3x^2 + 6x}{2y + 3}$. [3]

SAT PREP SP www.satprep.co.in

(b) Hence find the coordinates of the points on the curve at which the tangent is parallel to the x-axis. [5]

8 The variables x and y satisfy the differential equation

$$e^{4x}\frac{dy}{dx} = \cos^2 3y.$$

It is given that $y = 0$ when $x = 2$.

Solve the differential equation, obtaining an expression for y in terms of x. [7]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

9 Let $f(x) = \dfrac{17x^2 - 7x + 16}{(2 + 3x^2)(2 - x)}$.

(a) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^3. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) State the set of values of x for which the expansion in **(b)** is valid. Give your answer in an exact form. [1]

.....

.....

.....

.....

.....

© UCLES 2023

[Turn over

10



The diagram shows the curve $y = x\cos 2x$, for $x \geqslant 0$.

(a) Find the equation of the tangent to the curve at the point where $x = \frac{1}{2}\pi$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

(b) Find the exact area of the shaded region shown in the diagram, bounded by the curve and the x-axis. [5]

11 The line l has equation $\mathbf{r} = \mathbf{i} - 2\mathbf{j} - 3\mathbf{k} + \lambda(-\mathbf{i} + \mathbf{j} + 2\mathbf{k})$. The points A and B have position vectors $-2\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ and $3\mathbf{i} - \mathbf{j} + \mathbf{k}$ respectively.

(a) Find a unit vector in the direction of l. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The line m passes through the points A and B.

(b) Find a vector equation for m. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(c) Determine whether lines l and m are parallel, intersect or are skew. [5]

SAT PREP SP www.satprep.co.uk

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

2968750175

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **May/June 2023**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2023

1 Solve the equation

$$3e^{2x} - 4e^{-2x} = 5.$$

Give the answer correct to 3 decimal places. [3]

SAT PREP SP www.satprep.co.in

2 **(a)** Sketch the graph of $y = |2x + 3|$. [1]

(b) Solve the inequality $3x + 8 > |2x + 3|$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

3 Find the coefficient of x^3 in the binomial expansion of $(3 + x)\sqrt{1 + 4x}$. [4]

SAT PREP SP www.satprep.co.in

4 **(a)** Show that the equation $\sin 2\theta + \cos 2\theta = 2\sin^2\theta$ can be expressed in the form

$$\cos^2\theta + 2\sin\theta\cos\theta - 3\sin^2\theta = 0.$$ [2]

.....

.....

.....

.....

.....

.....

.....

(b) Hence solve the equation $\sin 2\theta + \cos 2\theta = 2\sin^2\theta$ for $0° < \theta < 180°$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP www.satprep.co.in

5 The equation of a curve is $x^2y - ay^2 = 4a^3$, where a is a non-zero constant.

(a) Show that $\frac{dy}{dx} = \frac{2xy}{2ay - x^2}$. [4]

(b) Hence find the coordinates of the points where the tangent to the curve is parallel to the y-axis. [4]

6 Relative to the origin O, the points A, B and C have position vectors given by

$$\overrightarrow{OA} = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}, \qquad \overrightarrow{OB} = \begin{pmatrix} 4 \\ 3 \\ 2 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OC} = \begin{pmatrix} 3 \\ -2 \\ -4 \end{pmatrix}.$$

The quadrilateral $ABCD$ is a parallelogram.

(a) Find the position vector of D. [3]

SAT PREP SP www.satprep.co.in

(b) The angle between BA and BC is θ.

Find the exact value of $\cos\theta$. [3]

.....

.....

.....

.....

.....

.....

.....

(c) Hence find the area of $ABCD$, giving your answer in the form $p\sqrt{q}$, where p and q are integers. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

[Turn over

7 The variables x and y satisfy the differential equation

$$\cos 2x \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{4 \tan 2x}{\sin^2 3y},$$

where $0 \leqslant x < \frac{1}{4}\pi$. It is given that $y = 0$ when $x = \frac{1}{6}\pi$.

Solve the differential equation to obtain the value of x when $y = \frac{1}{6}\pi$. Give your answer correct to 3 decimal places. [8]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

8 Let $f(x) = \dfrac{3 - 3x^2}{(2x + 1)(x + 2)^2}$.

(a) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence find the exact value of $\int_0^4 \mathrm{f}(x)\,\mathrm{d}x$, giving your answer in the form $a + b\ln c$, where a, b and c are integers. [5]

SAT PREP SP www.satprep.co.in

9 The constant a is such that $\int_0^a xe^{-2x}\,dx = \frac{1}{8}$.

(a) Show that $a = \frac{1}{2}\ln(4a + 2)$. [5]

SAT PREP SP www.satprep.co.in

(b) Verify by calculation that a lies between 0.5 and 1. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in **(a)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

[Turn over

10 The polynomial $x^3 + 5x^2 + 31x + 75$ is denoted by $\mathrm{p}(x)$.

(a) Show that $(x + 3)$ is a factor of $\mathrm{p}(x)$. [2]

.....

.....

.....

.....

.....

.....

.....

(b) Show that $z = -1 + 2\sqrt{6}\mathrm{i}$ is a root of $\mathrm{p}(z) = 0$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(c) Hence find the complex numbers z which are roots of $p(z^2) = 0$. [7]

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

7943719452

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **May/June 2023**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2023

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Solve the inequality $|5x - 3| < 2|3x - 7|$. [4]

SAT PREP SP www.satprep.co.in

2 Solve the equation $\ln(2x^2 - 3) = 2\ln x - \ln 2$, giving your answer in an exact form. [3]

SAT PREP SP www.satprep.co.in

3 **(a)** On an Argand diagram, sketch the locus of points representing complex numbers z satisfying $|z + 3 - 2i| = 2$. [2]

(b) Find the least value of $|z|$ for points on this locus, giving your answer in an exact form. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

[Turn over

4 Solve the equation $2\cos x - \cos \frac{1}{2}x = 1$ for $0 \leqslant x \leqslant 2\pi$. [5]

SAT PREP SP www.satprep.co.uk

5 The complex number $2 + yi$ is denoted by a, where y is a real number and $y < 0$. It is given that $f(a) = a^3 - a^2 - 2a$.

(a) Find a simplified expression for $f(a)$ in terms of y. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Given that $\text{Re}\big(f(a)\big) = -20$, find $\arg a$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

6 The equation $\cot \frac{1}{2}x = 3x$ has one root in the interval $0 < x < \pi$, denoted by α.

(a) Show by calculation that α lies between 0.5 and 1. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show that, if a sequence of positive values given by the iterative formula

$$x_{n+1} = \frac{1}{3}\left(x_n + 4\tan^{-1}\left(\frac{1}{3x_n}\right)\right)$$

converges, then it converges to α. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

(c) Use this iterative formula to calculate α correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

SAT PREP SP www.satprep.co.in

7 The equation of a curve is $3x^2 + 4xy + 3y^2 = 5$.

(a) Show that $\frac{dy}{dx} = -\frac{3x + 2y}{2x + 3y}$. [4]

SAT PREP SP www.satprep.co.in

(b) Hence find the exact coordinates of the two points on the curve at which the tangent is parallel to $y + 2x = 0$. [5]

© UCLES 2023

[Turn over

8 **(a)** The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{4 + 9y^2}{\mathrm{e}^{2x+1}}.$$

It is given that $y = 0$ when $x = 1$.

Solve the differential equation, obtaining an expression for y in terms of x. [7]

SAT PREP SP www.satprep.co.uk

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State what happens to the value of y as x tends to infinity. Give your answer in an exact form. [1]

.....

.....

.....

.....

.....

9 Let $f(x) = \dfrac{2x^2 + 17x - 17}{(1 + 2x)(2 - x)^2}$.

(a) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence show that $\int_0^1 f(x)\,dx = \frac{5}{2} - \ln 72$. [5]

SAT PREP SP www.satprep.co.in

10



The diagram shows the curve $y = (x + 5)\sqrt{3 - 2x}$ and its maximum point M.

(a) Find the exact coordinates of M. [5]

SAT PREP SP www.satprep.co.in

(b) Using the substitution $u = 3 - 2x$, find by integration the area of the shaded region bounded by the curve and the x-axis. Give your answer in the form $a\sqrt{13}$, where a is a rational number. [5]

11 The points A and B have position vectors $\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}$ and $2\mathbf{i} - \mathbf{j} + \mathbf{k}$ respectively. The line l has equation $\mathbf{r} = \mathbf{i} - \mathbf{j} + 3\mathbf{k} + \mu(2\mathbf{i} - 3\mathbf{j} + 4\mathbf{k})$.

(a) Show that l does not intersect the line passing through A and B. [5]

SAT PREP SP www.satprep.co.in

(b) Find the position vector of the foot of the perpendicular from A to l. [4]

SAT PREP SP www.satprep.co.uk

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2023

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

5471210840

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **May/June 2023**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages.

© UCLES 2023

1 Solve the equation $\ln(x + 5) = 5 + \ln x$. Give your answer correct to 3 decimal places. [4]

SAT PREP SP www.satprep.co.uk

2 Find the quotient and remainder when $2x^4 - 27$ is divided by $x^2 + x + 3$. [3]

SAT PREP SP www.satprep.co.in

3 On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 3 - \mathrm{i}| \leqslant 3$ and $|z| \geqslant |z - 4\mathrm{i}|$. [4]

SAT PREP SP www.satprep.co.in

4 The parametric equations of a curve are

$$x = \frac{\cos\theta}{2 - \sin\theta}, \qquad y = \theta + 2\cos\theta.$$

Show that $\frac{\mathrm{d}y}{\mathrm{d}x} = (2 - \sin\theta)^2$. [5]

SAT PREP SP www.satprep.co.in

5



The diagram shows the part of the curve $y = x^2 \cos 3x$ for $0 \leqslant x \leqslant \frac{1}{6}\pi$, and its maximum point M, where $x = a$.

(a) Show that a satisfies the equation $a = \frac{1}{3}\tan^{-1}\left(\frac{2}{3a}\right)$. [3]

SAT PREP SP www.satprep.co.in

(b) Use an iterative formula based on the equation in **(a)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

SAT PREP SP www.satprep.co.in

6 **(a)** Express $3\cos x + 2\cos(x - 60°)$ in the form $R\cos(x - \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. State the exact value of R and give α correct to 2 decimal places. [4]

SAT PREP SP www.satprep.co.in

(b) Hence solve the equation

$$3\cos 2\theta + 2\cos(2\theta - 60°) = 2.5$$

for $0° < \theta < 180°$. [4]

SAT PREP SP www.satprep.co.in

7 **(a)** Use the substitution $u = \cos x$ to show that

$$\int_0^{\pi} \sin 2x \, e^{2\cos x} \, dx = \int_{-1}^{1} 2ue^{2u} \, du. \quad [4]$$

SAT PREP SP www.satprep.co.in

(b) Hence find the exact value of $\int_0^{\pi} \sin 2x \, e^{2\cos x} \, dx$. [4]

SAT PREP SP www.satprep.co.in

8 The variables x and y satisfy the differential equation

$$\frac{dy}{dx} = \frac{y^2 + 4}{x(y + 4)}$$

for $x > 0$. It is given that $x = 4$ when $y = 2\sqrt{3}$.

Solve the differential equation to obtain the value of x when $y = 2$. [8]

SAT PREP SP www.satprep.co.uk

SAT PREP SP www.satprep.co.in

9 The lines l and m have equations

$$l: \quad \mathbf{r} = a\mathbf{i} + 3\mathbf{j} + b\mathbf{k} + \lambda(c\mathbf{i} - 2\mathbf{j} + 4\mathbf{k}),$$
$$m: \quad \mathbf{r} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k} + \mu(2\mathbf{i} - 3\mathbf{j} + \mathbf{k}).$$

Relative to the origin O, the position vector of the point P is $4\mathbf{i} + 7\mathbf{j} - 2\mathbf{k}$.

(a) Given that l is perpendicular to m and that P lies on l, find the values of the constants a, b and c. [4]

SAT PREP SP www.satprep.co.in

(b) The perpendicular from P meets line m at Q. The point R lies on PQ extended, with $PQ : QR = 2 : 3$.

Find the position vector of R. [6]

SAT PREP SP www.satprep.co.in

10 Let $\mathrm{f}(x) = \dfrac{21 - 8x - 2x^2}{(1 + 2x)(3 - x)^2}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

11 The complex number z is defined by $z = \dfrac{5a - 2\mathrm{i}}{3 + a\mathrm{i}}$, where a is an integer. It is given that $\arg z = -\frac{1}{4}\pi$.

(a) Find the value of a and hence express z in the form $x + \mathrm{i}y$, where x and y are real. [6]

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Express z^3 in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$. Give the simplified exact values of r and θ. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

9337495756

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **February/March 2023**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages.

© UCLES 2023

1 It is given that $x = \ln(2y - 3) - \ln(y + 4)$.

Express y in terms of x. [3]

SAT PREP SP www.satprep.co.in

2 **(a)** On an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $-\frac{1}{3}\pi \leqslant \arg(z - 1 - 2\mathrm{i}) \leqslant \frac{1}{3}\pi$ and $\mathrm{Re}\, z \leqslant 3$. [3]

(b) Calculate the least value of $\arg z$ for points in the region from **(a)**. Give your answer in radians correct to 3 decimal places. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2023

3 The polynomial $2x^4 + ax^3 + bx - 1$, where a and b are constants, is denoted by $p(x)$. When $p(x)$ is divided by $x^2 - x + 1$ the remainder is $3x + 2$.

Find the values of a and b. [5]

SAT PREP SP www.satprep.co.in

4 Solve the equation

$$\frac{5z}{1+2\mathrm{i}} - zz^* + 30 + 10\mathrm{i} = 0,$$

giving your answers in the form $x + \mathrm{i}y$, where x and y are real. [5]

SAT PREP SP www.satprep.co.in

5 The parametric equations of a curve are

$$x = te^{2t}, \qquad y = t^2 + t + 3.$$

(a) Show that $\frac{dy}{dx} = e^{-2t}$. [3]

SAT PREP SP www.satprep.co.in

(b) Hence show that the normal to the curve, where $t = -1$, passes through the point $\left(0,\ 3 - \frac{1}{e^4}\right)$. [3]

SAT PREP SP www.satprep.co.in

6 **(a)** Express $5\sin\theta + 12\cos\theta$ in the form $R\cos(\theta - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. [3]

SAT PREP SP www.satprep.co.uk

(b) Hence solve the equation $5 \sin 2x + 12 \cos 2x = 6$ for $0 \leqslant x \leqslant \pi$. [4]

SAT PREP SP www.satprep.co.in

7



The diagram shows a circle with centre O and radius r. The angle of the **minor** sector AOB of the circle is x radians. The area of the **major** sector of the circle is 3 times the area of the shaded region.

(a) Show that $x = \frac{3}{4}\sin x + \frac{1}{2}\pi$. [4]

SAT PREP www.satprep.co.uk

(b) Show by calculation that the root of the equation in **(a)** lies between 2 and 2.5. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in **(a)** to calculate this root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

8



The diagram shows the curve $y = x^3 \ln x$, for $x > 0$, and its minimum point M.

(a) Find the exact coordinates of M. [4]

SAT PREP SP www.satprep.co.in

(b) Find the exact area of the shaded region bounded by the curve, the x-axis and the line $x = \frac{1}{2}$. [5]

SAT PREP SP www.satprep.co.uk

9 The variables x and y satisfy the differential equation

$$\frac{dy}{dx} = e^{3y} \sin^2 2x.$$

It is given that $y = 0$ when $x = 0$.

Solve the differential equation and find the value of y when $x = \frac{1}{2}$. [7]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

10 With respect to the origin O, the points A, B, C and D have position vectors given by

$$\overrightarrow{OA} = \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 1 \\ 2 \\ -3 \end{pmatrix}, \quad \overrightarrow{OC} = \begin{pmatrix} 1 \\ -2 \\ 5 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OD} = \begin{pmatrix} 5 \\ -6 \\ 11 \end{pmatrix}.$$

(a) Find the obtuse angle between the vectors $\overrightarrow{OA}$ and $\overrightarrow{OB}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The line l passes through the points A and B.

(b) Find a vector equation for the line l. [2]

.....

.....

.....

.....

.....

.....

© UCLES 2023

(c) Find the position vector of the point of intersection of the line l and the line passing through C and D. [4]

11 Let $\mathrm{f}(x) = \dfrac{5x^2 + x + 11}{(4 + x^2)(1 + x)}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence show that $\int_0^2 \mathrm{f}(x)\,\mathrm{d}x = \ln 54 - \frac{1}{8}\pi$. [5]

© UCLES 2023

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2023

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER			

1087265794

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **October/November 2022**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2022

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 **(a)** Sketch the graph of $y = |2x + 1|$. [1]

(b) Solve the inequality $3x + 5 < |2x + 1|$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

2 On a sketch of an Argand diagram shade the region whose points represent complex numbers z satisfying the inequalities $|z| \leqslant 3$, $\mathrm{Re}\, z \geqslant -2$ and $\frac{1}{4}\pi \leqslant \arg z \leqslant \pi$. [4]

SAT PREP SP www.satprep.co.in

3 Solve the equation $2^{3x-1} = 5(3^{-x})$. Give your answer in the form $\frac{\ln a}{\ln b}$, where a and b are integers. [4]

SAT PREP SP www.satprep.co.in

4 Solve the equation $\tan(x + 45°) = 2\cot x$ for $0° < x < 180°$. [5]

SAT PREP SP www.satprep.co.in

5 The complex numbers u and w are defined by $u = 2e^{\frac{1}{4}\pi i}$ and $w = 3e^{\frac{1}{3}\pi i}$.

(a) Find $\dfrac{u^2}{w}$, giving your answer in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$. Give the exact values of r and θ. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State the least positive integer n such that both $\text{Im}\, w^n = 0$ and $\text{Re}\, w^n > 0$. [1]

.....

.....

.....

.....

.....

.....

.....

6 **(a)** Prove the identity $\cos 4\theta + 4\cos 2\theta + 3 \equiv 8\cos^4 \theta$. [4]

SAT PREP SP www.satprep.co.in

(b) Hence solve the equation $\cos 4\theta + 4\cos 2\theta = 4$ for $0° \leqslant \theta \leqslant 180°$. [3]

SAT PREP SP www.satprep.co.uk

7 The equation of a curve is $y = \dfrac{x}{\cos^2 x}$, for $0 \leqslant x < \frac{1}{2}\pi$. At the point where $x = a$, the tangent to the curve has gradient equal to 12.

(a) Show that $a = \cos^{-1}\left(\sqrt[3]{\dfrac{\cos a + 2a \sin a}{12}}\right)$. [3]

SAT PREP SP www.satprep.co.in

(b) Verify by calculation that a lies between 0.9 and 1. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in part **(a)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

8 In a certain chemical reaction the amount, x grams, of a substance is increasing. The differential equation satisfied by x and t, the time in seconds since the reaction began, is

$$\frac{\mathrm{d}x}{\mathrm{d}t} = kx\mathrm{e}^{-0.1t},$$

where k is a positive constant. It is given that $x = 20$ at the start of the reaction.

(a) Solve the differential equation, obtaining a relation between x, t and k. [5]

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Given that $x = 40$ when $t = 10$, find the value of k and find the value approached by x as t becomes large. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

9



The diagram shows part of the curve $y = (3 - x)e^{-\frac{1}{3}x}$ for $x \geqslant 0$, and its minimum point M.

(a) Find the exact coordinates of M. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

(b) Find the area of the shaded region bounded by the curve and the axes, giving your answer in terms of e. [5]

10 Let $\mathrm{f}(x) = \dfrac{2x^2 + 7x + 8}{(1 + x)(2 + x)^2}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

© UCLES 2022

11



In the diagram, $OABCD$ is a solid figure in which $OA = OB = 4$ units and $OD = 3$ units. The edge OD is vertical, DC is parallel to OB and $DC = 1$ unit. The base, OAB, is horizontal and angle $AOB = 90°$. Unit vectors **i**, **j** and **k** are parallel to OA, OB and OD respectively. The midpoint of AB is M and the point N on BC is such that $CN = 2NB$.

(a) Express vectors $\overrightarrow{MD}$ and $\overrightarrow{ON}$ in terms of **i**, **j** and **k**. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

(b) Calculate the angle in degrees between the directions of $\overrightarrow{MD}$ and $\overrightarrow{ON}$. [3]

(c) Show that the length of the perpendicular from M to ON is $\sqrt{\frac{22}{5}}$. [4]

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

6217921484

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **October/November 2022**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2022

1 Solve the equation $2^{3x-1} = 5(3^{1-x})$. Give your answer in the form $\frac{\ln a}{\ln b}$ where a and b are integers. [4]

SAT PREP SP www.satprep.co.in

2 The polynomial $2x^3 - x^2 + a$, where a is a constant, is denoted by $\mathrm{p}(x)$. It is given that $(2x + 3)$ is a factor of $\mathrm{p}(x)$.

(a) Find the value of a. [2]

.....

.....

.....

.....

.....

.....

.....

.....

(b) When a has this value, solve the inequality $\mathrm{p}(x) < 0$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

3 The equation of a curve is $y = \sin x \sin 2x$. The curve has a stationary point in the interval $0 < x < \frac{1}{2}\pi$.

Find the x-coordinate of this point, giving your answer correct to 3 significant figures. [6]

SAT PREP SP www.satprep.co.in

4 **(a)** Express $4\cos x - \sin x$ in the form $R\cos(x + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. State the exact value of R and give α correct to 2 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence solve the equation $4\cos 2x - \sin 2x = 3$ for $0° < x < 180°$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

[Turn over

5 **(a)** Solve the equation $z^2 - 6iz - 12 = 0$, giving the answers in the form $x + iy$, where x and y are real and exact. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) On a sketch of an Argand diagram with origin O, show points A and B representing the roots of the equation in part **(a)**. [1]

SAT PREP SP www.satprep.co.in

(c) Find the exact modulus and argument of each root. [3]

(d) Hence show that the triangle OAB is equilateral. [1]

SAT PREP SP www.satprep.co.in

6 Relative to the origin O, the points A, B and C have position vectors given by

$$\overrightarrow{OA} = \begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 3 \\ 1 \\ 2 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OC} = \begin{pmatrix} 5 \\ 3 \\ -2 \end{pmatrix}.$$

(a) Using a scalar product, find the cosine of angle BAC. [4]

SAT PREP SP www.satprep.co.in

(b) Hence find the area of triangle ABC. Give your answer in a simplified exact form. [4]

SAT PREP SP www.satprep.co.in

7 The variables x and θ satisfy the differential equation

$$x \sin^2 \theta \frac{\mathrm{d}x}{\mathrm{d}\theta} = \tan^2 \theta - 2 \cot \theta,$$

for $0 < \theta < \frac{1}{2}\pi$ and $x > 0$. It is given that $x = 2$ when $\theta = \frac{1}{4}\pi$.

(a) Show that $\frac{\mathrm{d}}{\mathrm{d}\theta}(\cot^2 \theta) = -\frac{2 \cot \theta}{\sin^2 \theta}$.

(You may assume without proof that the derivative of $\cot \theta$ with respect to θ is $-\operatorname{cosec}^2 \theta$.) [1]

.....

.....

.....

.....

.....

.....

.....

.....

(b) Solve the differential equation and find the value of x when $\theta = \frac{1}{6}\pi$. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

SAT PREP SP www.satprep.co.in

8



The diagram shows part of the curve $y = \sin\sqrt{x}$. This part of the curve intersects the x-axis at the point where $x = a$.

(a) State the exact value of a. [1]

.....

.....

.....

(b) Using the substitution $u = \sqrt{x}$, find the exact area of the shaded region in the first quadrant bounded by this part of the curve and the x-axis. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

9



The diagram shows a semicircle with diameter AB, centre O and radius r. The shaded region is the minor segment on the chord AC and its area is one third of the area of the semicircle. The angle CAB is θ radians.

(a) Show that $\theta = \frac{1}{3}(\pi - 1.5 \sin 2\theta)$. [4]

SAT PREP SP www.satprep.co.in

(b) Verify by calculation that $0.5 < \theta < 0.7$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in part **(a)** to determine θ correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP www.satprep.co.in

10 Let $\mathrm{f}(x) = \dfrac{4 - x + x^2}{(1 + x)(2 + x^2)}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Find the exact value of $\int_0^4 \mathrm{f}(x)\,\mathrm{d}x$. Give your answer as a single logarithm. [5]

© UCLES 2022

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

SAT PREP SP www.satprep.co.in

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

9938938839

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **October/November 2022**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2022

1 Solve the equation $\ln(2x - 1) = 2\ln(x + 1) - \ln x$. Give your answer correct to 3 decimal places. [4]

SAT PREP SP www.satprep.co.uk

2 Expand $\sqrt{\dfrac{1+2x}{1-2x}}$ in ascending powers of x, up to and including the term in x^2, simplifying the coefficients. [5]

SAT PREP SP www.satprep.co.in

3 Find the exact value of $\int_0^{\frac{1}{4}\pi} x \sec^2 x \, \mathrm{d}x$. [5]

SJ SAT PREP www.satprep.co.in

4 The parametric equations of a curve are

$$x = 2t - \tan t, \qquad y = \ln(\sin 2t),$$

for $0 < t < \frac{1}{2}\pi$.

Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \cot t$. [5]

SAT PREP SP www.satprep.co.uk

5 **(a)** On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z + 2| \leqslant 2$ and $\text{Im}\, z \geqslant 1$. [4]

(b) Find the greatest value of $\arg z$ for points in the shaded region. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

6 Solve the quadratic equation $(1 - 3\mathrm{i})z^2 - (2 + \mathrm{i})z + \mathrm{i} = 0$, giving your answers in the form $x + \mathrm{i}y$, where x and y are real. [6]

© UCLES 2022

7 **(a)** Show that the equation $\sqrt{5}\sec x + \tan x = 4$ can be expressed as $R\cos(x + \alpha) = \sqrt{5}$, where $R > 0$ and $0° < \alpha < 90°$. Give the exact value of R and the value of α correct to 2 decimal places. [4]

© UCLES 2022

(b) Hence solve the equation $\sqrt{5}\sec 2x + \tan 2x = 4$, for $0° < x < 180°$. [4]

SAT PREP SP www.satprep.co.in

8 The curve with equation $y = \dfrac{x^3}{e^x - 1}$ has a stationary point at $x = p$, where $p > 0$.

(a) Show that $p = 3(1 - e^{-p})$. [3]

SAT PREP SP www.satprep.co.in

(b) Verify by calculation that p lies between 2.5 and 3. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in part **(a)** to determine p correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

[Turn over

9 With respect to the origin O, the position vectors of the points A, B and C are given by

$$\overrightarrow{OA} = \begin{pmatrix} 0 \\ 5 \\ 2 \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OC} = \begin{pmatrix} 4 \\ -3 \\ -2 \end{pmatrix}.$$

The midpoint of AC is M and the point N lies on BC, between B and C, and is such that $BN = 2NC$.

(a) Find the position vectors of M and N. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find a vector equation for the line through M and N. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

(c) Find the position vector of the point Q where the line through M and N intersects the line through A and B. [4]

10 A gardener is filling an ornamental pool with water, using a hose that delivers 30 litres of water per minute. Initially the pool is empty. At time t minutes after filling begins the volume of water in the pool is V litres. The pool has a small leak and loses water at a rate of $0.01V$ litres per minute.

The differential equation satisfied by V and t is of the form $\frac{\mathrm{d}V}{\mathrm{d}t} = a - bV$.

(a) Write down the values of the constants a and b. [1]

.....

.....

.....

.....

(b) Solve the differential equation and find the value of t when $V = 1000$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Obtain an expression for V in terms of t and hence state what happens to V as t becomes large. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

11 Let $\mathrm{f}(x) = \dfrac{5 - x + 6x^2}{(3 - x)(1 + 3x^2)}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Find the exact value of $\int_0^1 \mathrm{f}(x)\,\mathrm{d}x$, simplifying your answer. [5]

SP SAT PREP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

5794103083

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **May/June 2022**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2022

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Solve the equation $2(3^{2x-1}) = 4^{x+1}$, giving your answer correct to 2 decimal places. [4]

SAT PREP SP www.satprep.co.uk

2 **(a)** Expand $(2 - x^2)^{-2}$ in ascending powers of x, up to and including the term in x^4, simplifying the coefficients. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State the set of values of x for which the expansion is valid. [1]

.....

.....

.....

.....

.....

© UCLES 2022

3 Solve the equation $2\cot 2x + 3\cot x = 5$, for $0° < x < 180°$. [6]

SAT PREP SP www.satprep.co.in

4 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{xy}{1+x^2},$$

and $y = 2$ when $x = 0$.

Solve the differential equation, obtaining a simplified expression for y in terms of x. [7]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

5 The polynomial $ax^3 - 10x^2 + bx + 8$, where a and b are constants, is denoted by $\mathrm{p}(x)$. It is given that $(x - 2)$ is a factor of both $\mathrm{p}(x)$ and $\mathrm{p}'(x)$.

(a) Find the values of a and b. [5]

SAT PREP SP www.satprep.co.in

(b) When a and b have these values, factorise $p(x)$ completely. [3]

SAT PREP SP www.satprep.co.uk

6 Let $I = \int_0^3 \frac{27}{(9+x^2)^2}\,\mathrm{d}x$.

(a) Using the substitution $x = 3\tan\theta$, show that $I = \int_0^{\frac{1}{4}\pi} \cos^2\theta\,\mathrm{d}\theta$. [4]

SAT PREP SP www.satprep.co.in

(b) Hence find the exact value of I. [4]

SAT PREP SP www.satprep.co.uk

7 The complex number u is defined by $u = \dfrac{\sqrt{2} - a\sqrt{2}\mathrm{i}}{1 + 2\mathrm{i}}$, where a is a positive integer.

(a) Express u in terms of a, in the form $x + \mathrm{i}y$, where x and y are real and exact. [3]

SAT PREP SP www.satprep.co.in

It is now given that $a = 3$.

(b) Express u in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$, giving the exact values of r and θ. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Using your answer to part **(b)**, find the two square roots of u. Give your answers in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$, giving the exact values of r and θ. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

8 The equation of a curve is $x^3 + y^3 + 2xy + 8 = 0$.

(a) Express $\frac{dy}{dx}$ in terms of x and y. [4]

SAT PREP SP www.satprep.co.in

The tangent to the curve at the point where $x = 0$ and the tangent at the point where $y = 0$ intersect at the acute angle α.

(b) Find the exact value of $\tan \alpha$. [5]

9



In the diagram, $OABCDEFG$ is a cuboid in which $OA = 2$ units, $OC = 4$ units and $OG = 2$ units. Unit vectors **i**, **j** and **k** are parallel to OA, OC and OG respectively. The point M is the midpoint of DF. The point N on AB is such that $AN = 3NB$.

(a) Express the vectors $\overrightarrow{OM}$ and $\overrightarrow{MN}$ in terms of **i**, **j** and **k**. [3]

.....

.....

.....

.....

.....

.....

.....

(b) Find a vector equation for the line through M and N. [2]

.....

.....

.....

.....

.....

.....

© UCLES 2022

(c) Show that the length of the perpendicular from O to the line through M and N is $\sqrt{\frac{53}{6}}$. [4]

SAT PREP SP www.satprep.co.in

10



The curve $y = x\sqrt{\sin x}$ has one stationary point in the interval $0 < x < \pi$, where $x = a$ (see diagram).

(a) Show that $\tan a = -\frac{1}{2}a$. [4]

SAT PREP SP www.satprep.co.in

(b) Verify by calculation that a lies between 2 and 2.5. [2]

.....

.....

.....

.....

.....

.....

.....

(c) Show that if a sequence of values in the interval $0 < x < \pi$ given by the iterative formula $x_{n+1} = \pi - \tan^{-1}\left(\frac{1}{2}x_n\right)$ converges, then it converges to a, the root of the equation in part **(a)**. [2]

.....

.....

.....

.....

.....

.....

.....

(d) Use the iterative formula given in part **(c)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2022

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

0788354854

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **May/June 2022**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2022

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Solve the equation $\ln(e^{2x} + 3) = 2x + \ln 3$, giving your answer correct to 3 decimal places. [4]

SAT PREP SP www.satprep.co.in

2 Solve the equation $3\cos 2\theta = 3\cos\theta + 2$, for $0° \leqslant \theta \leqslant 360°$. [5]

SAT PREP SP www.satprep.co.in

3 The polynomial $ax^3 + x^2 + bx + 3$ is denoted by $p(x)$. It is given that $p(x)$ is divisible by $(2x - 1)$ and that when $p(x)$ is divided by $(x + 2)$ the remainder is 5.

Find the values of a and b. [5]

SAT PREP SP www.satprep.co.in

4 The equation of a curve is $y = \cos^3 x\sqrt{\sin x}$. It is given that the curve has one stationary point in the interval $0 < x < \frac{1}{2}\pi$.

Find the x-coordinate of this stationary point, giving your answer correct to 3 significant figures. [6]

© UCLES 2022

5 **(a)** By sketching a suitable pair of graphs, show that the equation $\ln x = 3x - x^2$ has one real root. [2]

(b) Verify by calculation that the root lies between 2 and 2.8. [2]

.....

.....

.....

.....

.....

.....

.....

(c) Use the iterative formula $x_{n+1} = \sqrt{3x_n - \ln x_n}$ to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

6 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = x\mathrm{e}^{y-x},$$

and $y = 0$ when $x = 0$.

(a) Solve the differential equation, obtaining an expression for y in terms of x. [7]

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the value of y when $x = 1$, giving your answer in the form $a - \ln b$, where a and b are integers. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 The equation of a curve is $x^3 + 3x^2y - y^3 = 3$.

(a) Show that $\frac{dy}{dx} = \frac{x^2 + 2xy}{y^2 - x^2}$. [4]

SAT PREP SP www.satprep.co.in

(b) Find the coordinates of the points on the curve where the tangent is parallel to the x-axis. [5]

SAT PREP SP www.satprep.co.in

8 Let $f(x) = \dfrac{x^2 + 9x}{(3x - 1)(x^2 + 3)}$.

(a) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence find $\int_1^3 \mathrm{f}(x)\,\mathrm{d}x$, giving your answer in a simplified exact form. [5]

SAT PREP SP www.satprep.co.in

9 The lines l and m have vector equations

$$\mathbf{r} = -\mathbf{i} + 3\mathbf{j} + 4\mathbf{k} + \lambda(2\mathbf{i} - \mathbf{j} - \mathbf{k}) \quad \text{and} \quad \mathbf{r} = 5\mathbf{i} + 4\mathbf{j} + 3\mathbf{k} + \mu(a\mathbf{i} + b\mathbf{j} + \mathbf{k})$$

respectively, where a and b are constants.

(a) Given that l and m intersect, show that $2b - a = 4$. [4]

SAT PREP SP www.satprep.co.in

(b) Given also that l and m are perpendicular, find the values of a and b. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) When a and b have these values, find the position vector of the point of intersection of l and m. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

10 The complex number $-1 + \sqrt{7}\mathrm{i}$ is denoted by u. It is given that u is a root of the equation

$$2x^3 + 3x^2 + 14x + k = 0,$$

where k is a real constant.

(a) Find the value of k. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the other two roots of the equation. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

(c) On an Argand diagram, sketch the locus of points representing complex numbers z satisfying the equation $|z - u| = 2$. [2]

(d) Determine the greatest value of $\arg z$ for points on this locus, giving your answer in radians. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

9937725522

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **May/June 2022**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2022

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Find, in terms of a, the set of values of x satisfying the inequality

$$2|3x + a| < |2x + 3a|,$$

where a is a positive constant. [4]

SAT PREP SP www.satprep.co.in

2 Solve the equation $\cos(\theta - 60°) = 3\sin\theta$, for $0° \leqslant \theta \leqslant 360°$. [5]

SAT PREP SP www.satprep.co.in

3 **(a)** Show that the equation $\log_3(2x + 1) = 1 + 2\log_3(x - 1)$ can be written as a quadratic equation in x. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence solve the equation $\log_3(4y + 1) = 1 + 2\log_3(2y - 1)$, giving your answer correct to 2 decimal places. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

[Turn over

4 The curve $y = e^{-4x} \tan x$ has two stationary points in the interval $0 \leqslant x < \frac{1}{2}\pi$.

(a) Obtain an expression for $\frac{dy}{dx}$ and show it can be written in the form $\sec^2 x(a + b \sin 2x)e^{-4x}$, where a and b are constants. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(b) Hence find the exact x-coordinates of the two stationary points. [3]

SAT PREP SP www.satprep.co.uk

5 The complex number 3 – i is denoted by u.

(a) Show, on an Argand diagram with origin O, the points A, B and C representing the complex numbers u, u^* and $u^* - u$ respectively.

State the type of quadrilateral formed by the points O, A, B and C. [3]

.....

.....

.....

(b) Express $\frac{u^*}{u}$ in the form $x + iy$, where x and y are real. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(c) By considering the argument of $\frac{u^*}{u}$, or otherwise, prove that $\tan^{-1}\left(\frac{3}{4}\right) = 2\tan^{-1}\left(\frac{1}{3}\right)$. [2]

SAT PREP SP www.satprep.co.in

6 The parametric equations of a curve are $x = \dfrac{1}{\cos t}$, $y = \ln \tan t$, where $0 < t < \frac{1}{2}\pi$.

(a) Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{\cos t}{\sin^2 t}$. [5]

SAT PREP SP www.satprep.co.in

(b) Find the equation of the tangent to the curve at the point where $y = 0$. [3]

SAT PREP SP www.satprep.co.uk

7 Let $f(x) = \dfrac{5x^2 + 8x - 3}{(x - 2)(2x^2 + 3)}$.

(a) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

8 At time t days after the start of observations, the number of insects in a population is N. The variation in the number of insects is modelled by a differential equation of the form $\frac{\mathrm{d}N}{\mathrm{d}t} = kN^{\frac{3}{2}} \cos 0.02t$, where k is a constant and N is a continuous variable. It is given that when $t = 0$, $N = 100$.

(a) Solve the differential equation, obtaining a relation between N, k and t. [5]

SAT PREP SP www.satprep.co.in

(b) Given also that $N = 625$ when $t = 50$, find the value of k. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Obtain an expression for N in terms of t, and find the greatest value of N predicted by this model. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

9 With respect to the origin O, the point A has position vector given by $\overrightarrow{OA} = \mathbf{i} + 5\mathbf{j} + 6\mathbf{k}$. The line l has vector equation $\mathbf{r} = 4\mathbf{i} + \mathbf{k} + \lambda(-\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$.

(a) Find in degrees the acute angle between the directions of OA and l. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the position vector of the foot of the perpendicular from A to l. [4]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Hence find the position vector of the reflection of A in l. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.uk

10 The constant a is such that $\int_1^a x^2 \ln x \, dx = 4$.

(a) Show that $a = \left(\frac{35}{3 \ln a - 1}\right)^{\frac{1}{3}}$. [5]

SAT PREP SP www.satprep.co.uk

(b) Verify by calculation that a lies between 2.4 and 2.8. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in part **(a)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2022

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

3201784670

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **February/March 2022**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2022

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Solve the inequality $|2x + 3| > 3|x + 2|$. [4]

SAT PREP SP www.satprep.co.in

2 On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z + 2 - 3i| \leqslant 2$ and $\arg z \leqslant \frac{3}{4}\pi$. [4]

SAT PREP SP www.satprep.co.in

3



The variables x and y satisfy the equation $x^n y^2 = C$, where n and C are constants. The graph of $\ln y$ against $\ln x$ is a straight line passing through the points (0.31, 1.21) and (1.06, 0.91), as shown in the diagram.

Find the value of n and find the value of C correct to 2 decimal places. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

4 The parametric equations of a curve are

$$x = 1 - \cos\theta, \qquad y = \cos\theta - \tfrac{1}{4}\cos 2\theta.$$

Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = -2\sin^2\left(\tfrac{1}{2}\theta\right)$. [5]

SAT PREP SP www.satprep.co.uk

5 The angles α and β lie between $0°$ and $180°$ and are such that

$$\tan(\alpha + \beta) = 2 \quad \text{and} \quad \tan\alpha = 3\tan\beta.$$

Find the possible values of α and β. [6]

SAT PREP SP www.satprep.co.in

6 Find the complex numbers w which satisfy the equation $w^2 + 2iw^* = 1$ and are such that $\text{Re}\, w \leqslant 0$. Give your answers in the form $x + iy$, where x and y are real. [6]

SAT PREP SP www.satprep.co.in

7 **(a)** By sketching a suitable pair of graphs, show that the equation $4 - x^2 = \sec \frac{1}{2}x$ has exactly one root in the interval $0 \leqslant x < \pi$. [2]

(b) Verify by calculation that this root lies between 1 and 2. [2]

.....

.....

.....

.....

.....

.....

(c) Use the iterative formula $x_{n+1} = \sqrt{4 - \sec \frac{1}{2}x_n}$ to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

© UCLES 2022

8 **(a)** Find the quotient and remainder when $8x^3 + 4x^2 + 2x + 7$ is divided by $4x^2 + 1$. [3]

SAT PREP SP www.satprep.co.in

(b) Hence find the exact value of $\int_0^{\frac{1}{2}} \frac{8x^3 + 4x^2 + 2x + 7}{4x^2 + 1}\,\mathrm{d}x$. [5]

SAT PREP SP www.satprep.co.in

9 The variables x and y satisfy the differential equation

$$(x + 1)(3x + 1)\frac{dy}{dx} = y,$$

and it is given that $y = 1$ when $x = 1$.

Solve the differential equation and find the exact value of y when $x = 3$, giving your answer in a simplified form. [9]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

10 The points A and B have position vectors $2\mathbf{i} + \mathbf{j} + \mathbf{k}$ and $\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}$ respectively. The line l has vector equation $\mathbf{r} = \mathbf{i} + 2\mathbf{j} - 3\mathbf{k} + \mu(\mathbf{i} - 3\mathbf{j} - 2\mathbf{k})$.

(a) Find a vector equation for the line through A and B. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the acute angle between the directions of AB and l, giving your answer in degrees. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

(c) Show that the line through A and B does not intersect the line l. [4]

SAT PREP SP www.satprep.co.uk

11



The diagram shows the curve $y = \sin x \cos 2x$ for $0 \leqslant x \leqslant \frac{1}{2}\pi$, and its maximum point M.

(a) Find the x-coordinate of M, giving your answer correct to 3 significant figures. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

(b) Using the substitution $u = \cos x$, find the area of the shaded region enclosed by the curve and the x-axis in the first quadrant, giving your answer in a simplified exact form. [5]

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2022

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

0746866051

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **October/November 2021**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages.

© UCLES 2021

1 Solve the equation $4|5^x - 1| = 5^x$, giving your answers correct to 3 decimal places. [4]

SAT PREP SP www.satprep.co.in

2 **(a)** Express $5\sin x - 3\cos x$ in the form $R\sin(x - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. Give the exact value of R and give α correct to 2 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence state the greatest and least possible values of $(5\sin x - 3\cos x)^2$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

3 The curve with equation $y = xe^{1-2x}$ has one stationary point.

(a) Find the coordinates of this point. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Determine whether the stationary point is a maximum or a minimum. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

4 Using the substitution $u = \sqrt{x}$, find the exact value of

$$\int_3^\infty \frac{1}{(x+1)\sqrt{x}}\,\mathrm{d}x. \qquad [6]$$

SAT PREP SP www.satprep.co.in

5 **(a)** Show that the equation

$$\cot 2\theta + \cot \theta = 2$$

can be expressed as a quadratic equation in $\tan \theta$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence solve the equation $\cot 2\theta + \cot \theta = 2$, for $0 < \theta < \pi$, giving your answers correct to 3 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

6 When $(a + bx)\sqrt{1 + 4x}$, where a and b are constants, is expanded in ascending powers of x, the coefficients of x and x^2 are 3 and −6 respectively.

Find the values of a and b. [6]

SAT PREP SP www.satprep.co.in

7 **(a)** Given that $y = \ln(\ln x)$, show that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{x \ln x}.$$ [1]

.....

.....

.....

.....

.....

The variables x and t satisfy the differential equation

$$x \ln x + t\frac{\mathrm{d}x}{\mathrm{d}t} = 0.$$

It is given that $x = \mathrm{e}$ when $t = 2$.

(b) Solve the differential equation obtaining an expression for x in terms of t, simplifying your answer. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Hence state what happens to the value of x as t tends to infinity. [1]

.....

.....

.....

.....

8 The constant a is such that $\int_1^a \frac{\ln x}{\sqrt{x}}\,\mathrm{d}x = 6$.

(a) Show that $a = \exp\left(\frac{1}{\sqrt{a}} + 2\right)$. [5]

[$\exp(x)$ is an alternative notation for e^x.]

SAT PREP SP www.satprep.co.uk

(b) Verify by calculation that a lies between 9 and 11. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in part **(a)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP www.satprep.co.in

9 Two lines l and m have equations $\mathbf{r} = 3\mathbf{i} + 2\mathbf{j} + 5\mathbf{k} + s(4\mathbf{i} - \mathbf{j} + 3\mathbf{k})$ and $\mathbf{r} = \mathbf{i} - \mathbf{j} - 2\mathbf{k} + t(-\mathbf{i} + 2\mathbf{j} + 2\mathbf{k})$ respectively.

(a) Show that l and m are perpendicular. [2]

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show that l and m intersect and state the position vector of the point of intersection. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

(c) Show that the length of the perpendicular from the origin to the line m is $\frac{1}{3}\sqrt{5}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

10 The complex number $1 + 2i$ is denoted by u. The polynomial $2x^3 + ax^2 + 4x + b$, where a and b are real constants, is denoted by $p(x)$. It is given that u is a root of the equation $p(x) = 0$.

(a) Find the values of a and b. [4]

(b) State a second complex root of this equation. [1]

SAT PREP SP www.satprep.co.in

(c) Find the real factors of $\mathrm{p}(x)$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) **(i)** On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - u| \leqslant \sqrt{5}$ and $\arg z \leqslant \frac{1}{4}\pi$. [4]

(ii) Find the least value of $\mathrm{Im}\, z$ for points in the shaded region. Give your answer in an exact form. [1]

.....

.....

.....

.....

© UCLES 2021

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

4325784148

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **October/November 2021**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2021

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Find the value of x for which $3(2^{1-x}) = 7^x$. Give your answer in the form $\frac{\ln a}{\ln b}$, where a and b are integers. [4]

SAT PREP SP www.satprep.co.in

2 Solve the inequality $|3x - a| > 2|x + 2a|$, where a is a positive constant. [4]

SAT PREP SP www.satprep.co.uk

3 **(a)** Given the complex numbers $u = a + \mathrm{i}b$ and $w = c + \mathrm{i}d$, where a, b, c and d are real, prove that $(u + w)^* = u^* + w^*$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

(b) Solve the equation $(z + 2 + \mathrm{i})^* + (2 + \mathrm{i})z = 0$, giving your answer in the form $x + \mathrm{i}y$ where x and y are real. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

[Turn over

4 Express $\dfrac{4x^2 - 13x + 13}{(2x - 1)(x - 3)}$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

5 **(a)** On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 3 - 2i| \leqslant 1$ and $\text{Im}\, z \geqslant 2$. [4]

(b) Find the greatest value of $\arg z$ for points in the shaded region, giving your answer in degrees. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

6 **(a)** Using the expansions of $\sin(3x + 2x)$ and $\sin(3x - 2x)$, show that

$$\tfrac{1}{2}(\sin 5x + \sin x) \equiv \sin 3x \cos 2x. \qquad [3]$$

SAT PREP SP www.satprep.co.in

(b) Hence show that $\int_0^{\frac{1}{4}\pi} \sin 3x \cos 2x \, \mathrm{d}x = \frac{1}{5}(3 - \sqrt{2})$. [3]

7 The variables x and y satisfy the differential equation

$$e^{2x}\frac{dy}{dx} = 4xy^2,$$

and it is given that $y = 1$ when $x = 0$.

Solve the differential equation, obtaining an expression for y in terms of x. [7]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

8 **(a)** By first expanding $\left(\cos^2\theta + \sin^2\theta\right)^2$, show that

$$\cos^4\theta + \sin^4\theta \equiv 1 - \tfrac{1}{2}\sin^2 2\theta.$$ [3]

SAT PREP SP www.satprep.co.in

(b) Hence solve the equation

$$\cos^4\theta + \sin^4\theta = \tfrac{5}{9},$$

for $0° < \theta < 180°$. [4]

SAT PREP SP www.satprep.co.in

9 The equation of a curve is $ye^{2x} - y^2e^x = 2$.

(a) Show that $\frac{dy}{dx} = \frac{2ye^x - y^2}{2y - e^x}$. [4]

SP SAT PREP www.satprep.co.in

(b) Find the exact coordinates of the point on the curve where the tangent is parallel to the y-axis. [4]

10 With respect to the origin O, the position vectors of the points A and B are given by $\overrightarrow{OA} = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$ and $\overrightarrow{OB} = \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix}$.

(a) Find a vector equation for the line l through A and B. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The point C lies on l and is such that $\overrightarrow{AC} = 3\overrightarrow{AB}$.

Find the position vector of C. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(c) Find the possible position vectors of the point P on l such that $OP = \sqrt{14}$. [5]

SAT PREP SP www.satprep.co.uk

11 The equation of a curve is $y = \sqrt{\tan x}$, for $0 \leqslant x < \frac{1}{2}\pi$.

(a) Express $\frac{dy}{dx}$ in terms of $\tan x$, and verify that $\frac{dy}{dx} = 1$ when $x = \frac{1}{4}\pi$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

The value of $\frac{dy}{dx}$ is also 1 at another point on the curve where $x = a$, as shown in the diagram.



(b) Show that $t^3 + t^2 + 3t - 1 = 0$, where $t = \tan a$. [4]

.....

.....

.....

.....

.....

© UCLES 2021

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use the iterative formula

$$a_{n+1} = \tan^{-1}\left(\tfrac{1}{3}(1 - \tan^2 a_n - \tan^3 a_n)\right)$$

to determine a correct to 2 decimal places, giving the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

* 1 4 7 2 6 4 1 3 2 3 *

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **October/November 2021**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2021

1 Find the quotient and remainder when $2x^4 + 1$ is divided by $x^2 - x + 2$. [3]

SAT PREP SP www.satprep.co.in

2 **(a)** Sketch the graph of $y = |2x - 3|$. [1]

(b) Solve the inequality $|2x - 3| < 3x + 2$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

[Turn over

3 Solve the equation $4^{x-2} = 4^x - 4^2$, giving your answer correct to 3 decimal places. [4]

SAT PREP SP www.satprep.co.uk

4 Find the exact value of $\int_{\frac{1}{3}\pi}^{\pi} x \sin \frac{1}{2}x \, \mathrm{d}x$. [5]

SAT PREP SP www.satprep.co.in

5 Solve the equation $\sin\theta = 3\cos 2\theta + 2$, for $0° \leqslant \theta \leqslant 360°$. [5]

SAT PREP SP www.satprep.co.in

6 **(a)** By first expanding $\cos(x - 60°)$, show that the expression

$$2\cos(x - 60°) + \cos x$$

can be written in the form $R\cos(x - \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. Give the exact value of R and the value of α correct to 2 decimal places. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence find the value of x in the interval $0° < x < 360°$ for which $2\cos(x - 60°) + \cos x$ takes its least possible value. [2]

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.uk

7 The equation of a curve is $\ln(x + y) = x - 2y$.

(a) Show that $\dfrac{dy}{dx} = \dfrac{x + y - 1}{2(x + y) + 1}$. [4]

SAT PREP SP www.satprep.co.in

(b) Find the coordinates of the point on the curve where the tangent is parallel to the x-axis. [3]

SAT PREP SP www.satprep.co.uk

8



In the diagram, $OABCD$ is a pyramid with vertex D. The horizontal base $OABC$ is a square of side 4 units. The edge OD is vertical and $OD = 4$ units. The unit vectors $\mathbf{i}$, $\mathbf{j}$ and $\mathbf{k}$ are parallel to OA, OC and OD respectively.

The midpoint of AB is M and the point N on CD is such that $DN = 3NC$.

(a) Find a vector equation for the line through M and N. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

(b) Show that the length of the perpendicular from O to MN is $\frac{1}{3}\sqrt{82}$. [4]

SAT PREP SP www.satprep.co.uk

9 Let $f(x) = \dfrac{1}{(9-x)\sqrt{x}}$.

(a) Find the x-coordinate of the stationary point of the curve with equation $y = f(x)$. [4]

SAT PREP SP www.satprep.co.in

(b) Using the substitution $u = \sqrt{x}$, show that $\int_0^4 \mathrm{f}(x)\,\mathrm{d}x = \frac{1}{3}\ln 5$. [6]

10 A large plantation of area 20 km^2 is becoming infected with a plant disease. At time t years the area infected is x km^2 and the rate of increase of x is proportional to the ratio of the area infected to the area not yet infected.

When $t = 0$, $x = 1$ and $\frac{dx}{dt} = 1$.

(a) Show that x and t satisfy the differential equation

$$\frac{dx}{dt} = \frac{19x}{20 - x}.$$ [2]

.....

.....

.....

.....

.....

.....

.....

(b) Solve the differential equation and show that when $t = 1$ the value of x satisfies the equation $x = e^{0.9+0.05x}$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in part **(b)**, with an initial value of 2, to determine x correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Calculate the value of t at which the entire plantation becomes infected. [1]

.....

.....

.....

.....

© UCLES 2021

11 The complex number $-\sqrt{3} + \mathrm{i}$ is denoted by u.

(a) Express u in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$, giving the exact values of r and θ. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence show that u^6 is real and state its value. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(c) **(i)** On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $0 \leqslant \arg(z - u) \leqslant \frac{1}{4}\pi$ and $\mathrm{Re}\, z \leqslant 2$. [4]

(ii) Find the greatest value of $|z|$ for points in the shaded region. Give your answer correct to 3 significant figures. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

SAT PREP SP www.satprep.co.in

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

3602942828

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **May/June 2021**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2021

1 Solve the inequality $2|3x - 1| < |x + 1|$. [4]

SAT PREP SP www.satprep.co.in

2 Find the real root of the equation $\frac{2e^x + e^{-x}}{2 + e^x} = 3$, giving your answer correct to 3 decimal places. Your working should show clearly that the equation has only one real root. [5]

SAT PREP SP www.satprep.co.in

3 **(a)** Given that $\cos(x - 30°) = 2\sin(x + 30°)$, show that $\tan x = \dfrac{2 - \sqrt{3}}{1 - 2\sqrt{3}}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence solve the equation

$$\cos(x - 30°) = 2\sin(x + 30°),$$

for $0° < x < 360°$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

4 **(a)** Prove that $\dfrac{1-\cos 2\theta}{1+\cos 2\theta} \equiv \tan^2\theta$. [2]

(b) Hence find the exact value of $\displaystyle\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \frac{1-\cos 2\theta}{1+\cos 2\theta}\,\mathrm{d}\theta$. [4]

SAT PREP www.satprep.co.in

5 **(a)** Solve the equation $z^2 - 2piz - q = 0$, where p and q are real constants. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

In an Argand diagram with origin O, the roots of this equation are represented by the distinct points A and B.

(b) Given that A and B lie on the imaginary axis, find a relation between p and q. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

(c) Given instead that triangle OAB is equilateral, express q in terms of p. [3]

SAT PREP SP www.satprep.co.uk

6 The parametric equations of a curve are

$$x = \ln(2 + 3t), \qquad y = \frac{t}{2 + 3t}.$$

(a) Show that the gradient of the curve is always positive. [5]

SAT PREP SP www.satprep.co.in

(b) Find the equation of the tangent to the curve at the point where it intersects the y-axis. [3]

SAT PREP SP www.satprep.co.in

7



The diagram shows the curve $y = \dfrac{\tan^{-1} x}{\sqrt{x}}$ and its maximum point M where $x = a$.

(a) Show that a satisfies the equation

$$a = \tan\left(\frac{2a}{1 + a^2}\right).$$ [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

(b) Verify by calculation that a lies between 1.3 and 1.5. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use an iterative formula based on the equation in part **(a)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP www.satprep.co.in

8 With respect to the origin O, the points A and B have position vectors given by $\overrightarrow{OA} = \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$ and $\overrightarrow{OB} = \begin{pmatrix} 3 \\ 1 \\ -2 \end{pmatrix}$. The line l has equation $\mathbf{r} = \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}$.

(a) Find the acute angle between the directions of AB and l. [4]

SAT PREP SP www.satprep.co.uk

(b) Find the position vector of the point P on l such that $AP = BP$. [5]

SAT PREP SP www.satprep.co.uk

9 The equation of a curve is $y = x^{-\frac{2}{3}} \ln x$ for $x > 0$. The curve has one stationary point.

(a) Find the exact coordinates of the stationary point. [5]

SAT PREP SP www.satprep.co.uk

(b) Show that $\int_1^8 y \, dx = 18 \ln 2 - 9$. [5]

SAT PREP SP www.satprep.co.in

10 The variables x and t satisfy the differential equation $\frac{\mathrm{d}x}{\mathrm{d}t} = x^2(1 + 2x)$, and $x = 1$ when $t = 0$.

Using partial fractions, solve the differential equation, obtaining an expression for t in terms of x. [11]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

4535261400

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **May/June 2021**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2021

1 Solve the inequality $|2x - 1| < 3|x + 1|$. [4]

SAT PREP SP www.satprep.co.in

2 On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z + 1 - i| \leqslant 1$ and $\arg(z - 1) \leqslant \frac{3}{4}\pi$. [4]

SAT PREP SP www.satprep.co.in

3 The variables x and y satisfy the equation $x = A(3^{-y})$, where A is a constant.

(a) Explain why the graph of y against $\ln x$ is a straight line and state the exact value of the gradient of the line. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

It is given that the line intersects the y-axis at the point where $y = 1.3$.

(b) Calculate the value of A, giving your answer correct to 2 decimal places. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

4 Using integration by parts, find the exact value of $\int_0^2 \tan^{-1}\left(\frac{1}{2}x\right) \mathrm{d}x$. [5]

SAT PREP SP www.satprep.co.in

5 The complex number u is given by $u = 10 - 4\sqrt{6}\mathrm{i}$.

Find the two square roots of u, giving your answers in the form $a + \mathrm{i}b$, where a and b are real and exact. [5]

SAT PREP SP www.satprep.co.in

6 **(a)** Prove that $\operatorname{cosec} 2\theta - \cot 2\theta \equiv \tan \theta$. [3]

(b) Hence show that $\int_{\frac{1}{4}\pi}^{\frac{1}{3}\pi} (\operatorname{cosec} 2\theta - \cot 2\theta)\, \mathrm{d}\theta = \frac{1}{2} \ln 2$. [4]

SAT PREP www.satprep.co.uk

7 A curve is such that the gradient at a general point with coordinates (x, y) is proportional to $\frac{y}{\sqrt{x+1}}$. The curve passes through the points with coordinates $(0, 1)$ and $(3, e)$.

By setting up and solving a differential equation, find the equation of the curve, expressing y in terms of x. [7]

SAT PREP SP www.satprep.co.uk

8 The equation of a curve is $y = e^{-5x} \tan^2 x$ for $-\frac{1}{2}\pi < x < \frac{1}{2}\pi$.

Find the x-coordinates of the stationary points of the curve. Give your answers correct to 3 decimal places where appropriate. [8]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

9 Let $f(x) = \dfrac{14 - 3x + 2x^2}{(2 + x)(3 + x^2)}$.

(a) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

10



The diagram shows a trapezium $ABCD$ in which $AD = BC = r$ and $AB = 2r$. The acute angles BAD and ABC are both equal to x radians. Circular arcs of radius r with centres A and B meet at M, the midpoint of AB.

(a) Given that the sum of the areas of the shaded sectors is 90% of the area of the trapezium, show that x satisfies the equation $x = 0.9(2 - \cos x)\sin x$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Verify by calculation that x lies between 0.5 and 0.7. [2]

.....

.....

.....

.....

.....

.....

© UCLES 2021

(c) Show that if a sequence of values in the interval $0 < x < \frac{1}{2}\pi$ given by the iterative formula

$$x_{n+1} = \cos^{-1}\left(2 - \frac{x_n}{0.9 \sin x_n}\right)$$

converges, then it converges to the root of the equation in part **(a)**. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Use this iterative formula to determine x correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

11 With respect to the origin O, the points A and B have position vectors given by $\overrightarrow{OA} = 2\mathbf{i} - \mathbf{j}$ and $\overrightarrow{OB} = \mathbf{j} - 2\mathbf{k}$.

(a) Show that $OA = OB$ and use a scalar product to calculate angle AOB in degrees. [4]

SAT PREP SP www.satprep.co.in

The midpoint of AB is M. The point P on the line through O and M is such that $PA : OA = \sqrt{7} : 1$.

(b) Find the possible position vectors of P. [6]

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

2912457036

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **May/June 2021**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2021

1 Expand $(1 + 3x)^{\frac{2}{3}}$ in ascending powers of x, up to and including the term in x^3, simplifying the coefficients. [4]

SAT PREP SP www.satprep.co.in

2 Solve the equation $4^x = 3 + 4^{-x}$. Give your answer correct to 3 decimal places. [5]

SAT PREP SP www.satprep.co.uk

3 The parametric equations of a curve are

$$x = t + \ln(t + 2), \qquad y = (t - 1)e^{-2t},$$

where $t > -2$.

(a) Express $\frac{dy}{dx}$ in terms of t, simplifying your answer. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the exact y-coordinate of the stationary point of the curve. [2]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

4 Let $\mathrm{f}(x) = \dfrac{15 - 6x}{(1 + 2x)(4 - x)}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence find $\int_1^2 \mathrm{f}(x)\,\mathrm{d}x$, giving your answer in the form $\ln\left(\dfrac{a}{b}\right)$, where a and b are integers. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

5 **(a)** By first expanding $\tan(2\theta + 2\theta)$, show that the equation $\tan 4\theta = \frac{1}{2}\tan\theta$ may be expressed as $\tan^4\theta + 2\tan^2\theta - 7 = 0$. [4]

SAT PREP SP www.satprep.co.in

(b) Hence solve the equation $\tan 4\theta = \frac{1}{2}\tan\theta$, for $0° < \theta < 180°$. [3]

SAT PREP SP www.satprep.co.uk

6 **(a)** By sketching a suitable pair of graphs, show that the equation $\cot \frac{1}{2}x = 1 + e^{-x}$ has exactly one root in the interval $0 < x \leqslant \pi$. [2]

(b) Verify by calculation that this root lies between 1 and 1.5. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(c) Use the iterative formula $x_{n+1} = 2\tan^{-1}\left(\frac{1}{1+e^{-x_n}}\right)$ to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

SAT PREP SP www.satprep.co.in

7



For the curve shown in the diagram, the normal to the curve at the point P with coordinates (x, y) meets the x-axis at N. The point M is the foot of the perpendicular from P to the x-axis.

The curve is such that for all values of x in the interval $0 \leqslant x < \frac{1}{2}\pi$, the area of triangle PMN is equal to $\tan x$.

(a) **(i)** Show that $\dfrac{MN}{y} = \dfrac{\mathrm{d}y}{\mathrm{d}x}$. [1]

.....

.....

.....

.....

.....

(ii) Hence show that x and y satisfy the differential equation $\frac{1}{2}y^2\dfrac{\mathrm{d}y}{\mathrm{d}x} = \tan x$. [2]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

(b) Given that $y = 1$ when $x = 0$, solve this differential equation to find the equation of the curve, expressing y in terms of x. [6]

SAT PREP SP www.satprep.co.in

8



The diagram shows the curve $y = \frac{\ln x}{x^4}$ and its maximum point M.

(a) Find the exact coordinates of M. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

(b) By using integration by parts, show that for all $a > 1$, $\int_1^a \frac{\ln x}{x^4}\,\mathrm{d}x < \frac{1}{9}$. [6]

9 The quadrilateral $ABCD$ is a trapezium in which AB and DC are parallel. With respect to the origin O, the position vectors of A, B and C are given by $\overrightarrow{OA} = -\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$, $\overrightarrow{OB} = \mathbf{i} + 3\mathbf{j} + \mathbf{k}$ and $\overrightarrow{OC} = 2\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$.

(a) Given that $\overrightarrow{DC} = 3\overrightarrow{AB}$, find the position vector of D. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State a vector equation for the line through A and B. [1]

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(c) Find the distance between the parallel sides and hence find the area of the trapezium. [5]

10 **(a)** Verify that $-1 + \sqrt{2}\mathrm{i}$ is a root of the equation $z^4 + 3z^2 + 2z + 12 = 0$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the other roots of this equation. [7]

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER			

8483881097

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **February/March 2021**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2021

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Solve the equation $\ln(x^3 - 3) = 3\ln x - \ln 3$. Give your answer correct to 3 significant figures. [3]

SAT PREP SP www.satprep.co.in

2 The polynomial $ax^3 + 5x^2 - 4x + b$, where a and b are constants, is denoted by $\mathrm{p}(x)$. It is given that $(x + 2)$ is a factor of $\mathrm{p}(x)$ and that when $\mathrm{p}(x)$ is divided by $(x + 1)$ the remainder is 2.

Find the values of a and b. [5]

SAT PREP SP www.satprep.co.in

3 By first expressing the equation $\tan(x + 45°) = 2\cot x + 1$ as a quadratic equation in $\tan x$, solve the equation for $0° < x < 180°$. [6]

© UCLES 2021

[Turn over

4 The variables x and y satisfy the differential equation

$$(1 - \cos x)\frac{\mathrm{d}y}{\mathrm{d}x} = y \sin x.$$

It is given that $y = 4$ when $x = \pi$.

(a) Solve the differential equation, obtaining an expression for y in terms of x. [6]

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Sketch the graph of y against x for $0 < x < 2\pi$. [1]

SAT PREP SP www.satprep.co.in

5 **(a)** Express $\sqrt{7}\sin x + 2\cos x$ in the form $R\sin(x + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. State the exact value of R and give α correct to 2 decimal places. [3]

SAT PREP SP www.satprep.co.in

(b) Hence solve the equation $\sqrt{7}\sin 2\theta + 2\cos 2\theta = 1$, for $0° < \theta < 180°$. [5]

SAT PREP SP www.satprep.co.in

6 Let $\mathrm{f}(x) = \dfrac{5a}{(2x-a)(3a-x)}$, where a is a positive constant.

(a) Express $\mathrm{f}(x)$ in partial fractions. [3]

SAT PREP SP www.satprep.co.in

(b) Hence show that $\int_a^{2a} \mathrm{f}(x)\,\mathrm{d}x = \ln 6$. [4]

SAT PREP SP www.satprep.co.in

7 Two lines have equations $\mathbf{r} = \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix} + s \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix}$ and $\mathbf{r} = \begin{pmatrix} 2 \\ 1 \\ 4 \end{pmatrix} + t \begin{pmatrix} 1 \\ -1 \\ 4 \end{pmatrix}$.

(a) Show that the lines are skew. [5]

SAT PREP SP www.satprep.co.in

(b) Find the acute angle between the directions of the two lines. [3]

SAT PREP SP www.satprep.co.uk

8 The complex numbers u and v are defined by $u = -4 + 2\mathrm{i}$ and $v = 3 + \mathrm{i}$.

(a) Find $\frac{u}{v}$ in the form $x + \mathrm{i}y$, where x and y are real. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence express $\frac{u}{v}$ in the form $re^{\mathrm{i}\theta}$, where r and θ are exact. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.uk

In an Argand diagram, with origin O, the points A, B and C represent the complex numbers u, v and $2u + v$ respectively.

(c) State fully the geometrical relationship between OA and BC. [2]

.....

.....

.....

.....

.....

.....

.....

.....

(d) Prove that angle $AOB = \frac{3}{4}\pi$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

9 Let $f(x) = \dfrac{e^{2x} + 1}{e^{2x} - 1}$, for $x > 0$.

(a) The equation $x = f(x)$ has one root, denoted by a.

Verify by calculation that a lies between 1 and 1.5. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Use an iterative formula based on the equation in part **(a)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

(c) Find $f'(x)$. Hence find the exact value of x for which $f'(x) = -8$. [6]

SAT PREP SP www.satprep.co.uk

10



The diagram shows the curve $y = \sin 2x \cos^2 x$ for $0 \leqslant x \leqslant \frac{1}{2}\pi$, and its maximum point M.

(a) Using the substitution $u = \sin x$, find the exact area of the region bounded by the curve and the x-axis. [5]

SP SAT PREP www.satprep.co.in

(b) Find the exact x-coordinate of M. [6]

SAT PREP SP www.satprep.co.uk

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2021

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

5907089706

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **October/November 2020**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Blank pages are indicated.

© UCLES 2020

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Solve the inequality $2 - 5x > 2|x - 3|$. [4]

SAT PREP SP www.satprep.co.in

2 On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z| \geqslant 2$ and $|z - 1 + \mathrm{i}| \leqslant 1$. [4]

SAT PREP SP www.satprep.co.in

3 The parametric equations of a curve are

$$x = 3 - \cos 2\theta, \quad y = 2\theta + \sin 2\theta,$$

for $0 < \theta < \frac{1}{2}\pi$.

Show that $\frac{\mathrm{d}y}{\mathrm{d}x} = \cot\theta$. [5]

SAT PREP SP www.satprep.co.uk

4 Solve the equation

$$\log_{10}(2x + 1) = 2\log_{10}(x + 1) - 1.$$

Give your answers correct to 3 decimal places. [6]

SAT PREP SP www.satprep.co.in

5 **(a)** By sketching a suitable pair of graphs, show that the equation $\operatorname{cosec} x = 1 + \mathrm{e}^{-\frac{1}{2}x}$ has exactly two roots in the interval $0 < x < \pi$. [2]

.....

.....

(b) The sequence of values given by the iterative formula

$$x_{n+1} = \pi - \sin^{-1}\left(\frac{1}{\mathrm{e}^{-\frac{1}{2}x_n} + 1}\right),$$

with initial value $x_1 = 2$, converges to one of these roots.

Use the formula to determine this root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 **(a)** Express $\sqrt{6}\cos\theta + 3\sin\theta$ in the form $R\cos(\theta - \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. State the exact value of R and give α correct to 2 decimal places. [3]

© UCLES 2020

(b) Hence solve the equation $\sqrt{6}\cos\frac{1}{3}x + 3\sin\frac{1}{3}x = 2.5$, for $0° < x < 360°$. [4]

SAT PREP SP www.satprep.co.uk

7 **(a)** Verify that $-1 + \sqrt{5}\mathrm{i}$ is a root of the equation $2x^3 + x^2 + 6x - 18 = 0$. [3]

SAT PREP SP www.satprep.co.uk

(b) Find the other roots of this equation. [4]

SAT PREP SP www.satprep.co.uk

8 The coordinates (x, y) of a general point of a curve satisfy the differential equation

$$x\frac{\mathrm{d}y}{\mathrm{d}x} = (1 - 2x^2)y,$$

for $x > 0$. It is given that $y = 1$ when $x = 1$.

Solve the differential equation, obtaining an expression for y in terms of x. [6]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

9 Let $f(x) = \dfrac{8 + 5x + 12x^2}{(1 - x)(2 + 3x)^2}$.

(a) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

10



The diagram shows the curve $y = (2 - x)e^{-\frac{1}{2}x}$, and its minimum point M.

(a) Find the exact coordinates of M. [5]

SAT PREP SP www.satprep.co.in

(b) Find the area of the shaded region bounded by the curve and the axes. Give your answer in terms of e. [5]

11 Two lines have equations $\mathbf{r} = \mathbf{i} + 2\mathbf{j} + \mathbf{k} + \lambda(a\mathbf{i} + 2\mathbf{j} - \mathbf{k})$ and $\mathbf{r} = 2\mathbf{i} + \mathbf{j} - \mathbf{k} + \mu(2\mathbf{i} - \mathbf{j} + \mathbf{k})$, where a is a constant.

(a) Given that the two lines intersect, find the value of a and the position vector of the point of intersection. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(b) Given instead that the acute angle between the directions of the two lines is $\cos^{-1}\left(\frac{1}{6}\right)$, find the two possible values of a. [6]

© UCLES 2020

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER			

7687020674

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **October/November 2020**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Blank pages are indicated.

© UCLES 2020

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Solve the equation

$$\ln(1 + e^{-3x}) = 2.$$

Give the answer correct to 3 decimal places. [3]

SAT PREP SP www.satprep.co.in

2 **(a)** Expand $\sqrt[3]{1+6x}$ in ascending powers of x, up to and including the term in x^3, simplifying the coefficients. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State the set of values of x for which the expansion is valid. [1]

.....

.....

.....

.....

.....

© UCLES 2020

3 The variables x and y satisfy the relation $2^y = 3^{1-2x}$.

(a) By taking logarithms, show that the graph of y against x is a straight line. State the exact value of the gradient of this line. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the exact x-coordinate of the point of intersection of this line with the line $y = 3x$. Give your answer in the form $\frac{\ln a}{\ln b}$, where a and b are integers. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

4 **(a)** Show that the equation $\tan(\theta + 60°) = 2\cot\theta$ can be written in the form

$$\tan^2\theta + 3\sqrt{3}\tan\theta - 2 = 0.$$ [3]

SAT PREP SP www.satprep.co.uk

(b) Hence solve the equation $\tan(\theta + 60°) = 2\cot\theta$, for $0° < \theta < 180°$. [3]

SAT PREP SP www.satprep.co.uk

5



The diagram shows the curve with parametric equations

$$x = \tan\theta, \quad y = \cos^2\theta,$$

for $-\frac{1}{2}\pi < \theta < \frac{1}{2}\pi$.

(a) Show that the gradient of the curve at the point with parameter θ is $-2\sin\theta\cos^3\theta$. [3]

SAT PREP SP www.satprep.co.uk

The gradient of the curve has its maximum value at the point P.

(b) Find the exact value of the x-coordinate of P. [4]

SAT PREP SP www.satprep.co.in

6 The complex number u is defined by

$$u = \frac{7 + \mathrm{i}}{1 - \mathrm{i}}.$$

(a) Express u in the form $x + \mathrm{i}y$, where x and y are real. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show on a sketch of an Argand diagram the points A, B and C representing u, $7 + \mathrm{i}$ and $1 - \mathrm{i}$ respectively. [2]

SAT PREP SP www.satprep.co.in

(c) By considering the arguments of $7 + i$ and $1 - i$, show that

$$\tan^{-1}\left(\tfrac{4}{3}\right) = \tan^{-1}\left(\tfrac{1}{7}\right) + \tfrac{1}{4}\pi. \quad [3]$$

SAT PREP SP www.satprep.co.uk

7 The variables x and t satisfy the differential equation

$$e^{3t}\frac{dx}{dt} = \cos^2 2x,$$

for $t \geqslant 0$. It is given that $x = 0$ when $t = 0$.

(a) Solve the differential equation and obtain an expression for x in terms of t. [7]

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State what happens to the value of x when t tends to infinity. [1]

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

8 With respect to the origin O, the position vectors of the points A, B, C and D are given by

$$\overrightarrow{OA} = \begin{pmatrix} 2 \\ 1 \\ 5 \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 4 \\ -1 \\ 1 \end{pmatrix}, \quad \overrightarrow{OC} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OD} = \begin{pmatrix} 3 \\ 2 \\ 3 \end{pmatrix}.$$

(a) Show that $AB = 2CD$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the angle between the directions of $\overrightarrow{AB}$ and $\overrightarrow{CD}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

(c) Show that the line through A and B does not intersect the line through C and D. [4]

SAT PREP SP www.satprep.co.uk

9 Let $\mathrm{f}(x) = \dfrac{7x + 18}{(3x + 2)(x^2 + 4)}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence find the exact value of $\int_0^2 \mathrm{f}(x)\,\mathrm{d}x$. [6]

10



The diagram shows the curve $y = \sqrt{x}\cos x$, for $0 \leqslant x \leqslant \frac{3}{2}\pi$, and its minimum point M, where $x = a$. The shaded region between the curve and the x-axis is denoted by R.

(a) Show that a satisfies the equation $\tan a = \dfrac{1}{2a}$. [3]

.....

.....

.....

.....

.....

.....

(b) The sequence of values given by the iterative formula $a_{n+1} = \pi + \tan^{-1}\left(\dfrac{1}{2a_n}\right)$, with initial value $x_1 = 3$, converges to a.

Use this formula to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

(c) Find the volume of the solid obtained when the region R is rotated completely about the x-axis. Give your answer in terms of π. [6]

© UCLES 2020

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2020

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

2085422030

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **October/November 2020**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Blank pages are indicated.

© UCLES 2020

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Solve the inequality $2 - 5x > 2|x - 3|$. [4]

SAT PREP SP www.satprep.co.in

2 On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z| \geqslant 2$ and $|z - 1 + \mathrm{i}| \leqslant 1$. [4]

SAT PREP SP www.satprep.co.in

3 The parametric equations of a curve are

$$x = 3 - \cos 2\theta, \quad y = 2\theta + \sin 2\theta,$$

for $0 < \theta < \frac{1}{2}\pi$.

Show that $\frac{\mathrm{d}y}{\mathrm{d}x} = \cot\theta$. [5]

SAT PREP SP www.satprep.co.uk

4 Solve the equation

$$\log_{10}(2x + 1) = 2\log_{10}(x + 1) - 1.$$

Give your answers correct to 3 decimal places. [6]

SAT PREP SP www.satprep.co.in

5 **(a)** By sketching a suitable pair of graphs, show that the equation $\operatorname{cosec} x = 1 + \mathrm{e}^{-\frac{1}{2}x}$ has exactly two roots in the interval $0 < x < \pi$. [2]

.....

.....

(b) The sequence of values given by the iterative formula

$$x_{n+1} = \pi - \sin^{-1}\left(\frac{1}{\mathrm{e}^{-\frac{1}{2}x_n} + 1}\right),$$

with initial value $x_1 = 2$, converges to one of these roots.

Use the formula to determine this root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 **(a)** Express $\sqrt{6}\cos\theta + 3\sin\theta$ in the form $R\cos(\theta - \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. State the exact value of R and give α correct to 2 decimal places. [3]

(b) Hence solve the equation $\sqrt{6}\cos\frac{1}{3}x + 3\sin\frac{1}{3}x = 2.5$, for $0° < x < 360°$. [4]

SAT PREP SP www.satprep.co.uk

7 **(a)** Verify that $-1 + \sqrt{5}\mathrm{i}$ is a root of the equation $2x^3 + x^2 + 6x - 18 = 0$. [3]

SAT PREP SP www.satprep.co.uk

(b) Find the other roots of this equation. [4]

SAT PREP SP www.satprep.co.in

8 The coordinates (x, y) of a general point of a curve satisfy the differential equation

$$x\frac{\mathrm{d}y}{\mathrm{d}x} = (1 - 2x^2)y,$$

for $x > 0$. It is given that $y = 1$ when $x = 1$.

Solve the differential equation, obtaining an expression for y in terms of x. [6]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

9 Let $f(x) = \dfrac{8 + 5x + 12x^2}{(1 - x)(2 + 3x)^2}$.

(a) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

10



The diagram shows the curve $y = (2 - x)e^{-\frac{1}{2}x}$, and its minimum point M.

(a) Find the exact coordinates of M. [5]

SAT PREP SP www.satprep.co.in

(b) Find the area of the shaded region bounded by the curve and the axes. Give your answer in terms of e. [5]

11 Two lines have equations $\mathbf{r} = \mathbf{i} + 2\mathbf{j} + \mathbf{k} + \lambda(a\mathbf{i} + 2\mathbf{j} - \mathbf{k})$ and $\mathbf{r} = 2\mathbf{i} + \mathbf{j} - \mathbf{k} + \mu(2\mathbf{i} - \mathbf{j} + \mathbf{k})$, where a is a constant.

(a) Given that the two lines intersect, find the value of a and the position vector of the point of intersection. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(b) Given instead that the acute angle between the directions of the two lines is $\cos^{-1}\left(\frac{1}{6}\right)$, find the two possible values of a. [6]

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

1214647753

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **May/June 2020**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Blank pages are indicated.

© UCLES 2020

1 Find the set of values of x for which $2(3^{1-2x}) < 5^x$. Give your answer in a simplified exact form. [4]

SAT PREP SP www.satprep.co.uk

2 **(a)** Expand $(2 - 3x)^{-2}$ in ascending powers of x, up to and including the term in x^2, simplifying the coefficients. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State the set of values of x for which the expansion is valid. [1]

.....

.....

.....

.....

.....

© UCLES 2020

[Turn over

3 Express the equation $\tan(\theta + 60°) = 2 + \tan(60° - \theta)$ as a quadratic equation in $\tan\theta$, and hence solve the equation for $0° \leqslant \theta \leqslant 180°$. [6]

SAT PREP SP www.satprep.co.in

4 The curve with equation $y = e^{2x}(\sin x + 3\cos x)$ has a stationary point in the interval $0 \leqslant x \leqslant \pi$.

(a) Find the x-coordinate of this point, giving your answer correct to 2 decimal places. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Determine whether the stationary point is a maximum or a minimum. [2]

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

5 **(a)** Find the quotient and remainder when $2x^3 - x^2 + 6x + 3$ is divided by $x^2 + 3$. [3]

SAT PREP SP www.satprep.co.in

(b) Using your answer to part **(a)**, find the exact value of $\int_1^3 \frac{2x^3 - x^2 + 6x + 3}{x^2 + 3}\,\mathrm{d}x$. [5]

SAT PREP SP www.satprep.co.in

6



The diagram shows a circle with centre O and radius r. The tangents to the circle at the points A and B meet at T, and angle AOB is $2x$ radians. The shaded region is bounded by the tangents AT and BT, and by the minor arc AB. The area of the shaded region is equal to the area of the circle.

(a) Show that x satisfies the equation $\tan x = \pi + x$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

(b) This equation has one root in the interval $0 < x < \frac{1}{2}\pi$. Verify by calculation that this root lies between 1 and 1.4. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use the iterative formula

$$x_{n+1} = \tan^{-1}(\pi + x_n)$$

to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

7 Let $f(x) = \dfrac{\cos x}{1 + \sin x}$.

(a) Show that $f'(x) < 0$ for all x in the interval $-\frac{1}{2}\pi < x < \frac{3}{2}\pi$. [4]

SAT PREP SP www.satprep.co.in

(b) Find $\int_{\frac{1}{6}\pi}^{\frac{1}{2}\pi} \mathrm{f}(x)\,\mathrm{d}x$. Give your answer in a simplified exact form. [4]

8 A certain curve is such that its gradient at a point (x, y) is proportional to $\frac{y}{x\sqrt{x}}$. The curve passes through the points with coordinates $(1, 1)$ and $(4, e)$.

(a) By setting up and solving a differential equation, find the equation of the curve, expressing y in terms of x. [8]

SAT PREP SP www.satprep.co.uk

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Describe what happens to y as x tends to infinity. [1]

.....

.....

.....

.....

.....

9 With respect to the origin O, the vertices of a triangle ABC have position vectors

$$\overrightarrow{OA} = 2\mathbf{i} + 5\mathbf{k}, \quad \overrightarrow{OB} = 3\mathbf{i} + 2\mathbf{j} + 3\mathbf{k} \quad \text{and} \quad \overrightarrow{OC} = \mathbf{i} + \mathbf{j} + \mathbf{k}.$$

(a) Using a scalar product, show that angle ABC is a right angle. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show that triangle ABC is isosceles. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(c) Find the exact length of the perpendicular from O to the line through B and C. [4]

SAT PREP SP www.satprep.co.uk

10 **(a)** The complex number u is defined by $u = \dfrac{3\mathrm{i}}{a + 2\mathrm{i}}$, where a is real.

(i) Express u in the Cartesian form $x + \mathrm{i}y$, where x and y are in terms of a. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the exact value of a for which $\arg u^* = \frac{1}{3}\pi$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(b) **(i)** On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 2i| \leqslant |z - 1 - i|$ and $|z - 2 - i| \leqslant 2$. [4]

(ii) Calculate the least value of $\arg z$ for points in this region. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

4124991549

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **May/June 2020**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Blank pages are indicated.

© UCLES 2020

1 Find the quotient and remainder when $6x^4 + x^3 - x^2 + 5x - 6$ is divided by $2x^2 - x + 1$. [3]

SAT PREP SP www.satprep.co.in

2



The variables x and y satisfy the equation $y^2 = Ae^{kx}$, where A and k are constants. The graph of $\ln y$ against x is a straight line passing through the points (1.5, 1.2) and (5.24, 2.7) as shown in the diagram.

Find the values of A and k correct to 2 decimal places. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 Find the exact value of

$$\int_1^4 x^{\frac{3}{2}} \ln x \, \mathrm{d}x. \qquad [5]$$

SAT PREP SP www.satprep.co.in

4 A curve has equation $y = \cos x \sin 2x$.

Find the x-coordinate of the stationary point in the interval $0 < x < \frac{1}{2}\pi$, giving your answer correct to 3 significant figures. [6]

5 **(a)** Express $\sqrt{2}\cos x - \sqrt{5}\sin x$ in the form $R\cos(x + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. Give the exact value of R and the value of α correct to 3 decimal places. [3]

SAT PREP SP www.satprep.co.in

(b) Hence solve the equation $\sqrt{2}\cos 2\theta - \sqrt{5}\sin 2\theta = 1$, for $0° < \theta < 180°$. [4]

SAT PREP SP www.satprep.co.uk

6



The diagram shows the curve $y = \dfrac{x}{1 + 3x^4}$, for $x \geqslant 0$, and its maximum point M.

(a) Find the x-coordinate of M, giving your answer correct to 3 decimal places. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

(b) Using the substitution $u = \sqrt{3}x^2$, find by integration the exact area of the shaded region bounded by the curve, the x-axis and the line $x = 1$. [5]

7 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{y-1}{(x+1)(x+3)}.$$

It is given that $y = 2$ when $x = 0$.

Solve the differential equation, obtaining an expression for y in terms of x. [9]

SAT PREP SP www.satprep.co.uk

SAT PREP SP www.satprep.co.in

8 **(a)** Solve the equation $(1 + 2\mathrm{i})w + \mathrm{i}w^* = 3 + 5\mathrm{i}$. Give your answer in the form $x + \mathrm{i}y$, where x and y are real. [4]

SAT PREP SP www.satprep.co.in

(b) **(i)** On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 2 - 2i| \leqslant 1$ and $\arg(z - 4i) \geqslant -\frac{1}{4}\pi$. [4]

(ii) Find the least value of $\mathrm{Im}\, z$ for points in this region, giving your answer in an exact form. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

9



The diagram shows the curves $y = \cos x$ and $y = \dfrac{k}{1+x}$, where k is a constant, for $0 \leqslant x \leqslant \frac{1}{2}\pi$. The curves touch at the point where $x = p$.

(a) Show that p satisfies the equation $\tan p = \dfrac{1}{1+p}$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Use the iterative formula $p_{n+1} = \tan^{-1}\left(\frac{1}{1+p_n}\right)$ to determine the value of p correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Hence find the value of k correct to 2 decimal places. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

10 With respect to the origin O, the points A and B have position vectors given by $\overrightarrow{OA} = 6\mathbf{i} + 2\mathbf{j}$ and $\overrightarrow{OB} = 2\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$. The midpoint of OA is M. The point N lying on AB, between A and B, is such that $AN = 2NB$.

(a) Find a vector equation for the line through M and N. [5]

SAT PREP SP www.satprep.co.uk

The line through M and N intersects the line through O and B at the point P.

(b) Find the position vector of P. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Calculate angle OPM, giving your answer in degrees. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

1862605169

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **May/June 2020**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Blank pages are indicated.

© UCLES 2020

1 Solve the inequality $|2x - 1| > 3|x + 2|$. [4]

SAT PREP SP www.satprep.co.in

2 Find the exact value of $\int_0^1 (2-x)e^{-2x}\,dx$. [5]

3 **(a)** Show that the equation

$$\ln(1 + e^{-x}) + 2x = 0$$

can be expressed as a quadratic equation in e^x. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Hence solve the equation $\ln(1 + e^{-x}) + 2x = 0$, giving your answer correct to 3 decimal places. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

4 The equation of a curve is $y = x\tan^{-1}\left(\frac{1}{2}x\right)$.

(a) Find $\frac{\mathrm{d}y}{\mathrm{d}x}$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The tangent to the curve at the point where $x = 2$ meets the y-axis at the point with coordinates $(0, p)$.

Find p. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

[Turn over

5 By first expressing the equation

$$\tan\theta\tan(\theta + 45°) = 2\cot 2\theta$$

as a quadratic equation in $\tan\theta$, solve the equation for $0° < \theta < 90°$. [6]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

6 **(a)** By sketching a suitable pair of graphs, show that the equation $x^5 = 2 + x$ has exactly one real root. [2]

.....

.....

(b) Show that if a sequence of values given by the iterative formula

$$x_{n+1} = \frac{4x_n^5 + 2}{5x_n^4 - 1}$$

converges, then it converges to the root of the equation in part **(a)**. [2]

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use the iterative formula with initial value $x_1 = 1.5$ to calculate the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

7 Let $\mathrm{f}(x) = \dfrac{2}{(2x-1)(2x+1)}$.

(a) Express $\mathrm{f}(x)$ in partial fractions. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Using your answer to part **(a)**, show that

$$\big(\mathrm{f}(x)\big)^2 = \frac{1}{(2x-1)^2} - \frac{1}{2x-1} + \frac{1}{2x+1} + \frac{1}{(2x+1)^2}.$$ [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

(c) Hence show that $\int_1^2 \left(\mathrm{f}(x)\right)^2 \mathrm{d}x = \frac{2}{5} + \frac{1}{2}\ln\left(\frac{5}{9}\right)$. [5]

8 Relative to the origin O, the points A, B and D have position vectors given by

$$\overrightarrow{OA} = \mathbf{i} + 2\mathbf{j} + \mathbf{k}, \quad \overrightarrow{OB} = 2\mathbf{i} + 5\mathbf{j} + 3\mathbf{k} \quad \text{and} \quad \overrightarrow{OD} = 3\mathbf{i} + 2\mathbf{k}.$$

A fourth point C is such that $ABCD$ is a parallelogram.

(a) Find the position vector of C and verify that the parallelogram is not a rhombus. [5]

(b) Find angle BAD, giving your answer in degrees. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find the area of the parallelogram correct to 3 significant figures. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

9 **(a)** The complex numbers u and w are such that

$$u - w = 2\mathrm{i} \quad \text{and} \quad uw = 6.$$

Find u and w, giving your answers in the form $x + \mathrm{i}y$, where x and y are real and exact. [5]

SAT PREP SP www.satprep.co.in

(b) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities

$$|z - 2 - 2\mathrm{i}| \leqslant 2, \quad 0 \leqslant \arg z \leqslant \tfrac{1}{4}\pi \quad \text{and} \quad \mathrm{Re}\, z \leqslant 3. \qquad [5]$$

SAT PREP SP www.satprep.co.in

10



A tank containing water is in the form of a hemisphere. The axis is vertical, the lowest point is A and the radius is r, as shown in the diagram. The depth of water at time t is h. At time $t = 0$ the tank is full and the depth of the water is r. At this instant a tap at A is opened and water begins to flow out at a rate proportional to $\sqrt{h}$. The tank becomes empty at time $t = 14$.

The volume of water in the tank is V when the depth is h. It is given that $V = \frac{1}{3}\pi(3rh^2 - h^3)$.

(a) Show that h and t satisfy a differential equation of the form

$$\frac{\mathrm{d}h}{\mathrm{d}t} = -\frac{B}{2rh^{\frac{1}{2}} - h^{\frac{3}{2}}},$$

where B is a positive constant. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

(b) Solve the differential equation and obtain an expression for t in terms of h and r. [8]

SAT PREP SP www.satprep.co.uk

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

2601216071

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **February/March 2020**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Blank pages are indicated.

© UCLES 2020

1 **(a)** Sketch the graph of $y = |x - 2|$. [1]

(b) Solve the inequality $|x - 2| < 3x - 4$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

2 Solve the equation $\ln 3 + \ln(2x + 5) = 2\ln(x + 2)$. Give your answer in a simplified exact form. [4]

SAT PREP SP www.satprep.co.uk

3 **(a)** By sketching a suitable pair of graphs, show that the equation $\sec x = 2 - \frac{1}{2}x$ has exactly one root in the interval $0 \leqslant x < \frac{1}{2}\pi$. [2]

.....

.....

(b) Verify by calculation that this root lies between 0.8 and 1. [2]

.....

.....

.....

.....

.....

(c) Use the iterative formula $x_{n+1} = \cos^{-1}\left(\frac{2}{4 - x_n}\right)$ to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

4 Find $\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} x \sec^2 x \, \mathrm{d}x$. Give your answer in a simplified exact form. [7]

SAT PREP SP www.satprep.co.in

5 **(a)** Show that $\frac{\cos 3x}{\sin x} + \frac{\sin 3x}{\cos x} = 2\cot 2x$. [4]

© UCLES 2020

(b) Hence solve the equation $\frac{\cos 3x}{\sin x} + \frac{\sin 3x}{\cos x} = 4$, for $0 < x < \pi$. [3]

SAT PREP SP www.satprep.co.in

6 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1 + 4y^2}{\mathrm{e}^x}.$$

It is given that $y = 0$ when $x = 1$.

(a) Solve the differential equation, obtaining an expression for y in terms of x. [7]

SAT PREP SP www.satprep.co.uk

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) State what happens to the value of y as x tends to infinity. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

7 The equation of a curve is $x^3 + 3xy^2 - y^3 = 5$.

(a) Show that $\frac{dy}{dx} = \frac{x^2 + y^2}{y^2 - 2xy}$. [4]

(b) Find the coordinates of the points on the curve where the tangent is parallel to the y-axis. [5]

SAT PREP SP www.satprep.co.uk

8



In the diagram, $OABCDEFG$ is a cuboid in which $OA = 2$ units, $OC = 3$ units and $OD = 2$ units. Unit vectors **i**, **j** and **k** are parallel to OA, OC and OD respectively. The point M on AB is such that $MB = 2AM$. The midpoint of FG is N.

(a) Express the vectors $\overrightarrow{OM}$ and $\overrightarrow{MN}$ in terms of **i**, **j** and **k**. [3]

.....

.....

.....

.....

.....

(b) Find a vector equation for the line through M and N. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

(c) Find the position vector of P, the foot of the perpendicular from D to the line through M and N. [4]

9 Let $f(x) = \dfrac{2 + 11x - 10x^2}{(1 + 2x)(1 - 2x)(2 + x)}$.

(a) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

10 (a) The complex numbers v and w satisfy the equations

$$v + \mathrm{i}w = 5 \qquad \text{and} \qquad (1 + 2\mathrm{i})v - w = 3\mathrm{i}.$$

Solve the equations for v and w, giving your answers in the form $x + \mathrm{i}y$, where x and y are real. [6]

SAT PREP SP www.satprep.co.uk

(b) **(i)** On an Argand diagram, sketch the locus of points representing complex numbers z satisfying $|z - 2 - 3i| = 1$. [2]

(ii) Calculate the least value of $\arg z$ for points on this locus. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2020

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge International A Level

Cambridge Assessment International Education
Cambridge International Advanced Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER			

6182489980

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2019**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2019

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Given that $\ln\left(1 + e^{2y}\right) = x$, express y in terms of x. [3]

SAT PREP SP www.satprep.co.in

2 Solve the inequality $|2x - 3| > 4|x + 1|$. [4]

SAT PREP SP www.satprep.co.in

3 The parametric equations of a curve are

$$x = 2t + \sin 2t, \quad y = \ln(1 - \cos 2t).$$

Show that $\frac{\mathrm{d}y}{\mathrm{d}x} = \operatorname{cosec} 2t$. [5]

SAT PREP SP www.satprep.co.uk

4 The number of insects in a population t weeks after the start of observations is denoted by N. The population is decreasing at a rate proportional to $Ne^{-0.02t}$. The variables N and t are treated as continuous, and it is given that when $t = 0$, $N = 1000$ and $\frac{dN}{dt} = -10$.

(i) Show that N and t satisfy the differential equation

$$\frac{dN}{dt} = -0.01e^{-0.02t}N. \quad [1]$$

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Solve the differential equation and find the value of t when $N = 800$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) State what happens to the value of N as t becomes large. [1]

.....

.....

.....

.....

.....

.....

5 The curve with equation $y = e^{-2x} \ln(x - 1)$ has a stationary point when $x = p$.

(i) Show that p satisfies the equation $x = 1 + \exp\left(\frac{1}{2(x-1)}\right)$, where $\exp(x)$ denotes e^x. [3]

SAT PREP SP www.satprep.co.in

(ii) Verify by calculation that p lies between 2.2 and 2.6. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use an iterative formula based on the equation in part **(i)** to determine p correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

[Turn over

6 **(i)** By differentiating $\frac{\cos x}{\sin x}$, show that if $y = \cot x$ then $\frac{\mathrm{d}y}{\mathrm{d}x} = -\operatorname{cosec}^2 x$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Show that $\int_{\frac{1}{4}\pi}^{\frac{1}{2}\pi} x \operatorname{cosec}^2 x \, \mathrm{d}x = \frac{1}{4}(\pi + \ln 4)$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

SAT PREP SP www.satprep.co.in

7 Two lines l and m have equations $\mathbf{r} = a\mathbf{i} + 2\mathbf{j} + 3\mathbf{k} + \lambda(\mathbf{i} - 2\mathbf{j} + 3\mathbf{k})$ and $\mathbf{r} = 2\mathbf{i} + \mathbf{j} + 2\mathbf{k} + \mu(2\mathbf{i} - \mathbf{j} + \mathbf{k})$ respectively, where a is a constant. It is given that the lines intersect.

(i) Find the value of a. [4]

SAT PREP SP www.satprep.co.in

(ii) When a has this value, find the equation of the plane containing l and m. [5]

SAT PREP SP www.satprep.co.uk

8 Let $\mathrm{f}(x) = \dfrac{x^2 + x + 6}{x^2(x + 2)}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence, showing full working, show that the exact value of $\int_1^4 \mathrm{f}(x)\,\mathrm{d}x$ is $\frac{9}{4}$. [5]

9 **(i)** By first expanding $\cos(2x + x)$, show that $\cos 3x \equiv 4\cos^3 x - 3\cos x$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence solve the equation $\cos 3x + 3\cos x + 1 = 0$, for $0 \leqslant x \leqslant \pi$. [2]

.....

.....

.....

.....

.....

© UCLES 2019

(iii) Find the exact value of $\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \cos^3 x \, \mathrm{d}x$. [4]

SAT PREP SP www.satprep.co.in

10 **(a)** The complex number u is given by $u = -3 - (2\sqrt{10})\mathrm{i}$. Showing all necessary working and without using a calculator, find the square roots of u. Give your answers in the form $a + \mathrm{i}b$, where the numbers a and b are real and exact. [5]

SAT PREP SP www.satprep.co.in

(b) On a sketch of an Argand diagram shade the region whose points represent complex numbers z satisfying the inequalities $|z - 3 - \mathrm{i}| \leqslant 3$, $\arg z \geqslant \frac{1}{4}\pi$ and $\mathrm{Im}\, z \geqslant 2$, where $\mathrm{Im}\, z$ denotes the imaginary part of the complex number z. [5]

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2019

Cambridge International A Level

Cambridge Assessment International Education
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

4978410875

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2019**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2019

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Solve the equation $5\ln(4 - 3^x) = 6$. Show all necessary working and give the answer correct to 3 decimal places. [3]

SAT PREP SP www.satprep.co.in

2 The curve with equation $y = \dfrac{e^{-2x}}{1-x^2}$ has a stationary point in the interval $-1 < x < 1$. Find $\dfrac{dy}{dx}$ and hence find the x-coordinate of this stationary point, giving the answer correct to 3 decimal places. [5]

SAT PREP SP www.satprep.co.in

3 The polynomial $x^4 + 3x^3 + ax + b$, where a and b are constants, is denoted by $p(x)$. When $p(x)$ is divided by $x^2 + x - 1$ the remainder is $2x + 3$. Find the values of a and b. [5]

SAT PREP SP www.satprep.co.in

4 **(i)** Express $(\sqrt{6})\sin x + \cos x$ in the form $R\sin(x + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. State the exact value of R and give α correct to 3 decimal places. [3]

© UCLES 2019

(ii) Hence solve the equation $(\sqrt{6})\sin 2\theta + \cos 2\theta = 2$, for $0° < \theta < 180°$. [4]

SAT PREP SP www.satprep.co.uk

5 The equation of a curve is $2x^2y - xy^2 = a^3$, where a is a positive constant. Show that there is only one point on the curve at which the tangent is parallel to the x-axis and find the y-coordinate of this point. [7]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

6 The variables x and θ satisfy the differential equation

$$\sin \tfrac{1}{2}\theta \frac{\mathrm{d}x}{\mathrm{d}\theta} = (x + 2)\cos \tfrac{1}{2}\theta$$

for $0 < \theta < \pi$. It is given that $x = 1$ when $\theta = \frac{1}{3}\pi$. Solve the differential equation and obtain an expression for x in terms of $\cos\theta$. [8]

© UCLES 2019

SAT PREP SP www.satprep.co.in

7 **(a)** Find the complex number z satisfying the equation

$$z + \frac{\mathrm{i}z}{z^*} - 2 = 0,$$

where z^* denotes the complex conjugate of z. Give your answer in the form $x + \mathrm{i}y$, where x and y are real. [5]

SAT PREP SP www.satprep.co.in

(b) **(i)** On a single Argand diagram sketch the loci given by the equations $|z - 2i| = 2$ and $\text{Im}\, z = 3$, where $\text{Im}\, z$ denotes the imaginary part of z. [2]

(ii) In the first quadrant the two loci intersect at the point P. Find the exact argument of the complex number represented by P. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

8 Let $\mathrm{f}(x) = \dfrac{2x^2 + x + 8}{(2x - 1)(x^2 + 2)}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence, showing full working, find $\int_1^5 \mathrm{f}(x)\,\mathrm{d}x$, giving the answer in the form $\ln c$, where c is an integer. [5]

SAT PREP SP www.satprep.co.in

9 It is given that $\int_0^a x \cos \frac{1}{3}x \, \mathrm{d}x = 3$, where the constant a is such that $0 < a < \frac{3}{2}\pi$.

(i) Show that a satisfies the equation

$$a = \frac{4 - 3\cos \frac{1}{3}a}{\sin \frac{1}{3}a}. \qquad [5]$$

SAT PREP SP www.satprep.co.uk

(ii) Verify by calculation that a lies between 2.5 and 3. [2]

(iii) Use an iterative formula based on the equation in part **(i)** to calculate a correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

SAT PREP SP www.satprep.co.in

10 The line l has equation $\mathbf{r} = \mathbf{i} + 3\mathbf{j} - 2\mathbf{k} + \lambda(\mathbf{i} - 2\mathbf{j} + 3\mathbf{k})$. The plane p has equation $2x + y - 3z = 5$.

(i) Find the position vector of the point of intersection of l and p. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Calculate the acute angle between l and p. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(iii) A second plane q is perpendicular to the plane p and contains the line l. Find the equation of q, giving your answer in the form $ax + by + cz = d$. [5]

SP SAT PREP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2019

Cambridge International A Level

Cambridge Assessment International Education
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

4339107679

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2019**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2019

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Solve the inequality $2|x + 2| > |3x - 1|$. [4]

SAT PREP SP www.satprep.co.in

2 The polynomial $6x^3 + ax^2 + bx - 2$, where a and b are constants, is denoted by $\mathrm{p}(x)$. It is given that $(2x + 1)$ is a factor of $\mathrm{p}(x)$ and that when $\mathrm{p}(x)$ is divided by $(x + 2)$ the remainder is -24. Find the values of a and b. [5]

SAT PREP SP www.satprep.co.in

3 Showing all necessary working, solve the equation $\frac{3^{2x} + 3^{-x}}{3^{2x} - 3^{-x}} = 4$. Give your answer correct to 3 decimal places. [4]

SAT PREP SP www.satprep.co.in

4 **(i)** By first expanding $\tan(2x + x)$, show that the equation $\tan 3x = 3\cot x$ can be written in the form $\tan^4 x - 12\tan^2 x + 3 = 0$. [4]

© UCLES 2019

(ii) Hence solve the equation $\tan 3x = 3\cot x$ for $0° < x < 90°$. [3]

SAT PREP SP www.satprep.co.uk

5 **(i)** By sketching a suitable pair of graphs, show that the equation $\ln(x + 2) = 4e^{-x}$ has exactly one real root. [2]

(ii) Show by calculation that this root lies between $x = 1$ and $x = 1.5$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.uk

(iii) Use the iterative formula $x_{n+1} = \ln\left(\frac{4}{\ln(x_n + 2)}\right)$ to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

SAT PREP SP www.satprep.co.in

6 Throughout this question the use of a calculator is not permitted.

The complex number with modulus 1 and argument $\frac{1}{3}\pi$ is denoted by w.

(i) Express w in the form $x + iy$, where x and y are real and exact. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

The complex number $1 + 2i$ is denoted by u. The complex number v is such that $|v| = 2|u|$ and $\arg v = \arg u + \frac{1}{3}\pi$.

(ii) Sketch an Argand diagram showing the points representing u and v. [2]

SAT PREP SP www.satprep.co.in

(iii) Explain why v can be expressed as $2uw$. Hence find v, giving your answer in the form $a + ib$, where a and b are real and exact. [4]

SAT PREP SP www.satprep.co.in

7 The plane m has equation $x + 4y - 8z = 2$. The plane n is parallel to m and passes through the point P with coordinates $(5, 2, -2)$.

(i) Find the equation of n, giving your answer in the form $ax + by + cz = d$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Calculate the perpendicular distance between m and n. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

(iii) The line l lies in the plane n, passes through the point P and is perpendicular to OP, where O is the origin. Find a vector equation for l. [4]

SAT PREP SP www.satprep.co.in

8



The diagram shows the graph of $y = \sec x$ for $0 \leqslant x < \frac{1}{2}\pi$.

(i) Use the trapezium rule with 2 intervals to estimate the value of $\int_0^{1.2} \sec x \, dx$, giving your answer correct to 2 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

(ii) Explain, with reference to the diagram, whether the trapezium rule gives an overestimate or an underestimate of the true value of the integral in part **(i)**. [1]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

(iii) P is the point on the part of the curve $y = \sec x$ for $0 \leqslant x < \frac{1}{2}\pi$ at which the gradient is 2. By first differentiating $\dfrac{1}{\cos x}$, find the x-coordinate of P, giving your answer correct to 3 decimal places. [6]

SAT PREP SP www.satprep.co.in

9 The variables x and t satisfy the differential equation $5\frac{dx}{dt} = (20 - x)(40 - x)$. It is given that $x = 10$ when $t = 0$.

(i) Using partial fractions, solve the differential equation, obtaining an expression for x in terms of t. [9]

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) State what happens to the value of x when t becomes large. [1]

.....

.....

.....

.....

.....

10



The diagram shows the graph of $y = e^{\cos x} \sin^3 x$ for $0 \leqslant x \leqslant \pi$, and its maximum point M. The shaded region R is bounded by the curve and the x-axis.

(i) Find the x-coordinate of M. Show all necessary working and give your answer correct to 2 decimal places. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

(ii) By first using the substitution $u = \cos x$, find the exact value of the area of R. [7]

SAT PREP SP www.satprep.co.uk

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2019

Cambridge International A Level

Cambridge Assessment International Education
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

2969009541

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2019**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **18** printed pages and **2** blank pages.

© UCLES 2019

1 Use the trapezium rule with 3 intervals to estimate the value of

$$\int_0^3 \left|2^x - 4\right| \mathrm{d}x. \qquad [3]$$

SAT PREP SP www.satprep.co.in

2 Showing all necessary working, solve the equation $\ln(2x - 3) = 2\ln x - \ln(x - 1)$. Give your answer correct to 2 decimal places. [4]

SAT PREP SP www.satprep.co.in

3 Find the gradient of the curve $x^3 + 3xy^2 - y^3 = 1$ at the point with coordinates (1, 3). [4]

SAT PREP SP www.satprep.co.in

4 By first expressing the equation $\cot\theta - \cot(\theta + 45°) = 3$ as a quadratic equation in $\tan\theta$, solve the equation for $0° < \theta < 180°$. [6]

SAT PREP SP www.satprep.co.in

5 **(i)** Differentiate $\dfrac{1}{\sin^2\theta}$ with respect to θ. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) The variables x and θ satisfy the differential equation

$$x\tan\theta\frac{\mathrm{d}x}{\mathrm{d}\theta} + \operatorname{cosec}^2\theta = 0,$$

for $0 < \theta < \frac{1}{2}\pi$ and $x > 0$. It is given that $x = 4$ when $\theta = \frac{1}{6}\pi$. Solve the differential equation, obtaining an expression for x in terms of θ. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

SAT PREP SP www.satprep.co.in

6 **(i)** By first expanding $\sin(2x + x)$, show that $\sin 3x \equiv 3\sin x - 4\sin^3 x$. [4]

SAT PREP SP www.satprep.co.uk

(ii) Hence, showing all necessary working, find the exact value of $\int_0^{\frac{1}{3}\pi} \sin^3 x \, \mathrm{d}x$. [4]

SJ SAT PREP www.satprep.co.in

7



The diagram shows the curves $y = 4\cos\frac{1}{2}x$ and $y = \dfrac{1}{4-x}$, for $0 \leqslant x < 4$. When $x = a$, the tangents to the curves are perpendicular.

(i) Show that $a = 4 - \sqrt{(2\sin\frac{1}{2}a)}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

(ii) Verify by calculation that a lies between 2 and 3. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use an iterative formula based on the equation in part **(i)** to determine a correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

[Turn over

8 Let $f(x) = \dfrac{16 - 17x}{(2 + x)(3 - x)^2}$.

(i) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

9



The diagram shows a set of rectangular axes Ox, Oy and Oz, and four points A, B, C and D with position vectors $\overrightarrow{OA} = 3\mathbf{i}$, $\overrightarrow{OB} = 3\mathbf{i} + 4\mathbf{j}$, $\overrightarrow{OC} = \mathbf{i} + 3\mathbf{j}$ and $\overrightarrow{OD} = 2\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}$.

(i) Find the equation of the plane BCD, giving your answer in the form $ax + by + cz = d$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

(ii) Calculate the acute angle between the planes BCD and $OABC$. [4]

SAT PREP SP www.satprep.co.uk

10 Throughout this question the use of a calculator is not permitted.

The complex number $(\sqrt{3}) + \mathrm{i}$ is denoted by u.

(i) Express u in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$, giving the exact values of r and θ. Hence or otherwise state the exact values of the modulus and argument of u^4. [5]

SAT PREP SP www.satprep.co.in

(ii) Verify that u is a root of the equation $z^3 - 8z + 8\sqrt{3} = 0$ and state the other complex root of this equation. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - u| \leqslant 2$ and $\text{Im}\, z \geqslant 2$, where $\text{Im}\, z$ denotes the imaginary part of z. [5]

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge International A Level

Cambridge Assessment International Education
Cambridge International Advanced Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER			

6271108655

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2019**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **18** printed pages and **2** blank pages.

© UCLES 2019

1 Find the coefficient of x^3 in the expansion of $(3 - x)(1 + 3x)^{\frac{1}{3}}$ in ascending powers of x. [4]

SAT PREP SP www.satprep.co.uk

2 Showing all necessary working, solve the equation $9^x = 3^x + 12$. Give your answer correct to 2 decimal places. [4]

SAT PREP SP www.satprep.co.in

3 Showing all necessary working, solve the equation $\cot 2\theta = 2\tan\theta$ for $0° < \theta < 180°$. [5]

SAT PREP SP www.satprep.co.uk

4 Find the exact coordinates of the point on the curve $y = \dfrac{x}{1 + \ln x}$ at which the gradient of the tangent is equal to $\frac{1}{4}$. [7]

SAT PREP SP www.satprep.co.in

5 Throughout this question the use of a calculator is not permitted.

It is given that the complex number $-1 + (\sqrt{3})\mathrm{i}$ is a root of the equation

$$kx^3 + 5x^2 + 10x + 4 = 0,$$

where k is a real constant.

(i) Write down another root of the equation. [1]

.....

.....

.....

.....

.....

(ii) Find the value of k and the third root of the equation. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

SAT PREP SP www.satprep.co.in

6



In the diagram, A is the mid-point of the semicircle with centre O and radius r. A circular arc with centre A meets the semicircle at B and C. The angle OAB is equal to x radians. The area of the shaded region bounded by AB, AC and the arc with centre A is equal to half the area of the semicircle.

(i) Use triangle OAB to show that $AB = 2r\cos x$. [1]

.....

.....

.....

.....

.....

(ii) Hence show that $x = \cos^{-1}\sqrt{\left(\frac{\pi}{16x}\right)}$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

(iii) Verify by calculation that x lies between 1 and 1.5. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iv) Use an iterative formula based on the equation in part **(ii)** to determine x correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

7 The variables x and y satisfy the differential equation $\frac{dy}{dx} = xe^{x+y}$. It is given that $y = 0$ when $x = 0$.

(i) Solve the differential equation, obtaining y in terms of x. [7]

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Explain why x can only take values that are less than 1. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

8 Let $\mathrm{f}(x) = \dfrac{10x + 9}{(2x + 1)(2x + 3)^2}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence show that $\int_0^1 \mathrm{f}(x)\,\mathrm{d}x = \frac{1}{2}\ln\frac{9}{5} + \frac{1}{5}$. [5]

SAT PREP SP www.satprep.co.in

9 The points A and B have position vectors $\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ and $3\mathbf{i} + \mathbf{j} + \mathbf{k}$ respectively. The line l has equation $\mathbf{r} = 2\mathbf{i} + \mathbf{j} + \mathbf{k} + \mu(\mathbf{i} + \mathbf{j} + 2\mathbf{k})$.

(i) Show that l does not intersect the line passing through A and B. [5]

SAT PREP SP www.satprep.co.in

(ii) The plane m is perpendicular to AB and passes through the mid-point of AB. The plane m intersects the line l at the point P. Find the equation of m and the position vector of P. [5]

10



The diagram shows the curve $y = \sin 3x \cos x$ for $0 \leqslant x \leqslant \frac{1}{2}\pi$ and its minimum point M. The shaded region R is bounded by the curve and the x-axis.

(i) By expanding $\sin(3x + x)$ and $\sin(3x - x)$ show that

$$\sin 3x \cos x = \tfrac{1}{2}(\sin 4x + \sin 2x).$$ [3]

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Using the result of part **(i)** and showing all necessary working, find the exact area of the region R. [4]

.....

.....

.....

.....

.....

.....

© UCLES 2019

.....

.....

.....

.....

.....

.....

.....

(iii) Using the result of part **(i)**, express $\frac{dy}{dx}$ in terms of $\cos 2x$ and hence find the x-coordinate of M, giving your answer correct to 2 decimal places. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge International A Level

Cambridge Assessment International Education
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

3275076176

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2019**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **20** printed pages.

© UCLES 2019

1 Use logarithms to solve the equation $5^{3-2x} = 4(7^x)$, giving your answer correct to 3 decimal places. [4]

SAT PREP SP www.satprep.co.in

2 Show that $\int_0^{\frac{1}{4}\pi} x^2 \cos 2x \, dx = \frac{1}{32}(\pi^2 - 8)$. [5]

© UCLES 2019

[Turn over

3 Let $f(\theta) = \dfrac{1 - \cos 2\theta + \sin 2\theta}{1 + \cos 2\theta + \sin 2\theta}$.

(i) Show that $f(\theta) = \tan \theta$. [3]

SAT PREP SP www.satprep.co.in

(ii) Hence show that $\int_{\frac{1}{6}\pi}^{\frac{1}{4}\pi} \mathrm{f}(\theta)\,\mathrm{d}\theta = \frac{1}{2}\ln\frac{3}{2}$. [4]

4 The equation of a curve is $y = \dfrac{1 + e^{-x}}{1 - e^{-x}}$, for $x > 0$.

(i) Show that $\dfrac{dy}{dx}$ is always negative. [3]

SAT PREP SP www.satprep.co.in

(ii) The gradient of the curve is equal to -1 when $x = a$. Show that a satisfies the equation $e^{2a} - 4e^{a} + 1 = 0$. Hence find the exact value of a. [4]

SP SAT PREP www.satprep.co.in

5 The variables x and y satisfy the differential equation

$$(x + 1)y\frac{\mathrm{d}y}{\mathrm{d}x} = y^2 + 5.$$

It is given that $y = 2$ when $x = 0$. Solve the differential equation obtaining an expression for y^2 in terms of x. [7]

SP SAT PREP www.satprep.co.in

6



The diagram shows the curve $y = x^4 - 2x^3 - 7x - 6$. The curve intersects the x-axis at the points $(a, 0)$ and $(b, 0)$, where $a < b$. It is given that b is an integer.

(i) Find the value of b. [1]

.....

.....

.....

.....

.....

.....

.....

(ii) Hence show that a satisfies the equation $a = -\frac{1}{3}(2 + a^2 + a^3)$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use an iterative formula based on the equation in part **(ii)** to determine a correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

[Turn over

7 The curve $y = \sin\left(x + \frac{1}{3}\pi\right)\cos x$ has two stationary points in the interval $0 \leqslant x \leqslant \pi$.

(i) Find $\frac{\mathrm{d}y}{\mathrm{d}x}$. [2]

(ii) By considering the formula for $\cos(A + B)$, show that, at the stationary points on the curve, $\cos\left(2x + \frac{1}{3}\pi\right) = 0$. [2]

SAT PREP SP www.satprep.co.in

(iii) Hence find the exact x-coordinates of the stationary points. [3]

SAT PREP SP www.satprep.co.uk

8 Throughout this question the use of a calculator is not permitted.

The complex number u is defined by

$$u = \frac{4\mathrm{i}}{1 - (\sqrt{3})\mathrm{i}}.$$

(i) Express u in the form $x + \mathrm{i}y$, where x and y are real and exact. [3]

SAT PREP SP www.satprep.co.in

(ii) Find the exact modulus and argument of u. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z| < 2$ and $|z - u| < |z|$. [4]

SAT PREP SP www.satprep.co.in

9 Let $f(x) = \dfrac{2x(5-x)}{(3+x)(1-x)^2}$.

(i) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence obtain the expansion of $f(x)$ in ascending powers of x up to and including the term in x^3. [5]

SAT PREP SP www.satprep.co.in

10 The line l has equation $\mathbf{r} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k} + \mu(2\mathbf{i} - \mathbf{j} - 2\mathbf{k})$.

(i) The point P has position vector $4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$. Find the length of the perpendicular from P to l. [5]

SAT PREP SP www.satprep.co.in

(ii) It is given that l lies in the plane with equation $ax + by + 2z = 13$, where a and b are constants. Find the values of a and b. [6]

© UCLES 2019

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2019

Cambridge International A Level

Cambridge Assessment International Education
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

4755885761

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **February/March 2019**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **17** printed pages and **3** blank pages.

© UCLES 2019

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 **(i)** Show that the equation $\log_{10}(x-4) = 2 - \log_{10} x$ can be written as a quadratic equation in x. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence solve the equation $\log_{10}(x-4) = 2 - \log_{10} x$, giving your answer correct to 3 significant figures. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

2 The sequence of values given by the iterative formula

$$x_{n+1} = \frac{2x_n^6 + 12x_n}{3x_n^5 + 8},$$

with initial value $x_1 = 2$, converges to α.

(i) Use the formula to calculate α correct to 4 decimal places. Give the result of each iteration to 6 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) State an equation satisfied by α and hence find the exact value of α. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

3 **(i)** Given that $\sin(\theta + 45°) + 2\cos(\theta + 60°) = 3\cos\theta$, find the exact value of $\tan\theta$ in a form involving surds. You need not simplify your answer. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence solve the equation $\sin(\theta + 45°) + 2\cos(\theta + 60°) = 3\cos\theta$ for $0° < \theta < 360°$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

4 Show that $\int_1^4 x^{-\frac{3}{2}} \ln x \, \mathrm{d}x = 2 - \ln 4$. [5]

SAT PREP SP www.satprep.co.in

5 The variables x and y satisfy the relation $\sin y = \tan x$, where $-\frac{1}{2}\pi < y < \frac{1}{2}\pi$. Show that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{\cos x \sqrt{(\cos 2x)}}.$$ [5]

SAT PREP SP www.satprep.co.in

6 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = ky^3\mathrm{e}^{-x},$$

where k is a constant. It is given that $y = 1$ when $x = 0$, and that $y = \sqrt{\mathrm{e}}$ when $x = 1$. Solve the differential equation, obtaining an expression for y in terms of x. [7]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

7 **(a)** Showing all working and without using a calculator, solve the equation

$$(1 + \mathrm{i})z^2 - (4 + 3\mathrm{i})z + 5 + \mathrm{i} = 0.$$

Give your answers in the form $x + \mathrm{i}y$, where x and y are real. [6]

SAT PREP SP www.satprep.co.in

(b) The complex number u is given by

$$u = -1 - \mathrm{i}.$$

On a sketch of an Argand diagram show the point representing u. Shade the region whose points represent complex numbers satisfying the inequalities $|z| < |z - 2\mathrm{i}|$ and $\frac{1}{4}\pi < \arg(z - u) < \frac{1}{2}\pi$. [4]

SAT PREP SP www.satprep.co.in

8 Let $f(x) = \dfrac{12 + 12x - 4x^2}{(2 + x)(3 - 2x)}$.

(i) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

9 Two planes have equations $2x + 3y - z = 1$ and $x - 2y + z = 3$.

(i) Find the acute angle between the planes. [4]

SAT PREP SP www.satprep.co.in

(ii) Find a vector equation for the line of intersection of the planes. [6]

SAT PREP SP www.satprep.co.in

10



The diagram shows the curve $y = \sin^3 x \sqrt{(\cos x)}$ for $0 \leqslant x \leqslant \frac{1}{2}\pi$, and its maximum point M.

(i) Using the substitution $u = \cos x$, find by integration the exact area of the shaded region bounded by the curve and the x-axis. [6]

SAT PREP SP www.satprep.co.uk

(ii) Showing all your working, find the x-coordinate of M, giving your answer correct to 3 decimal places. [6]

© UCLES 2019

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

3283715323

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2018**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **20** printed pages.

© UCLES 2018

1 Find the set of values of x satisfying the inequality $2|2x - a| < |x + 3a|$, where a is a positive constant. [4]

© UCLES 2018

2 Showing all necessary working, solve the equation $\frac{2e^x + e^{-x}}{e^x - e^{-x}} = 4$, giving your answer correct to 2 decimal places. [4]

SAT PREP SP www.satprep.co.in

3 **(i)** By sketching a suitable pair of graphs, show that the equation $x^3 = 3 - x$ has exactly one real root. [2]

(ii) Show that if a sequence of real values given by the iterative formula

$$x_{n+1} = \frac{2x_n^3 + 3}{3x_n^2 + 1}$$

converges, then it converges to the root of the equation in part **(i)**. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use this iterative formula to determine the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

[Turn over

4 The parametric equations of a curve are

$$x = 2\sin\theta + \sin 2\theta, \quad y = 2\cos\theta + \cos 2\theta,$$

where $0 < \theta < \pi$.

(i) Obtain an expression for $\frac{\mathrm{d}y}{\mathrm{d}x}$ in terms of θ. [3]

SAT PREP SP www.satprep.co.in

(ii) Hence find the exact coordinates of the point on the curve at which the tangent is parallel to the y-axis. [4]

5 The coordinates (x, y) of a general point on a curve satisfy the differential equation

$$x\frac{\mathrm{d}y}{\mathrm{d}x} = (2 - x^2)y.$$

The curve passes through the point $(1, 1)$. Find the equation of the curve, obtaining an expression for y in terms of x. [7]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

6 **(i)** Show that the equation $(\sqrt{2})\operatorname{cosec} x + \cot x = \sqrt{3}$ can be expressed in the form $R\sin(x - \alpha) = \sqrt{2}$, where $R > 0$ and $0° < \alpha < 90°$. [4]

(ii) Hence solve the equation $(\sqrt{2})\operatorname{cosec} x + \cot x = \sqrt{3}$, for $0° < x < 180°$. [4]

SAT PREP SP www.satprep.co.uk

7



The diagram shows the curve $y = 5\sin^2 x \cos^3 x$ for $0 \leqslant x \leqslant \frac{1}{2}\pi$, and its maximum point M. The shaded region R is bounded by the curve and the x-axis.

(i) Find the x-coordinate of M, giving your answer correct to 3 decimal places. [5]

SAT PREP SP www.satprep.co.in

(ii) Using the substitution $u = \sin x$ and showing all necessary working, find the exact area of R. [4]

SAT PREP SP www.satprep.co.in

8 **(a)** Showing all necessary working, express the complex number $\dfrac{2+3\mathrm{i}}{1-2\mathrm{i}}$ in the form $r\mathrm{e}^{\mathrm{i}\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$. Give the values of r and θ correct to 3 significant figures. [5]

SAT PREP SP www.satprep.co.in

(b) On an Argand diagram sketch the locus of points representing complex numbers z satisfying the equation $|z - 3 + 2i| = 1$. Find the least value of $|z|$ for points on this locus, giving your answer in an exact form. [4]

SAT PREP SP www.satprep.co.uk

9 Let $\mathrm{f}(x) = \dfrac{6x^2 + 8x + 9}{(2 - x)(3 + 2x)^2}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence, showing all necessary working, show that $\int_{-1}^{0} \mathrm{f}(x)\,\mathrm{d}x = 1 + \frac{1}{2}\ln\left(\frac{3}{4}\right)$. [5]

SAT PREP SP www.satprep.co.in

10 The planes m and n have equations $3x + y - 2z = 10$ and $x - 2y + 2z = 5$ respectively. The line l has equation $\mathbf{r} = 4\mathbf{i} + 2\mathbf{j} + \mathbf{k} + \lambda(\mathbf{i} + \mathbf{j} + 2\mathbf{k})$.

(i) Show that l is parallel to m. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Calculate the acute angle between the planes m and n. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

(iii) A point P lies on the line l. The perpendicular distance of P from the plane n is equal to 2. Find the position vectors of the two possible positions of P. [4]

SP SAT PREP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

9782733059

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2018**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2018

1 Solve the inequality $3|2x - 1| > |x + 4|$. [4]

SAT PREP SP www.satprep.co.in

2 Showing all necessary working, solve the equation $\sin(\theta - 30°) + \cos\theta = 2\sin\theta$, for $0° < \theta < 180°$. [4]

3 **(i)** Find $\int \frac{\ln x}{x^3}\,\mathrm{d}x$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence show that $\int_1^2 \frac{\ln x}{x^3}\,\mathrm{d}x = \frac{1}{16}(3 - \ln 4)$. [2]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

4 Showing all necessary working, solve the equation

$$\frac{e^x + e^{-x}}{e^x + 1} = 4,$$

giving your answer correct to 3 decimal places. [5]

SAT PREP SP www.satprep.co.in

5 The equation of a curve is $y = x\ln(8 - x)$. The gradient of the curve is equal to 1 at only one point, when $x = a$.

(i) Show that a satisfies the equation $x = 8 - \dfrac{8}{\ln(8 - x)}$. [3]

SAT PREP SP www.satprep.co.in

(ii) Verify by calculation that a lies between 2.9 and 3.1. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use an iterative formula based on the equation in part **(i)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP www.satprep.co.in

6 A certain curve is such that its gradient at a general point with coordinates (x, y) is proportional to $\frac{y^2}{x}$. The curve passes through the points with coordinates (1, 1) and (e, 2). By setting up and solving a differential equation, find the equation of the curve, expressing y in terms of x. [8]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

7 A curve has equation $y = \dfrac{3\cos x}{2 + \sin x}$, for $-\frac{1}{2}\pi \leqslant x \leqslant \frac{1}{2}\pi$.

(i) Find the exact coordinates of the stationary point of the curve. [6]

SAT PREP SP www.satprep.co.in

(ii) The constant a is such that $\int_0^a \frac{3\cos x}{2+\sin x}\,\mathrm{d}x = 1$. Find the value of a, giving your answer correct to 3 significant figures. [4]

SAT PREP SP www.satprep.co.uk

8 Let $f(x) = \dfrac{7x^2 - 15x + 8}{(1 - 2x)(2 - x)^2}$.

(i) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

9 **(a)** **(i)** Without using a calculator, express the complex number $\frac{2+6\mathrm{i}}{1-2\mathrm{i}}$ in the form $x+\mathrm{i}y$, where x and y are real. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence, without using a calculator, express $\frac{2+6\mathrm{i}}{1-2\mathrm{i}}$ in the form $r(\cos\theta+\mathrm{i}\sin\theta)$, where $r>0$ and $-\pi<\theta\leqslant\pi$, giving the exact values of r and θ. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

(b) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying both the inequalities $|z - 3i| \leqslant 1$ and $\mathrm{Re}\, z \leqslant 0$, where $\mathrm{Re}\, z$ denotes the real part of z. Find the greatest value of $\arg z$ for points in this region, giving your answer in radians correct to 2 decimal places. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

[Turn over

10 The line l has equation $\mathbf{r} = 5\mathbf{i} - 3\mathbf{j} - \mathbf{k} + \lambda(\mathbf{i} - 2\mathbf{j} + \mathbf{k})$. The plane p has equation

$$(\mathbf{r} - \mathbf{i} - 2\mathbf{j}).(3\mathbf{i} + \mathbf{j} + \mathbf{k}) = 0.$$

The line l intersects the plane p at the point A.

(i) Find the position vector of A. [3]

SAT PREP SP www.satprep.co.in

(ii) Calculate the acute angle between l and p. [4]

[Question 10 (iii) is printed on the next page.]

(iii) Find the equation of the line which lies in p and intersects l at right angles. [4]

SAT PREP SP www.satprep.co.uk

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

3426712677

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2018**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **20** printed pages.

© UCLES 2018

1 Find the set of values of x satisfying the inequality $2|2x - a| < |x + 3a|$, where a is a positive constant. [4]

© UCLES 2018

2 Showing all necessary working, solve the equation $\frac{2e^{x} + e^{-x}}{e^{x} - e^{-x}} = 4$, giving your answer correct to 2 decimal places. [4]

SAT PREP SP www.satprep.co.in

3 **(i)** By sketching a suitable pair of graphs, show that the equation $x^3 = 3 - x$ has exactly one real root. [2]

(ii) Show that if a sequence of real values given by the iterative formula

$$x_{n+1} = \frac{2x_n^3 + 3}{3x_n^2 + 1}$$

converges, then it converges to the root of the equation in part **(i)**. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use this iterative formula to determine the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

4 The parametric equations of a curve are

$$x = 2\sin\theta + \sin 2\theta, \quad y = 2\cos\theta + \cos 2\theta,$$

where $0 < \theta < \pi$.

(i) Obtain an expression for $\frac{\mathrm{d}y}{\mathrm{d}x}$ in terms of θ. [3]

SAT PREP SP www.satprep.co.in

(ii) Hence find the exact coordinates of the point on the curve at which the tangent is parallel to the y-axis. [4]

5 The coordinates (x, y) of a general point on a curve satisfy the differential equation

$$x\frac{\mathrm{d}y}{\mathrm{d}x} = (2 - x^2)y.$$

The curve passes through the point $(1, 1)$. Find the equation of the curve, obtaining an expression for y in terms of x. [7]

SAT PREP SP www.satprep.co.in

6 **(i)** Show that the equation $(\sqrt{2})\operatorname{cosec} x + \cot x = \sqrt{3}$ can be expressed in the form $R\sin(x - \alpha) = \sqrt{2}$, where $R > 0$ and $0° < \alpha < 90°$. [4]

(ii) Hence solve the equation $(\sqrt{2})\operatorname{cosec} x + \cot x = \sqrt{3}$, for $0° < x < 180°$. [4]

SAT PREP SP www.satprep.co.in

7



The diagram shows the curve $y = 5\sin^2 x \cos^3 x$ for $0 \leqslant x \leqslant \frac{1}{2}\pi$, and its maximum point M. The shaded region R is bounded by the curve and the x-axis.

(i) Find the x-coordinate of M, giving your answer correct to 3 decimal places. [5]

SAT PREP SP www.satprep.co.in

(ii) Using the substitution $u = \sin x$ and showing all necessary working, find the exact area of R. [4]

SAT PREP SP www.satprep.co.in

8 **(a)** Showing all necessary working, express the complex number $\dfrac{2+3\mathrm{i}}{1-2\mathrm{i}}$ in the form $r\mathrm{e}^{\mathrm{i}\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$. Give the values of r and θ correct to 3 significant figures. [5]

SAT PREP SP www.satprep.co.in

(b) On an Argand diagram sketch the locus of points representing complex numbers z satisfying the equation $|z - 3 + 2i| = 1$. Find the least value of $|z|$ for points on this locus, giving your answer in an exact form. [4]

SAT PREP SP www.satprep.co.uk

9 Let $\mathrm{f}(x) = \dfrac{6x^2 + 8x + 9}{(2 - x)(3 + 2x)^2}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence, showing all necessary working, show that $\int_{-1}^{0} \mathrm{f}(x)\,\mathrm{d}x = 1 + \frac{1}{2}\ln\left(\frac{3}{4}\right)$. [5]

SAT PREP SP www.satprep.co.in

10 The planes m and n have equations $3x + y - 2z = 10$ and $x - 2y + 2z = 5$ respectively. The line l has equation $\mathbf{r} = 4\mathbf{i} + 2\mathbf{j} + \mathbf{k} + \lambda(\mathbf{i} + \mathbf{j} + 2\mathbf{k})$.

(i) Show that l is parallel to m. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Calculate the acute angle between the planes m and n. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

(iii) A point P lies on the line l. The perpendicular distance of P from the plane n is equal to 2. Find the position vectors of the two possible positions of P. [4]

SP SAT PREP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

7084725728

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2018**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **18** printed pages and **2** blank pages.

© UCLES 2018

1 Showing all necessary working, solve the equation $\ln(x^4 - 4) = 4 \ln x - \ln 4$, giving your answer correct to 2 decimal places. [4]

© UCLES 2018

2 **(i)** Given that $\sin(x - 60°) = 3\cos(x - 45°)$, find the exact value of $\tan x$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence solve the equation $\sin(x - 60°) = 3\cos(x - 45°)$, for $0° < x < 360°$. [2]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

[Turn over

3 A curve has equation $y = \dfrac{e^{3x}}{\tan \frac{1}{2}x}$. Find the x-coordinates of the stationary points of the curve in the interval $0 < x < \pi$. Give your answers correct to 3 decimal places. [6]

SAT PREP SP www.satprep.co.uk

4 The polynomial $x^4 + 2x^3 + ax + b$, where a and b are constants, is divisible by $x^2 - x + 1$. Find the values of a and b. [5]

SAT PREP SP www.satprep.co.in

5 Let $I = \int_{\frac{1}{4}}^{\frac{3}{4}} \sqrt{\left(\frac{x}{1-x}\right)}\,\mathrm{d}x$.

(i) Using the substitution $x = \cos^2\theta$, show that $I = \int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} 2\cos^2\theta\,\mathrm{d}\theta$. [4]

SAT PREP SP www.satprep.co.in

(ii) Hence find the exact value of I. [4]

SAT PREP SP www.satprep.co.uk

6 In a certain chemical reaction the amount, x grams, of a substance is decreasing. The differential equation relating x and t, the time in seconds since the reaction started, is

$$\frac{\mathrm{d}x}{\mathrm{d}t} = -kx\sqrt{t},$$

where k is a positive constant. It is given that $x = 100$ at the start of the reaction.

(i) Solve the differential equation, obtaining a relation between x, t and k. [5]

SAT PREP SP www.satprep.co.in

(ii) Given that $t = 25$ when $x = 80$, find the value of t when $x = 40$. [3]

SAT PREP SP www.satprep.co.uk

7 **(i)** Showing all working and without using a calculator, solve the equation $z^2 + (2\sqrt{6})z + 8 = 0$, giving your answers in the form $x + iy$, where x and y are real and exact. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Sketch an Argand diagram showing the points representing the roots. [1]

SAT PREP SP www.satprep.co.in

(iii) The points representing the roots are A and B, and O is the origin. Find angle AOB. [3]

(iv) Prove that triangle AOB is equilateral. [1]

SAT PREP SP www.satprep.co.in

8 The positive constant a is such that $\int_0^a xe^{-\frac{1}{2}x}\,dx = 2$.

(i) Show that a satisfies the equation $a = 2\ln(a+2)$. [5]

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Verify by calculation that a lies between 3 and 3.5. [2]

.....

.....

.....

.....

.....

.....

(iii) Use an iteration based on the equation in part **(i)** to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

9 Let $f(x) = \dfrac{12x^2 + 4x - 1}{(x-1)(3x+2)}$.

(i) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

10 The point P has position vector $3\mathbf{i} - 2\mathbf{j} + \mathbf{k}$. The line l has equation $\mathbf{r} = 4\mathbf{i} + 2\mathbf{j} + 5\mathbf{k} + \mu(\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$.

(i) Find the length of the perpendicular from P to l, giving your answer correct to 3 significant figures. [5]

SAT PREP SP www.satprep.co.in

(ii) Find the equation of the plane containing l and P, giving your answer in the form $ax + by + cz = d$. [5]

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

BLANK PAGE

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	
CANDIDATE NUMBER	

9448820043

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2018**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **20** printed pages.

© UCLES 2018

CAMBRIDGE International Examinations

1 Showing all necessary working, solve the equation $3|2^x - 1| = 2^x$, giving your answers correct to 3 significant figures. [4]

© UCLES 2018

2 Showing all necessary working, solve the equation $\cot\theta + \cot(\theta + 45°) = 2$, for $0° < \theta < 180°$. [5]

SAT PREP SP www.satprep.co.uk

3



In the diagram, the tangent to a curve at the point P with coordinates (x, y) meets the x-axis at T. The point N is the foot of the perpendicular from P to the x-axis. The curve is such that, for all values of x, the gradient of the curve is positive and $TN = 2$.

(i) Show that the differential equation satisfied by x and y is $\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{2}y$. [1]

.....

.....

.....

.....

.....

.....

.....

The point with coordinates $(4, 3)$ lies on the curve.

(ii) Solve the differential equation to obtain the equation of the curve, expressing y in terms of x. [5]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

SAT PREP SP www.satprep.co.in

4 **(i)** Show that $\dfrac{2\sin x - \sin 2x}{1 - \cos 2x} \equiv \dfrac{\sin x}{1 + \cos x}$. [4]

SAT PREP SP www.satprep.co.in

(ii) Hence, showing all necessary working, find $\int_{\frac{1}{3}\pi}^{\frac{1}{2}\pi} \frac{2\sin x - \sin 2x}{1 - \cos 2x}\,\mathrm{d}x$, giving your answer in the form $\ln k$. [4]

SAT PREP SP www.satprep.co.in

5 The equation of a curve is $x^2(x + 3y) - y^3 = 3$.

(i) Show that $\frac{dy}{dx} = \frac{x^2 + 2xy}{y^2 - x^2}$. [4]

SJ PREP www.sajprep.co.in

(ii) Hence find the exact coordinates of the two points on the curve at which the gradient of the normal is 1. [4]

6



The diagram shows a triangle ABC in which $AB = AC = a$ and angle $BAC = \theta$ radians. Semicircles are drawn outside the triangle with AB and AC as diameters. A circular arc with centre A joins B and C. The area of the shaded segment is equal to the sum of the areas of the semicircles.

(i) Show that $\theta = \frac{1}{2}\pi + \sin\theta$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

(ii) Verify by calculation that θ lies between 2.2 and 2.4. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use an iterative formula based on the equation in part **(i)** to determine θ correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP www.satprep.co.in

7 Throughout this question the use of a calculator is not permitted.

The complex numbers $-3\sqrt{3} + \mathrm{i}$ and $\sqrt{3} + 2\mathrm{i}$ are denoted by u and v respectively.

(i) Find, in the form $x + \mathrm{i}y$, where x and y are real and exact, the complex numbers uv and $\frac{u}{v}$. [5]

SAT PREP SP www.satprep.co.in

(ii) On a sketch of an Argand diagram with origin O, show the points A and B representing the complex numbers u and v respectively. Prove that angle $AOB = \frac{2}{3}\pi$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

8



The diagram shows the curve $y = (x + 1)e^{-\frac{1}{3}x}$ and its maximum point M.

(i) Find the x-coordinate of M. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

(ii) Find the area of the shaded region enclosed by the curve and the axes, giving your answer in terms of e. [5]

9 Let $\mathrm{f}(x) = \dfrac{x - 4x^2}{(3 - x)(2 + x^2)}$.

(i) Express $\mathrm{f}(x)$ in the form $\dfrac{A}{3 - x} + \dfrac{Bx + C}{2 + x^2}$. [4]

SAT PREP SP www.satprep.co.in

(ii) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^3. [5]

© UCLES 2018

[Turn over

10 Two lines l and m have equations $\mathbf{r} = 2\mathbf{i} - \mathbf{j} + \mathbf{k} + s(2\mathbf{i} + 3\mathbf{j} - \mathbf{k})$ and $\mathbf{r} = \mathbf{i} + 3\mathbf{j} + 4\mathbf{k} + t(\mathbf{i} + 2\mathbf{j} + \mathbf{k})$ respectively.

(i) Show that the lines are skew. [4]

A plane p is parallel to the lines l and m.

(ii) Find a vector that is normal to p. [3]

© UCLES 2018

.....

.....

.....

.....

.....

.....

(iii) Given that p is equidistant from the lines l and m, find the equation of p. Give your answer in the form $ax + by + cz = d$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

6101943949

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2018**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2018

BLANK PAGE

SAT PREP SP www.satprep.co.in

1 Expand $\frac{4}{\sqrt{(4-3x)}}$ in ascending powers of x, up to and including the term in x^2, simplifying the coefficients. [4]

SAT PREP SP www.satprep.co.in

2 Showing all necessary working, solve the equation $5^{2x} = 5^x + 5$. Give your answer correct to 3 decimal places. [5]

SAT PREP SP www.satprep.co.in

3 Showing all necessary working, find the value of $\int_0^{\frac{1}{6}\pi} x\cos 3x\,\mathrm{d}x$, giving your answer in terms of π. [5]

SAT PREP SP www.satprep.co.in

4 The curve with equation $y = \dfrac{\ln x}{3 + x}$ has a stationary point at $x = p$.

(i) Show that p satisfies the equation $\ln x = 1 + \dfrac{3}{x}$. [3]

SAT PREP SP www.satprep.co.in

(ii) By sketching suitable graphs, show that the equation in part **(i)** has only one root. [2]

(iii) It is given that the equation in part **(i)** can be written in the form $x = \dfrac{3+x}{\ln x}$. Use an iterative formula based on this rearrangement to determine the value of p correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

5 **(i)** By first expanding $\left(\cos^2 x + \sin^2 x\right)^3$, or otherwise, show that

$$\cos^6 x + \sin^6 x = 1 - \tfrac{3}{4}\sin^2 2x.$$ [4]

SAT PREP SP www.satprep.co.in

(ii) Hence solve the equation

$$\cos^6 x + \sin^6 x = \tfrac{2}{3},$$

for $0° < x < 180°$. [4]

SAT PREP SP www.satprep.co.in

6 **(i)** Express $\frac{1}{4-y^2}$ in partial fractions. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) The variables x and y satisfy the differential equation

$$x\frac{dy}{dx} = 4 - y^2,$$

and $y = 1$ when $x = 1$. Solve the differential equation, obtaining an expression for y in terms of x. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

SAT PREP SP www.satprep.co.in

7 Throughout this question the use of a calculator is not permitted.

(i) Express $\cos\theta + 2\sin\theta$ in the form $R\cos(\theta - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. Give the exact values of R and $\tan\alpha$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence, showing all necessary working, show that $\int_0^{\frac{1}{4}\pi} \frac{15}{(\cos\theta + 2\sin\theta)^2}\,\mathrm{d}\theta = 5$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

SAT PREP SP www.satprep.co.in

8 The equation of a curve is $2x^3 - y^3 - 3xy^2 = 2a^3$, where a is a non-zero constant.

(i) Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{2x^2 - y^2}{y^2 + 2xy}$. [4]

SP SAT PREP www.satprep.co.in

(ii) Find the coordinates of the two points on the curve at which the tangent is parallel to the y-axis. [5]

© UCLES 2018

[Turn over

9 **(a)** Find the complex number z satisfying the equation

$$3z - \mathrm{i}z^* = 1 + 5\mathrm{i},$$

where z^* denotes the complex conjugate of z. [4]

SAT PREP SP www.satprep.co.in

(b) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z which satisfy both the inequalities $|z| \leqslant 3$ and $\mathrm{Im}\, z \geqslant 2$, where $\mathrm{Im}\, z$ denotes the imaginary part of z. Calculate the greatest value of $\arg z$ for points in this region. Give your answer in radians correct to 2 decimal places. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

[Turn over

10 The points A and B have position vectors $2\mathbf{i} + \mathbf{j} + 3\mathbf{k}$ and $4\mathbf{i} + \mathbf{j} + \mathbf{k}$ respectively. The line l has equation $\mathbf{r} = 4\mathbf{i} + 6\mathbf{j} + \mu(\mathbf{i} + 2\mathbf{j} - 2\mathbf{k})$.

(i) Show that l does not intersect the line passing through A and B. [5]

SAT PREP SP www.satprep.co.in

The point P, with parameter t, lies on l and is such that angle PAB is equal to 120°.

(ii) Show that $3t^2 + 8t + 4 = 0$. Hence find the position vector of P. [6]

SAT PREP SP www.satprep.co.uk

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

5197459152

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **February/March 2018**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **20** printed pages.

© UCLES 2018

1 Use the trapezium rule with three intervals to estimate the value of

$$\int_0^{\frac{1}{4}\pi} \sqrt{(1 - \tan x)}\, \mathrm{d}x,$$

giving your answer correct to 3 decimal places. [3]

SAT PREP SP www.satprep.co.uk

2 Expand $\sqrt[4]{(1 - 4x)}$ in ascending powers of x, up to and including the term in x^3, simplifying the coefficients. [4]

SAT PREP SP www.satprep.co.in

3 **(i)** Using the expansions of $\cos(3x + x)$ and $\cos(3x - x)$, show that

$$\tfrac{1}{2}(\cos 4x + \cos 2x) \equiv \cos 3x \cos x. \qquad [3]$$

SAT PREP SP www.satprep.co.in

(ii) Hence show that $\int_{-\frac{1}{6}\pi}^{\frac{1}{6}\pi} \cos 3x \cos x \, \mathrm{d}x = \frac{3}{8}\sqrt{3}$. [3]

SAT PREP SP www.satprep.co.in

4 The variables x and y satisfy the equation $y^n = Ax^3$, where n and A are constants. It is given that $y = 2.58$ when $x = 1.20$, and $y = 9.49$ when $x = 2.51$.

(i) Explain why the graph of $\ln y$ against $\ln x$ is a straight line. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the values of n and A, giving your answers correct to 2 decimal places. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

SAT PREP SP www.satprep.co.in

5 The parametric equations of a curve are

$$x = 2t + \sin 2t, \quad y = 1 - 2\cos 2t,$$

for $-\frac{1}{2}\pi < t < \frac{1}{2}\pi$.

(i) Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = 2\tan t$. [5]

SAT PREP SP www.satprep.co.uk

(ii) Hence find the x-coordinate of the point on the curve at which the gradient of the normal is 2. Give your answer correct to 3 significant figures. [2]

SP SAT PREP www.satprep.co.in

6 The variables x and θ satisfy the differential equation

$$x\cos^2\theta\frac{\mathrm{d}x}{\mathrm{d}\theta} = 2\tan\theta + 1,$$

for $0 \leqslant \theta < \frac{1}{2}\pi$ and $x > 0$. It is given that $x = 1$ when $\theta = \frac{1}{4}\pi$.

(i) Show that $\dfrac{\mathrm{d}}{\mathrm{d}\theta}(\tan^2\theta) = \dfrac{2\tan\theta}{\cos^2\theta}$. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Solve the differential equation and calculate the value of x when $\theta = \frac{1}{3}\pi$, giving your answer correct to 3 significant figures. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

SAT PREP SP www.satprep.co.in

7 **(i)** By sketching suitable graphs, show that the equation $e^{2x} = 6 + e^{-x}$ has exactly one real root. [2]

(ii) Verify by calculation that this root lies between 0.5 and 1. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(iii) Show that if a sequence of values given by the iterative formula

$$x_{n+1} = \tfrac{1}{3}\ln\left(1 + 6\mathrm{e}^{x_n}\right)$$

converges, then it converges to the root of the equation in part **(i)**. [2]

(iv) Use this iterative formula to calculate the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

© UCLES 2018

8 Let $\mathrm{f}(x) = \dfrac{5x^2 + x + 27}{(2x + 1)(x^2 + 9)}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence find $\int_0^4 \mathrm{f}(x)\,\mathrm{d}x$, giving your answer in the form $\ln c$, where c is an integer. [5]

SAT PREP SP www.satprep.co.in

9 The complex number $1 + 2i$ is denoted by u.

(i) It is given that u is a root of the equation $2x^3 - x^2 + 4x + k = 0$, where k is a constant.

(a) Showing all working and without using a calculator, find the value of k. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Showing all working and without using a calculator, find the other two roots of this equation. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2018

.....

.....

.....

.....

.....

.....

.....

(ii) On an Argand diagram sketch the locus of points representing complex numbers z satisfying the equation $|z - u| = 1$. Determine the least value of $\arg z$ for points on this locus. Give your answer in radians correct to 2 decimal places. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SATPREP www.satprep.co.in

10 The line l has equation $\mathbf{r} = 4\mathbf{i} + 3\mathbf{j} - \mathbf{k} + \mu(\mathbf{i} + 2\mathbf{j} - 2\mathbf{k})$. The plane p has equation $2x - 3y - z = 4$.

(i) Find the position vector of the point of intersection of l and p. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the acute angle between l and p. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

(iii) A second plane q is parallel to l, perpendicular to p and contains the point with position vector $4\mathbf{j} - \mathbf{k}$. Find the equation of q, giving your answer in the form $ax + by + cz = d$. [5]

© UCLES 2018

[Turn over

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME

CENTRE NUMBER | | | | | |

CANDIDATE NUMBER | | | | |

8927569197

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2017**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2017

CAMBRIDGE International Examinations

1 Find the quotient and remainder when x^4 is divided by $x^2 + 2x - 1$. [3]

SAT PREP SP www.satprep.co.uk

2 Two variable quantities x and y are believed to satisfy an equation of the form $y = C(a^x)$, where C and a are constants. An experiment produced four pairs of values of x and y. The table below gives the corresponding values of x and $\ln y$.

x	0.9	1.6	2.4	3.2
$\ln y$	1.7	1.9	2.3	2.6

By plotting $\ln y$ against x for these four pairs of values and drawing a suitable straight line, estimate the values of C and a. Give your answers correct to 2 significant figures. [5]



.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 The equation $x^3 = 3x + 7$ has one real root, denoted by α.

(i) Show by calculation that α lies between 2 and 3. [2]

SAT PREP SP www.satprep.co.in

Two iterative formulae, A and B, derived from this equation are as follows:

$$x_{n+1} = (3x_n + 7)^{\frac{1}{3}}, \qquad (A)$$

$$x_{n+1} = \frac{x_n^3 - 7}{3}. \qquad (B)$$

Each formula is used with initial value $x_1 = 2.5$.

(ii) Show that one of these formulae produces a sequence which fails to converge, and use the other formula to calculate α correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [4]

SAT PREP SP www.satprep.co.in

4 **(i)** Prove the identity $\tan(45° + x) + \tan(45° - x) \equiv 2\sec 2x$. [4]

SAT PREP SP www.satprep.co.in

(ii) Hence sketch the graph of $y = \tan(45° + x) + \tan(45° - x)$ for $0° \leqslant x \leqslant 90°$. [3]

SAT PREP SP www.satprep.co.in

5 The equation of a curve is $2x^4 + xy^3 + y^4 = 10$.

(i) Show that $\frac{dy}{dx} = -\frac{8x^3 + y^3}{3xy^2 + 4y^3}$. [4]

© UCLES 2017

(ii) Hence show that there are two points on the curve at which the tangent is parallel to the x-axis and find the coordinates of these points. [4]

SAT PREP SP www.satprep.co.in

6 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 4\cos^2 y \tan x,$$

for $0 \leqslant x < \frac{1}{2}\pi$, and $x = 0$ when $y = \frac{1}{4}\pi$. Solve this differential equation and find the value of x when $y = \frac{1}{3}\pi$. [8]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

7 **(a)** The complex number u is given by $u = 8 - 15\mathrm{i}$. Showing all necessary working, find the two square roots of u. Give answers in the form $a + \mathrm{i}b$, where the numbers a and b are real and exact. [5]

SAT PREP SP www.satprep.co.in

(b) On an Argand diagram, shade the region whose points represent complex numbers satisfying both the inequalities $|z - 2 - \mathrm{i}| \leqslant 2$ and $0 \leqslant \arg(z - \mathrm{i}) \leqslant \frac{1}{4}\pi$. [4]

SAT PREP SP www.satprep.co.in

8 Let $\mathrm{f}(x) = \dfrac{4x^2 + 9x - 8}{(x+2)(2x-1)}$.

(i) Express $\mathrm{f}(x)$ in the form $A + \dfrac{B}{x+2} + \dfrac{C}{2x-1}$. [4]

SAT PREP SP www.satprep.co.in

(ii) Hence show that $\int_1^4 \mathrm{f}(x)\,\mathrm{d}x = 6 + \frac{1}{2}\ln\left(\frac{16}{7}\right)$. [5]

SAT PREP SP www.satprep.co.in

9



The diagram shows the curve $y = (1 + x^2)e^{-\frac{1}{2}x}$ for $x \geqslant 0$. The shaded region R is enclosed by the curve, the x-axis and the lines $x = 0$ and $x = 2$.

(i) Find the exact values of the x-coordinates of the stationary points of the curve. [4]

SAT PREP SP www.satprep.co.uk

(ii) Show that the exact value of the area of R is $18 - \frac{42}{e}$. [5]

SAT PREP SP www.satprep.co.in

10 The equations of two lines l and m are $\mathbf{r} = 3\mathbf{i} - \mathbf{j} - 2\mathbf{k} + \lambda(-\mathbf{i} + \mathbf{j} + 4\mathbf{k})$ and $\mathbf{r} = 4\mathbf{i} + 4\mathbf{j} - 3\mathbf{k} + \mu(2\mathbf{i} + \mathbf{j} - 2\mathbf{k})$ respectively.

(i) Show that the lines do not intersect. [3]

(ii) Calculate the acute angle between the directions of the lines. [3]

SAT PREP SP www.satprep.co.in

(iii) Find the equation of the plane which passes through the point $(3, -2, -1)$ and which is parallel to both l and m. Give your answer in the form $ax + by + cz = d$. [5]

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

5462375726

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2017**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2017

1



The diagram shows a sketch of the curve $y = \dfrac{3}{\sqrt{(9 - x^3)}}$ for values of x from -1.2 to 1.2.

(i) Use the trapezium rule, with two intervals, to estimate the value of

$$\int_{-1.2}^{1.2} \frac{3}{\sqrt{(9 - x^3)}}\,\mathrm{d}x,$$

giving your answer correct to 2 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

(ii) Explain, with reference to the diagram, why the trapezium rule may be expected to give a good approximation to the true value of the integral in this case. [1]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

2 Showing all necessary working, solve the equation $2\log_2 x = 3 + \log_2(x + 1)$, giving your answer correct to 3 significant figures. [5]

© UCLES 2017

[Turn over

3 By expressing the equation $\tan(\theta + 60°) + \tan(\theta - 60°) = \cot\theta$ in terms of $\tan\theta$ only, solve the equation for $0° < \theta < 90°$. [5]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

4 The curve with equation $y = \dfrac{2 - \sin x}{\cos x}$ has one stationary point in the interval $-\frac{1}{2}\pi < x < \frac{1}{2}\pi$.

(i) Find the exact coordinates of this point. [5]

SAT PREP SP www.satprep.co.in

(ii) Determine whether this point is a maximum or a minimum point. [2]

SAT PREP SP www.satprep.co.uk

5 The variables x and y satisfy the differential equation

$$(x + 1)\frac{dy}{dx} = y(x + 2),$$

and it is given that $y = 2$ when $x = 1$. Solve the differential equation and obtain an expression for y in terms of x. [7]

SAT PREP SP www.satprep.co.in

6 The equation of a curve is $x^3y - 3xy^3 = 2a^4$, where a is a non-zero constant.

(i) Show that $\frac{dy}{dx} = \frac{3x^2y - 3y^3}{9xy^2 - x^3}$. [4]

(ii) Hence show that there are only two points on the curve at which the tangent is parallel to the x-axis and find the coordinates of these points. [4]

SAT PREP SP www.satprep.co.in

7 Throughout this question the use of a calculator is not permitted.

The complex number $1 - (\sqrt{3})\mathrm{i}$ is denoted by u.

(i) Find the modulus and argument of u. [2]

(ii) Show that $u^3 + 8 = 0$. [2]

SAT PREP SP www.satprep.co.in

(iii) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying both the inequalities $|z - u| \leqslant 2$ and $\mathrm{Re}\, z \geqslant 2$, where $\mathrm{Re}\, z$ denotes the real part of z. [4]

SAT PREP SP www.satprep.co.in

8 Let $f(x) = \dfrac{8x^2 + 9x + 8}{(1 - x)(2x + 3)^2}$.

(i) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

9 It is given that $\int_1^a x^{\frac{1}{2}} \ln x \, \mathrm{d}x = 2$, where $a > 1$.

(i) Show that $a^{\frac{3}{2}} = \dfrac{7 + 2a^{\frac{3}{2}}}{3 \ln a}$. [5]

SAT PREP SP www.satprep.co.in

(ii) Show by calculation that a lies between 2 and 4. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use the iterative formula

$$a_{n+1} = \left(\frac{7 + 2a_n^{\frac{3}{2}}}{3 \ln a_n} \right)^{\frac{2}{3}}$$

to determine a correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10 Two planes p and q have equations $x + y + 3z = 8$ and $2x - 2y + z = 3$ respectively.

(i) Calculate the acute angle between the planes p and q. [4]

(ii) The point A on the line of intersection of p and q has y-coordinate equal to 2. Find the equation of the plane which contains the point A and is perpendicular to both the planes p and q. Give your answer in the form $ax + by + cz = d$. [7]

SAT PREP SP www.sprep.co.in

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

9388870911

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2017**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2017

CAMBRIDGE International Examinations

1 Find the quotient and remainder when x^4 is divided by $x^2 + 2x - 1$. [3]

SAT PREP SP www.satprep.co.uk

2 Two variable quantities x and y are believed to satisfy an equation of the form $y = C(a^x)$, where C and a are constants. An experiment produced four pairs of values of x and y. The table below gives the corresponding values of x and $\ln y$.

x	0.9	1.6	2.4	3.2
$\ln y$	1.7	1.9	2.3	2.6

By plotting $\ln y$ against x for these four pairs of values and drawing a suitable straight line, estimate the values of C and a. Give your answers correct to 2 significant figures. [5]



.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 The equation $x^3 = 3x + 7$ has one real root, denoted by α.

(i) Show by calculation that α lies between 2 and 3. [2]

SAT PREP SP www.satprep.co.in

Two iterative formulae, A and B, derived from this equation are as follows:

$$x_{n+1} = (3x_n + 7)^{\frac{1}{3}}, \qquad (A)$$

$$x_{n+1} = \frac{x_n^3 - 7}{3}. \qquad (B)$$

Each formula is used with initial value $x_1 = 2.5$.

(ii) Show that one of these formulae produces a sequence which fails to converge, and use the other formula to calculate α correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [4]

SAT PREP SP www.satprep.co.in

4 **(i)** Prove the identity $\tan(45° + x) + \tan(45° - x) \equiv 2 \sec 2x$. [4]

SAT PREP SP www.satprep.co.uk

(ii) Hence sketch the graph of $y = \tan(45° + x) + \tan(45° - x)$ for $0° \leqslant x \leqslant 90°$. [3]

SAT PREP SP www.satprep.co.in

5 The equation of a curve is $2x^4 + xy^3 + y^4 = 10$.

(i) Show that $\frac{dy}{dx} = -\frac{8x^3 + y^3}{3xy^2 + 4y^3}$. [4]

© UCLES 2017

(ii) Hence show that there are two points on the curve at which the tangent is parallel to the x-axis and find the coordinates of these points. [4]

6 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 4\cos^2 y \tan x,$$

for $0 \leqslant x < \frac{1}{2}\pi$, and $x = 0$ when $y = \frac{1}{4}\pi$. Solve this differential equation and find the value of x when $y = \frac{1}{3}\pi$. [8]

SAT PREP SP www.satprep.co.in

SAT PREP SP www.satprep.co.in

7 **(a)** The complex number u is given by $u = 8 - 15\mathrm{i}$. Showing all necessary working, find the two square roots of u. Give answers in the form $a + \mathrm{i}b$, where the numbers a and b are real and exact. [5]

SAT PREP SP www.satprep.co.in

(b) On an Argand diagram, shade the region whose points represent complex numbers satisfying both the inequalities $|z - 2 - \mathrm{i}| \leqslant 2$ and $0 \leqslant \arg(z - \mathrm{i}) \leqslant \frac{1}{4}\pi$. [4]

SAT PREP SP www.satprep.co.in

8 Let $\mathrm{f}(x) = \dfrac{4x^2 + 9x - 8}{(x+2)(2x-1)}$.

(i) Express $\mathrm{f}(x)$ in the form $A + \dfrac{B}{x+2} + \dfrac{C}{2x-1}$. [4]

SAT PREP SP www.satprep.co.in

(ii) Hence show that $\int_1^4 \mathrm{f}(x)\,\mathrm{d}x = 6 + \frac{1}{2}\ln\left(\frac{16}{7}\right)$. [5]

SAT PREP SP www.satprep.co.in

9



The diagram shows the curve $y = (1 + x^2)e^{-\frac{1}{2}x}$ for $x \geqslant 0$. The shaded region R is enclosed by the curve, the x-axis and the lines $x = 0$ and $x = 2$.

(i) Find the exact values of the x-coordinates of the stationary points of the curve. [4]

SAT PREP SP www.satprep.co.uk

(ii) Show that the exact value of the area of R is $18 - \frac{42}{e}$. [5]

SAT PREP SP www.satprep.co.in

10 The equations of two lines l and m are $\mathbf{r} = 3\mathbf{i} - \mathbf{j} - 2\mathbf{k} + \lambda(-\mathbf{i} + \mathbf{j} + 4\mathbf{k})$ and $\mathbf{r} = 4\mathbf{i} + 4\mathbf{j} - 3\mathbf{k} + \mu(2\mathbf{i} + \mathbf{j} - 2\mathbf{k})$ respectively.

(i) Show that the lines do not intersect. [3]

(ii) Calculate the acute angle between the directions of the lines. [3]

SAT PREP SP www.satprep.co.in

(iii) Find the equation of the plane which passes through the point $(3, -2, -1)$ and which is parallel to both l and m. Give your answer in the form $ax + by + cz = d$. [5]

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

4382153981

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2017**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2017

1 Solve the inequality $|2x + 1| < 3|x - 2|$. [4]

SAT PREP SP www.satprep.co.uk

2 Expand $\frac{1}{\sqrt[3]{(1+6x)}}$ in ascending powers of x, up to and including the term in x^3, simplifying the coefficients. [4]

SAT PREP SP www.satprep.co.in

3 It is given that $x = \ln(1 - y) - \ln y$, where $0 < y < 1$.

(i) Show that $y = \dfrac{e^{-x}}{1 + e^{-x}}$. [2]

SAT PREP SP www.satprep.co.in

(ii) Hence show that $\int_0^1 y \, \mathrm{d}x = \ln\left(\frac{2\mathrm{e}}{\mathrm{e}+1}\right)$. [4]

© UCLES 2017

4 The parametric equations of a curve are

$$x = \ln\cos\theta, \quad y = 3\theta - \tan\theta,$$

where $0 \leqslant \theta < \frac{1}{2}\pi$.

(i) Express $\frac{\mathrm{d}y}{\mathrm{d}x}$ in terms of $\tan\theta$. [5]

SAT PREP SP www.satprep.co.uk

(ii) Find the exact y-coordinate of the point on the curve at which the gradient of the normal is equal to 1. [3]

SAT PREP SP www.satprep.co.in

5



The diagram shows a semicircle with centre O, radius r and diameter AB. The point P on its circumference is such that the area of the minor segment on AP is equal to half the area of the minor segment on BP. The angle AOP is x radians.

(i) Show that x satisfies the equation $x = \frac{1}{3}(\pi + \sin x)$. [3]

SAT PREP www.satprep.co.uk

(ii) Verify by calculation that x lies between 1 and 1.5. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use an iterative formula based on the equation in part **(i)** to determine x correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP www.satprep.co.in

6 The plane with equation $2x + 2y - z = 5$ is denoted by m. Relative to the origin O, the points A and B have coordinates (3, 4, 0) and (−1, 0, 2) respectively.

(i) Show that the plane m bisects AB at right angles. [5]

SAT PREP SP www.satprep.co.in

A second plane p is parallel to m and nearer to O. The perpendicular distance between the planes is 1.

(ii) Find the equation of p, giving your answer in the form $ax + by + cz = d$. [3]

SAT PREP SP www.satprep.co.in

7 Throughout this question the use of a calculator is not permitted.

The complex numbers u and w are defined by $u = -1 + 7\mathrm{i}$ and $w = 3 + 4\mathrm{i}$.

(i) Showing all your working, find in the form $x + \mathrm{i}y$, where x and y are real, the complex numbers $u - 2w$ and $\dfrac{u}{w}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

In an Argand diagram with origin O, the points A, B and C represent the complex numbers u, w and $u - 2w$ respectively.

(ii) Prove that angle $AOB = \frac{1}{4}\pi$. [2]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) State fully the geometrical relation between the line segments OB and CA. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

[Turn over

8 **(i)** By first expanding $2\sin(x - 30°)$, express $2\sin(x - 30°) - \cos x$ in the form $R\sin(x - \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. Give the exact value of R and the value of α correct to 2 decimal places. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence solve the equation

$$2\sin(x - 30°) - \cos x = 1,$$

for $0° < x < 180°$. [3]

SAT PREP SP www.satprep.co.in

9 **(i)** Express $\frac{1}{x(2x+3)}$ in partial fractions. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) The variables x and y satisfy the differential equation

$$x(2x+3)\frac{\mathrm{d}y}{\mathrm{d}x} = y,$$

and it is given that $y = 1$ when $x = 1$. Solve the differential equation and calculate the value of y when $x = 9$, giving your answer correct to 3 significant figures. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

10



The diagram shows the curve $y = \sin x \cos^2 2x$ for $0 \leqslant x \leqslant \frac{1}{4}\pi$ and its maximum point M.

(i) Using the substitution $u = \cos x$, find by integration the exact area of the shaded region bounded by the curve and the x-axis. [6]

SAT PREP SP www.satprep.co.in

(ii) Find the x-coordinate of M. Give your answer correct to 2 decimal places. [6]

SAT PREP SP www.satprep.co.uk

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

9035132693

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2017**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2017

1 Solve the equation $\ln(x^2 + 1) = 1 + 2\ln x$, giving your answer correct to 3 significant figures. [3]

SAT PREP SP www.satprep.co.uk

2 Solve the inequality $|x - 3| < 3x - 4$. [4]

SAT PREP SP www.satprep.co.in

3 **(i)** Express the equation $\cot\theta - 2\tan\theta = \sin 2\theta$ in the form $a\cos^4\theta + b\cos^2\theta + c = 0$, where a, b and c are constants to be determined. [3]

© UCLES 2017

(ii) Hence solve the equation $\cot\theta - 2\tan\theta = \sin 2\theta$ for $90° < \theta < 180°$. [2]

SAT PREP SP www.satprep.co.in

4 The parametric equations of a curve are

$$x = t^2 + 1, \quad y = 4t + \ln(2t - 1).$$

(i) Express $\frac{\mathrm{d}y}{\mathrm{d}x}$ in terms of t. [3]

(ii) Find the equation of the normal to the curve at the point where $t = 1$. Give your answer in the form $ax + by + c = 0$. [3]

SAT PREP SP www.satprep.co.in

5 In a certain chemical process a substance A reacts with and reduces a substance B. The masses of A and B at time t after the start of the process are x and y respectively. It is given that $\frac{dy}{dt} = -0.2xy$ and $x = \frac{10}{(1+t)^2}$. At the beginning of the process $y = 100$.

(i) Form a differential equation in y and t, and solve this differential equation. [6]

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the exact value approached by the mass of B as t becomes large. State what happens to the mass of A as t becomes large. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

[Turn over

6 Throughout this question the use of a calculator is not permitted.

The complex number $2 - i$ is denoted by u.

(i) It is given that u is a root of the equation $x^3 + ax^2 - 3x + b = 0$, where the constants a and b are real. Find the values of a and b. [4]

SAT PREP SP www.satprep.co.in

(ii) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying both the inequalities $|z - u| < 1$ and $|z| < |z + \mathrm{i}|$. [4]

SAT PREP SP www.satprep.co.in

7 **(i)** Prove that if $y = \dfrac{1}{\cos\theta}$ then $\dfrac{\mathrm{d}y}{\mathrm{d}\theta} = \sec\theta\tan\theta$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Prove the identity $\dfrac{1+\sin\theta}{1-\sin\theta} \equiv 2\sec^2\theta + 2\sec\theta\tan\theta - 1$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Hence find the exact value of $\int_0^{\frac{1}{4}\pi} \frac{1+\sin\theta}{1-\sin\theta}\,\mathrm{d}\theta$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

[Turn over

8 Let $f(x) = \dfrac{5x^2 - 7x + 4}{(3x + 2)(x^2 + 5)}$.

(i) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

9 Relative to the origin O, the point A has position vector given by $\overrightarrow{OA} = \mathbf{i} + 2\mathbf{j} + 4\mathbf{k}$. The line l has equation $\mathbf{r} = 9\mathbf{i} - \mathbf{j} + 8\mathbf{k} + \mu(3\mathbf{i} - \mathbf{j} + 2\mathbf{k})$.

(i) Find the position vector of the foot of the perpendicular from A to l. Hence find the position vector of the reflection of A in l. [5]

SAT PREP SP www.satprep.co.in

(ii) Find the equation of the plane through the origin which contains l. Give your answer in the form $ax + by + cz = d$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Find the exact value of the perpendicular distance of A from this plane. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

10



The diagram shows the curve $y = x^2 \cos 2x$ for $0 \leqslant x \leqslant \frac{1}{4}\pi$. The curve has a maximum point at M where $x = p$.

(i) Show that p satisfies the equation $p = \frac{1}{2}\tan^{-1}\left(\frac{1}{p}\right)$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Use the iterative formula $p_{n+1} = \frac{1}{2}\tan^{-1}\left(\frac{1}{p_n}\right)$ to determine the value of p correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

(iii) Find, showing all necessary working, the exact area of the region bounded by the curve and the x-axis. [5]

© UCLES 2017

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

5851032809

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2017**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2017

1 Prove the identity $\dfrac{\cot x - \tan x}{\cot x + \tan x} \equiv \cos 2x$. [3]

SAT PREP SP www.satprep.co.in

2 Expand $(3 + 2x)^{-3}$ in ascending powers of x up to and including the term in x^2, simplifying the coefficients. [4]

SAT PREP SP www.satprep.co.in

3 Using the substitution $u = e^x$, solve the equation $4e^{-x} = 3e^x + 4$. Give your answer correct to 3 significant figures. [4]

© UCLES 2017

4 Find the exact value of $\int_0^{\frac{1}{2}\pi} \theta \sin \frac{1}{2}\theta \, \mathrm{d}\theta$. [4]

© UCLES 2017

[Turn over

5 A curve has equation $y = \frac{2}{3}\ln(1 + 3\cos^2 x)$ for $0 \leqslant x \leqslant \frac{1}{2}\pi$.

(i) Express $\frac{\mathrm{d}y}{\mathrm{d}x}$ in terms of $\tan x$. [4]

SAT PREP SP www.satprep.co.in

(ii) Hence find the x-coordinate of the point on the curve where the gradient is -1. Give your answer correct to 3 significant figures. [2]

© UCLES 2017

6 The equation $\cot x = 1 - x$ has one root in the interval $0 < x < \pi$, denoted by α.

(i) Show by calculation that α is greater than 2.5. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Show that, if a sequence of values in the interval $0 < x < \pi$ given by the iterative formula $x_{n+1} = \pi + \tan^{-1}\left(\frac{1}{1 - x_n}\right)$ converges, then it converges to α. [2]

.....

.....

.....

.....

.....

.....

.....

.....

SAT PREP SP www.satprep.co.in

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use this iterative formula to determine α correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

[Turn over

7



The diagram shows a sketch of the curve $y = \dfrac{e^{\frac{1}{2}x}}{x}$ for $x > 0$, and its minimum point M.

(i) Find the x-coordinate of M. [4]

SJ SAT PREP www.satprep.co.in

(ii) Use the trapezium rule with two intervals to estimate the value of

$$\int_1^3 \frac{e^{\frac{1}{2}x}}{x}\,dx,$$

giving your answer correct to 2 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) The estimate found in part **(ii)** is denoted by E. Explain, without further calculation, whether another estimate found using the trapezium rule with four intervals would be greater than E or less than E. [1]

.....

.....

.....

.....

.....

.....

© UCLES 2017

[Turn over

8 In a certain chemical reaction, a compound A is formed from a compound B. The masses of A and B at time t after the start of the reaction are x and y respectively and the sum of the masses is equal to 50 throughout the reaction. At any time the rate of increase of the mass of A is proportional to the mass of B at that time.

(i) Explain why $\frac{\mathrm{d}x}{\mathrm{d}t} = k(50 - x)$, where k is a constant. [1]

.....

.....

.....

.....

.....

.....

It is given that $x = 0$ when $t = 0$, and $x = 25$ when $t = 10$.

(ii) Solve the differential equation in part **(i)** and express x in terms of t. [8]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

SAT PREP SP www.satprep.co.in

9 Let $\mathrm{f}(x) = \dfrac{3x^2 - 4}{x^2(3x + 2)}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence show that $\int_1^2 \mathrm{f}(x)\,\mathrm{d}x = \ln\left(\frac{25}{8}\right) - 1$. [5]

SAT PREP SP www.satprep.co.in

10 The points A and B have position vectors given by $\overrightarrow{OA} = \mathbf{i} - 2\mathbf{j} + 2\mathbf{k}$ and $\overrightarrow{OB} = 3\mathbf{i} + \mathbf{j} + \mathbf{k}$. The line l has equation $\mathbf{r} = 2\mathbf{i} + \mathbf{j} + m\mathbf{k} + \mu(\mathbf{i} - 2\mathbf{j} - 4\mathbf{k})$, where m is a constant.

(i) Given that the line l intersects the line passing through A and B, find the value of m. [5]

SAT PREP SP www.satprep.co.in

(ii) Find the equation of the plane which is parallel to $\mathbf{i} - 2\mathbf{j} - 4\mathbf{k}$ and contains the points A and B. Give your answer in the form $ax + by + cz = d$. [5]

SAT PREP SP www.satprep.co.in

11 Throughout this question the use of a calculator is not permitted.

(a) The complex numbers z and w satisfy the equations

$$z + (1 + \mathrm{i})w = \mathrm{i} \qquad \text{and} \qquad (1 - \mathrm{i})z + \mathrm{i}w = 1.$$

Solve the equations for z and w, giving your answers in the form $x + \mathrm{i}y$, where x and y are real. [6]

SAT PREP SP www.satprep.co.uk

(b) The complex numbers u and v are given by $u = 1 + (2\sqrt{3})\mathrm{i}$ and $v = 3 + 2\mathrm{i}$. In an Argand diagram, u and v are represented by the points A and B. A third point C lies in the first quadrant and is such that $BC = 2AB$ and angle $ABC = 90°$. Find the complex number z represented by C, giving your answer in the form $x + \mathrm{i}y$, where x and y are real and exact. [4]

SAT PREP SP www.satprep.co.uk

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

CANDIDATE NAME	
CENTRE NUMBER	CANDIDATE NUMBER

8450477717

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **February/March 2017**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

© UCLES 2017

1 Solve the equation $\ln(1 + 2^x) = 2$, giving your answer correct to 3 decimal places. [3]

SAT PREP SP www.satprep.co.uk

2 Solve the inequality $|x - 4| < 2|3x + 1|$. [4]

SAT PREP SP www.satprep.co.in

3 **(i)** By sketching suitable graphs, show that the equation $e^{-\frac{1}{2}x} = 4 - x^2$ has one positive root and one negative root. [2]

(ii) Verify by calculation that the negative root lies between -1 and -1.5. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

(iii) Use the iterative formula $x_{n+1} = -\sqrt{\left(4 - e^{-\frac{1}{2}x_n}\right)}$ to determine this root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

SAT PREP SP www.satprep.co.in

4 **(i)** Express $8\cos\theta - 15\sin\theta$ in the form $R\cos(\theta + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$, stating the exact value of R and giving the value of α correct to 2 decimal places. [3]

SAT PREP SP www.satprep.co.in

(ii) Hence solve the equation

$$8\cos 2x - 15\sin 2x = 4,$$

for $0° < x < 180°$. [4]

© UCLES 2017

5 The curve with equation $y = e^{-ax} \tan x$, where a is a positive constant, has only one point in the interval $0 < x < \frac{1}{2}\pi$ at which the tangent is parallel to the x-axis. Find the value of a and state the exact value of the x-coordinate of this point. [7]

SAT PREP SP www.satprep.co.uk

SAT PREP SP www.satprep.co.in

6 The line l has equation $\mathbf{r} = \mathbf{i} + 2\mathbf{j} - 3\mathbf{k} + \lambda(2\mathbf{i} - \mathbf{j} + \mathbf{k})$. The plane p has equation $3x + y - 5z = 20$.

(i) Show that the line l lies in the plane p. [3]

SAT PREP SP www.satprep.co.in

(ii) A second plane is parallel to l, perpendicular to p and contains the point with position vector $3\mathbf{i} - \mathbf{j} + 2\mathbf{k}$. Find the equation of this plane, giving your answer in the form $ax + by + cz = d$. [5]

SAT PREP SP www.satprep.co.in

7



A water tank has vertical sides and a horizontal rectangular base, as shown in the diagram. The area of the base is $2\,\text{m}^2$. At time $t = 0$ the tank is empty and water begins to flow into it at a rate of $1\,\text{m}^3$ per hour. At the same time water begins to flow out from the base at a rate of $0.2\sqrt{h}\,\text{m}^3$ per hour, where h m is the depth of water in the tank at time t hours.

(i) Form a differential equation satisfied by h and t, and show that the time T hours taken for the depth of water to reach 4 m is given by

$$T = \int_0^4 \frac{10}{5 - \sqrt{h}}\,\mathrm{d}h. \qquad [3]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

(ii) Using the substitution $u = 5 - \sqrt{h}$, find the value of T. [6]

SAT PREP SP www.satprep.co.uk

8 Throughout this question the use of a calculator is not permitted.

The polynomial $z^4 + 3z^2 + 6z + 10$ is denoted by $p(z)$. The complex number $-1 + i$ is denoted by u.

(i) Showing all your working, verify that u is a root of the equation $p(z) = 0$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the other three roots of the equation $p(z) = 0$. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2017

9 Let $f(x) = \dfrac{x(6-x)}{(2+x)(4+x^2)}$.

(i) Express $f(x)$ in partial fractions. [5]

SAT PREP SP www.satprep.co.in

(ii) Hence obtain the expansion of $f(x)$ in ascending powers of x, up to and including the term in x^2. [5]

SAT PREP SP www.satprep.co.in

10



The diagram shows the curve $y = (\ln x)^2$. The x-coordinate of the point P is equal to e, and the normal to the curve at P meets the x-axis at Q.

(i) Find the x-coordinate of Q. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Show that $\int \ln x \, dx = x \ln x - x + c$, where c is a constant. [1]

.....

.....

.....

.....

.....

© UCLES 2017

(iii) Using integration by parts, or otherwise, find the exact value of the area of the shaded region between the curve, the x-axis and the normal PQ. [5]

© UCLES 2017

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2016**

1 hour 45 minutes

Additional Materials: List of Formulae (MF9)

5711204026

READ THESE INSTRUCTIONS FIRST

An answer booklet is provided inside this question paper. You should follow the instructions on the front cover of the answer booklet. If you need additional answer paper ask the invigilator for a continuation booklet.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

SAT PREP www.satprep.co.in

This document consists of **3** printed pages, **1** blank page and **1** insert.

© UCLES 2016

CAMBRIDGE International Examinations

[Turn over

1 Solve the equation $\dfrac{3^x + 2}{3^x - 2} = 8$, giving your answer correct to 3 decimal places. [3]

2 Expand $(2 - x)(1 + 2x)^{-\frac{3}{2}}$ in ascending powers of x, up to and including the term in x^2, simplifying the coefficients. [4]

3 Express the equation $\sec\theta = 3\cos\theta + \tan\theta$ as a quadratic equation in $\sin\theta$. Hence solve this equation for $-90° < \theta < 90°$. [5]

4 The equation of a curve is $xy(x - 6y) = 9a^3$, where a is a non-zero constant. Show that there is only one point on the curve at which the tangent is parallel to the x-axis, and find the coordinates of this point. [7]

5 **(i)** Prove the identity $\tan 2\theta - \tan\theta \equiv \tan\theta \sec 2\theta$. [4]

(ii) Hence show that $\displaystyle\int_0^{\frac{1}{6}\pi} \tan\theta \sec 2\theta \, \mathrm{d}\theta = \tfrac{1}{2}\ln\tfrac{3}{2}$. [4]

6 **(i)** By sketching a suitable pair of graphs, show that the equation

$$\operatorname{cosec}\tfrac{1}{2}x = \tfrac{1}{3}x + 1$$

has one root in the interval $0 < x \leqslant \pi$. [2]

(ii) Show by calculation that this root lies between 1.4 and 1.6. [2]

(iii) Show that, if a sequence of values in the interval $0 < x \leqslant \pi$ given by the iterative formula

$$x_{n+1} = 2\sin^{-1}\left(\frac{3}{x_n + 3}\right)$$

converges, then it converges to the root of the equation in part **(i)**. [2]

(iv) Use this iterative formula to calculate the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

© UCLES 2016

7



The diagram shows part of the curve $y = (2x - x^2)e^{\frac{1}{2}x}$ and its maximum point M.

(i) Find the exact x-coordinate of M. [4]

(ii) Find the exact value of the area of the shaded region bounded by the curve and the positive x-axis. [5]

8 Two planes have equations $3x + y - z = 2$ and $x - y + 2z = 3$.

(i) Show that the planes are perpendicular. [3]

(ii) Find a vector equation for the line of intersection of the two planes. [6]

9 Throughout this question the use of a calculator is not permitted.

(a) Solve the equation $(1 + 2\mathrm{i})w^2 + 4w - (1 - 2\mathrm{i}) = 0$, giving your answers in the form $x + \mathrm{i}y$, where x and y are real. [5]

(b) On a sketch of an Argand diagram, shade the region whose points represent complex numbers satisfying the inequalities $|z - 1 - \mathrm{i}| \leqslant 2$ and $-\frac{1}{4}\pi \leqslant \arg z \leqslant \frac{1}{4}\pi$. [5]

10 A large field of area 4 km^2 is becoming infected with a soil disease. At time t years the area infected is x km^2 and the rate of growth of the infected area is given by the differential equation $\frac{\mathrm{d}x}{\mathrm{d}t} = kx(4 - x)$, where k is a positive constant. It is given that when $t = 0$, $x = 0.4$ and that when $t = 2$, $x = 2$.

(i) Solve the differential equation and show that $k = \frac{1}{4}\ln 3$. [9]

(ii) Find the value of t when 90% of the area of the field is infected. [2]

© UCLES 2016

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2016**

1 hour 45 minutes

Additional Materials: List of Formulae (MF9)

6635872224

READ THESE INSTRUCTIONS FIRST

An answer booklet is provided inside this question paper. You should follow the instructions on the front cover of the answer booklet. If you need additional answer paper ask the invigilator for a continuation booklet.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

SAT PREP www.satprep.co.in

This document consists of **3** printed pages, **1** blank page and **1** insert.

© UCLES 2016

CAMBRIDGE International Examinations

[Turn over

1 Solve the equation $\dfrac{3^x + 2}{3^x - 2} = 8$, giving your answer correct to 3 decimal places. [3]

2 Expand $(2 - x)(1 + 2x)^{-\frac{3}{2}}$ in ascending powers of x, up to and including the term in x^2, simplifying the coefficients. [4]

3 Express the equation $\sec\theta = 3\cos\theta + \tan\theta$ as a quadratic equation in $\sin\theta$. Hence solve this equation for $-90° < \theta < 90°$. [5]

4 The equation of a curve is $xy(x - 6y) = 9a^3$, where a is a non-zero constant. Show that there is only one point on the curve at which the tangent is parallel to the x-axis, and find the coordinates of this point. [7]

5 **(i)** Prove the identity $\tan 2\theta - \tan\theta \equiv \tan\theta \sec 2\theta$. [4]

(ii) Hence show that $\displaystyle\int_0^{\frac{1}{6}\pi} \tan\theta \sec 2\theta \, \mathrm{d}\theta = \tfrac{1}{2}\ln\tfrac{3}{2}$. [4]

6 **(i)** By sketching a suitable pair of graphs, show that the equation

$$\operatorname{cosec}\tfrac{1}{2}x = \tfrac{1}{3}x + 1$$

has one root in the interval $0 < x \leqslant \pi$. [2]

(ii) Show by calculation that this root lies between 1.4 and 1.6. [2]

(iii) Show that, if a sequence of values in the interval $0 < x \leqslant \pi$ given by the iterative formula

$$x_{n+1} = 2\sin^{-1}\left(\frac{3}{x_n + 3}\right)$$

converges, then it converges to the root of the equation in part **(i)**. [2]

(iv) Use this iterative formula to calculate the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

© UCLES 2016

7



The diagram shows part of the curve $y = (2x - x^2)e^{\frac{1}{2}x}$ and its maximum point M.

(i) Find the exact x-coordinate of M. [4]

(ii) Find the exact value of the area of the shaded region bounded by the curve and the positive x-axis. [5]

8 Two planes have equations $3x + y - z = 2$ and $x - y + 2z = 3$.

(i) Show that the planes are perpendicular. [3]

(ii) Find a vector equation for the line of intersection of the two planes. [6]

9 Throughout this question the use of a calculator is not permitted.

(a) Solve the equation $(1 + 2\mathrm{i})w^2 + 4w - (1 - 2\mathrm{i}) = 0$, giving your answers in the form $x + \mathrm{i}y$, where x and y are real. [5]

(b) On a sketch of an Argand diagram, shade the region whose points represent complex numbers satisfying the inequalities $|z - 1 - \mathrm{i}| \leqslant 2$ and $-\frac{1}{4}\pi \leqslant \arg z \leqslant \frac{1}{4}\pi$. [5]

10 A large field of area 4 km^2 is becoming infected with a soil disease. At time t years the area infected is x km^2 and the rate of growth of the infected area is given by the differential equation $\frac{\mathrm{d}x}{\mathrm{d}t} = kx(4 - x)$, where k is a positive constant. It is given that when $t = 0$, $x = 0.4$ and that when $t = 2$, $x = 2$.

(i) Solve the differential equation and show that $k = \frac{1}{4}\ln 3$. [9]

(ii) Find the value of t when 90% of the area of the field is infected. [2]

© UCLES 2016

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2016**

1 hour 45 minutes

Additional Materials: List of Formulae (MF9)

7240366882

READ THESE INSTRUCTIONS FIRST

An answer booklet is provided inside this question paper. You should follow the instructions on the front cover of the answer booklet. If you need additional answer paper ask the invigilator for a continuation booklet.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

SAT PREP www.satprep.co.in

This document consists of **4** printed pages and **1** insert.

© UCLES 2016

CAMBRIDGE International Examinations

1 It is given that $z = \ln(y + 2) - \ln(y + 1)$. Express y in terms of z. [3]

2 The equation of a curve is $y = \dfrac{\sin x}{1 + \cos x}$, for $-\pi < x < \pi$. Show that the gradient of the curve is positive for all x in the given interval. [4]

3 Express the equation $\cot 2\theta = 1 + \tan \theta$ as a quadratic equation in $\tan \theta$. Hence solve this equation for $0° < \theta < 180°$. [6]

4 The polynomial $4x^4 + ax^2 + 11x + b$, where a and b are constants, is denoted by $\mathrm{p}(x)$. It is given that $\mathrm{p}(x)$ is divisible by $x^2 - x + 2$.

(i) Find the values of a and b. [5]

(ii) When a and b have these values, find the real roots of the equation $\mathrm{p}(x) = 0$. [2]

5



The diagram shows a variable point P with coordinates (x, y) and the point N which is the foot of the perpendicular from P to the x-axis. P moves on a curve such that, for all $x \geqslant 0$, the gradient of the curve is equal in value to the area of the triangle OPN, where O is the origin.

(i) State a differential equation satisfied by x and y. [1]

The point with coordinates $(0, 2)$ lies on the curve.

(ii) Solve the differential equation to obtain the equation of the curve, expressing y in terms of x. [5]

(iii) Sketch the curve. [1]

6 Let $I = \displaystyle\int_1^4 \frac{(\sqrt{x}) - 1}{2(x + \sqrt{x})}\,\mathrm{d}x$.

(i) Using the substitution $u = \sqrt{x}$, show that $I = \displaystyle\int_1^2 \frac{u - 1}{u + 1}\,\mathrm{d}u$. [3]

(ii) Hence show that $I = 1 + \ln \frac{4}{9}$. [6]

© UCLES 2016

7 Throughout this question the use of a calculator is not permitted.

The complex number z is defined by $z = (\sqrt{2}) - (\sqrt{6})\mathrm{i}$. The complex conjugate of z is denoted by z^*.

(i) Find the modulus and argument of z. [2]

(ii) Express each of the following in the form $x + \mathrm{i}y$, where x and y are real and exact:

(a) $z + 2z^*$;

(b) $\dfrac{z^*}{\mathrm{i}z}$.

[4]

(iii) On a sketch of an Argand diagram with origin O, show the points A and B representing the complex numbers z^* and $\mathrm{i}z$ respectively. Prove that angle AOB is equal to $\frac{1}{6}\pi$. [3]

8 Let $\mathrm{f}(x) = \dfrac{3x^2 + x + 6}{(x + 2)(x^2 + 4)}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

(ii) Hence obtain the expansion of $\mathrm{f}(x)$ in ascending powers of x, up to and including the term in x^2. [5]

9



The diagram shows the curves $y = x\cos x$ and $y = \dfrac{k}{x}$, where k is a constant, for $0 < x \leqslant \frac{1}{2}\pi$. The curves touch at the point where $x = a$.

(i) Show that a satisfies the equation $\tan a = \dfrac{2}{a}$. [5]

(ii) Use the iterative formula $a_{n+1} = \tan^{-1}\left(\dfrac{2}{a_n}\right)$ to determine a correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

(iii) Hence find the value of k correct to 2 decimal places. [2]

[Question 10 is printed on the next page.]

10 The line l has vector equation $\mathbf{r} = \mathbf{i} + 2\mathbf{j} + \mathbf{k} + \lambda(2\mathbf{i} - \mathbf{j} + \mathbf{k})$.

(i) Find the position vectors of the two points on the line whose distance from the origin is $\sqrt{(10)}$. [5]

(ii) The plane p has equation $ax + y + z = 5$, where a is a constant. The acute angle between the line l and the plane p is equal to $\sin^{-1}\left(\frac{2}{3}\right)$. Find the possible values of a. [5]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2016**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

4146704902

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

© UCLES 2016

1 **(i)** Solve the equation $2|x - 1| = 3|x|$. [3]

(ii) Hence solve the equation $2|5^x - 1| = 3|5^x|$, giving your answer correct to 3 significant figures. [2]

2 Find the exact value of $\int_0^{\frac{1}{2}} xe^{-2x}\,dx$. [5]

3 By expressing the equation $\operatorname{cosec}\theta = 3\sin\theta + \cot\theta$ in terms of $\cos\theta$ only, solve the equation for $0° < \theta < 180°$. [5]

4 The variables x and y satisfy the differential equation

$$x\frac{dy}{dx} = y(1 - 2x^2),$$

and it is given that $y = 2$ when $x = 1$. Solve the differential equation and obtain an expression for y in terms of x in a form not involving logarithms. [6]

5 The curve with equation $y = \sin x \cos 2x$ has one stationary point in the interval $0 < x < \frac{1}{2}\pi$. Find the x-coordinate of this point, giving your answer correct to 3 significant figures. [6]

6 **(i)** By sketching a suitable pair of graphs, show that the equation

$$5e^{-x} = \sqrt{x}$$

has one root. [2]

(ii) Show that, if a sequence of values given by the iterative formula

$$x_{n+1} = \tfrac{1}{2}\ln\left(\frac{25}{x_n}\right)$$

converges, then it converges to the root of the equation in part **(i)**. [2]

(iii) Use this iterative formula, with initial value $x_1 = 1$, to calculate the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

7 The equation of a curve is $x^3 - 3x^2y + y^3 = 3$.

(i) Show that $\dfrac{dy}{dx} = \dfrac{x^2 - 2xy}{x^2 - y^2}$. [4]

(ii) Find the coordinates of the points on the curve where the tangent is parallel to the x-axis. [5]

© UCLES 2016

8 Let $\mathrm{f}(x) = \dfrac{4x^2 + 12}{(x+1)(x-3)^2}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

(ii) Hence obtain the expansion of $\mathrm{f}(x)$ in ascending powers of x, up to and including the term in x^2. [5]

9 With respect to the origin O, the points A, B, C, D have position vectors given by

$$\overrightarrow{OA} = \mathbf{i} + 3\mathbf{j} + 2\mathbf{k}, \quad \overrightarrow{OB} = 2\mathbf{i} + \mathbf{j} - \mathbf{k}, \quad \overrightarrow{OC} = 2\mathbf{i} + 4\mathbf{j} + \mathbf{k}, \quad \overrightarrow{OD} = -3\mathbf{i} + \mathbf{j} + 2\mathbf{k}.$$

(i) Find the equation of the plane containing A, B and C, giving your answer in the form $ax + by + cz = d$. [6]

(ii) The line through D parallel to OA meets the plane with equation $x + 2y - z = 7$ at the point P. Find the position vector of P and show that the length of DP is $2\sqrt{(14)}$. [5]

10 **(a)** Showing all your working and without the use of a calculator, find the square roots of the complex number $7 - (6\sqrt{2})\mathrm{i}$. Give your answers in the form $x + \mathrm{i}y$, where x and y are real and exact. [5]

(b) **(i)** On an Argand diagram, sketch the loci of points representing complex numbers w and z such that $|w - 1 - 2\mathrm{i}| = 1$ and $\arg(z - 1) = \frac{3}{4}\pi$. [4]

(ii) Calculate the least value of $|w - z|$ for points on these loci. [2]

© UCLES 2016

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2016**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

5402648439

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

www.satprep.org

This document consists of **3** printed pages and **1** blank page.

© UCLES 2016

1 Use logarithms to solve the equation $4^{3x-1} = 3(5^x)$, giving your answer correct to 3 decimal places. [4]

2 Expand $\dfrac{1}{\sqrt{(1-2x)}}$ in ascending powers of x, up to and including the term in x^3, simplifying the coefficients. [4]

3 Find the exact value of $\displaystyle\int_0^{\frac{1}{2}\pi} x^2 \sin 2x \, \mathrm{d}x$. [5]

4 The curve with equation $y = \dfrac{(\ln x)^2}{x}$ has two stationary points. Find the exact values of the coordinates of these points. [6]

5 **(i)** Prove the identity $\cos 4\theta - 4\cos 2\theta \equiv 8\sin^4\theta - 3$. [4]

(ii) Hence solve the equation

$$\cos 4\theta = 4\cos 2\theta + 3,$$

for $0° \leqslant \theta \leqslant 360°$. [4]

6 The variables x and θ satisfy the differential equation

$$(3 + \cos 2\theta)\frac{\mathrm{d}x}{\mathrm{d}\theta} = x \sin 2\theta,$$

and it is given that $x = 3$ when $\theta = \frac{1}{4}\pi$.

(i) Solve the differential equation and obtain an expression for x in terms of θ. [7]

(ii) State the least value taken by x. [1]

7 Let $\mathrm{f}(x) = \dfrac{4x^2 + 7x + 4}{(2x+1)(x+2)}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

(ii) Show that $\displaystyle\int_0^4 \mathrm{f}(x) \, \mathrm{d}x = 8 - \ln 3$. [5]

© UCLES 2016

8



The diagram shows the curve $y = \operatorname{cosec} x$ for $0 < x < \pi$ and part of the curve $y = \mathrm{e}^{-x}$. When $x = a$, the tangents to the curves are parallel.

(i) By differentiating $\dfrac{1}{\sin x}$, show that if $y = \operatorname{cosec} x$ then $\dfrac{\mathrm{d}y}{\mathrm{d}x} = -\operatorname{cosec} x \cot x$. [3]

(ii) By equating the gradients of the curves at $x = a$, show that

$$a = \tan^{-1}\left(\frac{\mathrm{e}^{a}}{\sin a}\right).$$ [2]

(iii) Verify by calculation that a lies between 1 and 1.5. [2]

(iv) Use an iterative formula based on the equation in part **(ii)** to determine a correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

9 The points A, B and C have position vectors, relative to the origin O, given by $\overrightarrow{OA} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$, $\overrightarrow{OB} = 4\mathbf{j} + \mathbf{k}$ and $\overrightarrow{OC} = 2\mathbf{i} + 5\mathbf{j} - \mathbf{k}$. A fourth point D is such that the quadrilateral $ABCD$ is a parallelogram.

(i) Find the position vector of D and verify that the parallelogram is a rhombus. [5]

(ii) The plane p is parallel to OA and the line BC lies in p. Find the equation of p, giving your answer in the form $ax + by + cz = d$. [5]

10 **(a)** Showing all necessary working, solve the equation $\mathrm{i}z^2 + 2z - 3\mathrm{i} = 0$, giving your answers in the form $x + \mathrm{i}y$, where x and y are real and exact. [5]

(b) **(i)** On a sketch of an Argand diagram, show the locus representing complex numbers satisfying the equation $|z| = |z - 4 - 3\mathrm{i}|$. [2]

(ii) Find the complex number represented by the point on the locus where $|z|$ is least. Find the modulus and argument of this complex number, giving the argument correct to 2 decimal places. [3]

© UCLES 2016

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2016**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

6768939761

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

www.satprep.org

This document consists of **3** printed pages and **1** blank page.

© UCLES 2016

1 Solve the inequality $2|x-2| > |3x+1|$. [4]

2 The variables x and y satisfy the relation $3^y = 4^{2-x}$.

(i) By taking logarithms, show that the graph of y against x is a straight line. State the exact value of the gradient of this line. [3]

(ii) Calculate the exact x-coordinate of the point of intersection of this line with the line with equation $y = 2x$, simplifying your answer. [2]

3 **(i)** Express $(\sqrt{5})\cos x + 2\sin x$ in the form $R\cos(x-\alpha)$, where $R > 0$ and $0° < \alpha < 90°$, giving the value of α correct to 2 decimal places. [3]

(ii) Hence solve the equation

$$(\sqrt{5})\cos\tfrac{1}{2}x + 2\sin\tfrac{1}{2}x = 1.2,$$

for $0° < x < 360°$. [3]

4 The parametric equations of a curve are

$$x = t + \cos t, \qquad y = \ln(1+\sin t),$$

where $-\frac{1}{2}\pi < t < \frac{1}{2}\pi$.

(i) Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \sec t$. [5]

(ii) Hence find the x-coordinates of the points on the curve at which the gradient is equal to 3. Give your answers correct to 3 significant figures. [3]

5 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \mathrm{e}^{-2y}\tan^2 x,$$

for $0 \leqslant x < \frac{1}{2}\pi$, and it is given that $y = 0$ when $x = 0$. Solve the differential equation and calculate the value of y when $x = \frac{1}{4}\pi$. [8]

6 The curve with equation $y = x^2\cos\frac{1}{2}x$ has a stationary point at $x = p$ in the interval $0 < x < \pi$.

(i) Show that p satisfies the equation $\tan\frac{1}{2}p = \dfrac{4}{p}$. [3]

(ii) Verify by calculation that p lies between 2 and 2.5. [2]

(iii) Use the iterative formula $p_{n+1} = 2\tan^{-1}\left(\dfrac{4}{p_n}\right)$ to determine the value of p correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

© UCLES 2016

7 Let $I = \int_0^1 \frac{x^5}{(1+x^2)^3}\,\mathrm{d}x$.

(i) Using the substitution $u = 1 + x^2$, show that $I = \int_1^2 \frac{(u-1)^2}{2u^3}\,\mathrm{d}u$. [3]

(ii) Hence find the exact value of I. [5]

8 The points A and B have position vectors, relative to the origin O, given by $\overrightarrow{OA} = \mathbf{i} + \mathbf{j} + \mathbf{k}$ and $\overrightarrow{OB} = 2\mathbf{i} + 3\mathbf{k}$. The line l has vector equation $\mathbf{r} = 2\mathbf{i} - 2\mathbf{j} - \mathbf{k} + \mu(-\mathbf{i} + 2\mathbf{j} + \mathbf{k})$.

(i) Show that the line passing through A and B does not intersect l. [4]

(ii) Show that the length of the perpendicular from A to l is $\frac{1}{\sqrt{2}}$. [5]

9 Throughout this question the use of a calculator is not permitted.

The complex numbers $-1 + 3\mathrm{i}$ and $2 - \mathrm{i}$ are denoted by u and v respectively. In an Argand diagram with origin O, the points A, B and C represent the numbers u, v and $u + v$ respectively.

(i) Sketch this diagram and state fully the geometrical relationship between OB and AC. [4]

(ii) Find, in the form $x + \mathrm{i}y$, where x and y are real, the complex number $\frac{u}{v}$. [3]

(iii) Prove that angle $AOB = \frac{3}{4}\pi$. [2]

10 Let $\mathrm{f}(x) = \frac{10x - 2x^2}{(x+3)(x-1)^2}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

(ii) Hence obtain the expansion of $\mathrm{f}(x)$ in ascending powers of x, up to and including the term in x^2. [5]

© UCLES 2016

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **February/March 2016**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

* 0 5 7 2 7 5 7 8 4 2 *

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

www.satprep.org

This document consists of **3** printed pages and **1** blank page.

© UCLES 2016

1 Solve the equation $\ln(x^2 + 4) = 2\ln x + \ln 4$, giving your answer in an exact form. [3]

2 Express the equation $\tan(\theta + 45°) - 2\tan(\theta - 45°) = 4$ as a quadratic equation in $\tan\theta$. Hence solve this equation for $0° \leqslant \theta \leqslant 180°$. [6]

3 The equation $x^5 - 3x^3 + x^2 - 4 = 0$ has one positive root.

(i) Verify by calculation that this root lies between 1 and 2. [2]

(ii) Show that the equation can be rearranged in the form

$$x = \sqrt[3]{\left(3x + \frac{4}{x^2} - 1\right)}.$$ [1]

(iii) Use an iterative formula based on this rearrangement to determine the positive root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

4 The polynomial $4x^3 + ax + 2$, where a is a constant, is denoted by $\mathrm{p}(x)$. It is given that $(2x + 1)$ is a factor of $\mathrm{p}(x)$.

(i) Find the value of a. [2]

(ii) When a has this value,

(a) factorise $\mathrm{p}(x)$, [2]

(b) solve the inequality $\mathrm{p}(x) > 0$, justifying your answer. [3]

5 Let $I = \displaystyle\int_0^1 \frac{9}{(3 + x^2)^2}\,\mathrm{d}x$.

(i) Using the substitution $x = (\sqrt{3})\tan\theta$, show that $I = \sqrt{3}\displaystyle\int_0^{\frac{1}{6}\pi} \cos^2\theta\,\mathrm{d}\theta$. [3]

(ii) Hence find the exact value of I. [4]

6 A curve has equation

$$\sin y \ln x = x - 2\sin y,$$

for $-\frac{1}{2}\pi \leqslant y \leqslant \frac{1}{2}\pi$.

(i) Find $\dfrac{\mathrm{d}y}{\mathrm{d}x}$ in terms of x and y. [5]

(ii) Hence find the exact x-coordinate of the point on the curve at which the tangent is parallel to the x-axis. [3]

© UCLES 2016

7 The variables x and y satisfy the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = x\mathrm{e}^{x+y},$$

and it is given that $y = 0$ when $x = 0$.

(i) Solve the differential equation and obtain an expression for y in terms of x. [7]

(ii) Explain briefly why x can only take values less than 1. [1]

8 The line l has equation $\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}$. The plane p has equation $\mathbf{r}.\begin{pmatrix} 2 \\ -1 \\ -1 \end{pmatrix} = 6$.

(i) Show that l is parallel to p. [3]

(ii) A line m lies in the plane p and is perpendicular to l. The line m passes through the point with coordinates (5, 3, 1). Find a vector equation for m. [6]

9 Let $\mathrm{f}(x) = \dfrac{3x^3 + 6x - 8}{x(x^2 + 2)}$.

(i) Express $\mathrm{f}(x)$ in the form $A + \dfrac{B}{x} + \dfrac{Cx + D}{x^2 + 2}$. [5]

(ii) Show that $\displaystyle\int_1^2 \mathrm{f}(x)\,\mathrm{d}x = 3 - \ln 4$. [5]

10 **(a)** Find the complex number z satisfying the equation $z^* + 1 = 2\mathrm{i}z$, where z^* denotes the complex conjugate of z. Give your answer in the form $x + \mathrm{i}y$, where x and y are real. [5]

(b) **(i)** On a sketch of an Argand diagram, shade the region whose points represent complex numbers satisfying the inequalities $|z + 1 - 3\mathrm{i}| \leqslant 1$ and $\mathrm{Im}\, z \geqslant 3$, where $\mathrm{Im}\, z$ denotes the imaginary part of z. [4]

(ii) Determine the difference between the greatest and least values of $\arg z$ for points lying in this region. [2]

© UCLES 2016

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2015**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

2390741568

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

© UCLES 2015

1 Solve the inequality $|2x - 5| > 3|2x + 1|$. [4]

2 Using the substitution $u = 3^x$, solve the equation $3^x + 3^{2x} = 3^{3x}$ giving your answer correct to 3 significant figures. [5]

3 The angles θ and ϕ lie between 0° and 180°, and are such that

$$\tan(\theta - \phi) = 3 \quad \text{and} \quad \tan\theta + \tan\phi = 1.$$

Find the possible values of θ and ϕ. [6]

4 The equation $x^3 - x^2 - 6 = 0$ has one real root, denoted by α.

(i) Find by calculation the pair of consecutive integers between which α lies. [2]

(ii) Show that, if a sequence of values given by the iterative formula

$$x_{n+1} = \sqrt{\left(x_n + \frac{6}{x_n}\right)}$$

converges, then it converges to α. [2]

(iii) Use this iterative formula to determine α correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

5 The equation of a curve is $y = \mathrm{e}^{-2x}\tan x$, for $0 \leqslant x < \frac{1}{2}\pi$.

(i) Obtain an expression for $\dfrac{\mathrm{d}y}{\mathrm{d}x}$ and show that it can be written in the form $\mathrm{e}^{-2x}(a + b\tan x)^2$, where a and b are constants. [5]

(ii) Explain why the gradient of the curve is never negative. [1]

(iii) Find the value of x for which the gradient is least. [1]

6 The polynomial $8x^3 + ax^2 + bx - 1$, where a and b are constants, is denoted by $\mathrm{p}(x)$. It is given that $(x + 1)$ is a factor of $\mathrm{p}(x)$ and that when $\mathrm{p}(x)$ is divided by $(2x + 1)$ the remainder is 1.

(i) Find the values of a and b. [5]

(ii) When a and b have these values, factorise $\mathrm{p}(x)$ completely. [3]

© UCLES 2015

7 The points A, B and C have position vectors, relative to the origin O, given by

$$\overrightarrow{OA} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 3 \\ 0 \\ 1 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OC} = \begin{pmatrix} 1 \\ 1 \\ 4 \end{pmatrix}.$$

The plane m is perpendicular to AB and contains the point C.

(i) Find a vector equation for the line passing through A and B. [2]

(ii) Obtain the equation of the plane m, giving your answer in the form $ax + by + cz = d$. [2]

(iii) The line through A and B intersects the plane m at the point N. Find the position vector of N and show that $CN = \sqrt{(13)}$. [5]

8 The variables x and θ satisfy the differential equation

$$\frac{\mathrm{d}x}{\mathrm{d}\theta} = (x + 2)\sin^2 2\theta,$$

and it is given that $x = 0$ when $\theta = 0$. Solve the differential equation and calculate the value of x when $\theta = \frac{1}{4}\pi$, giving your answer correct to 3 significant figures. [9]

9 The complex number $3 - \mathrm{i}$ is denoted by u. Its complex conjugate is denoted by u^*.

(i) On an Argand diagram with origin O, show the points A, B and C representing the complex numbers u, u^* and $u^* - u$ respectively. What type of quadrilateral is $OABC$? [4]

(ii) Showing your working and without using a calculator, express $\dfrac{u^*}{u}$ in the form $x + \mathrm{i}y$, where x and y are real. [3]

(iii) By considering the argument of $\dfrac{u^*}{u}$, prove that

$$\tan^{-1}\left(\tfrac{3}{4}\right) = 2\tan^{-1}\left(\tfrac{1}{3}\right).$$ [3]

10



The diagram shows the curve $y = \dfrac{x^2}{1 + x^3}$ for $x \geqslant 0$, and its maximum point M. The shaded region R is enclosed by the curve, the x-axis and the lines $x = 1$ and $x = p$.

(i) Find the exact value of the x-coordinate of M. [4]

(ii) Calculate the value of p for which the area of R is equal to 1. Give your answer correct to 3 significant figures. [6]

© UCLES 2015

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2015**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

* 0 1 0 7 1 0 7 4 5 *

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

© UCLES 2015

1 Solve the inequality $|2x - 5| > 3|2x + 1|$. [4]

2 Using the substitution $u = 3^x$, solve the equation $3^x + 3^{2x} = 3^{3x}$ giving your answer correct to 3 significant figures. [5]

3 The angles θ and ϕ lie between 0° and 180°, and are such that

$$\tan(\theta - \phi) = 3 \quad \text{and} \quad \tan\theta + \tan\phi = 1.$$

Find the possible values of θ and ϕ. [6]

4 The equation $x^3 - x^2 - 6 = 0$ has one real root, denoted by α.

(i) Find by calculation the pair of consecutive integers between which α lies. [2]

(ii) Show that, if a sequence of values given by the iterative formula

$$x_{n+1} = \sqrt{\left(x_n + \frac{6}{x_n}\right)}$$

converges, then it converges to α. [2]

(iii) Use this iterative formula to determine α correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

5 The equation of a curve is $y = e^{-2x}\tan x$, for $0 \leqslant x < \frac{1}{2}\pi$.

(i) Obtain an expression for $\frac{dy}{dx}$ and show that it can be written in the form $e^{-2x}(a + b\tan x)^2$, where a and b are constants. [5]

(ii) Explain why the gradient of the curve is never negative. [1]

(iii) Find the value of x for which the gradient is least. [1]

6 The polynomial $8x^3 + ax^2 + bx - 1$, where a and b are constants, is denoted by $p(x)$. It is given that $(x + 1)$ is a factor of $p(x)$ and that when $p(x)$ is divided by $(2x + 1)$ the remainder is 1.

(i) Find the values of a and b. [5]

(ii) When a and b have these values, factorise $p(x)$ completely. [3]

© UCLES 2015

7 The points A, B and C have position vectors, relative to the origin O, given by

$$\overrightarrow{OA} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 3 \\ 0 \\ 1 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OC} = \begin{pmatrix} 1 \\ 1 \\ 4 \end{pmatrix}.$$

The plane m is perpendicular to AB and contains the point C.

(i) Find a vector equation for the line passing through A and B. [2]

(ii) Obtain the equation of the plane m, giving your answer in the form $ax + by + cz = d$. [2]

(iii) The line through A and B intersects the plane m at the point N. Find the position vector of N and show that $CN = \sqrt{(13)}$. [5]

8 The variables x and θ satisfy the differential equation

$$\frac{\mathrm{d}x}{\mathrm{d}\theta} = (x + 2)\sin^2 2\theta,$$

and it is given that $x = 0$ when $\theta = 0$. Solve the differential equation and calculate the value of x when $\theta = \frac{1}{4}\pi$, giving your answer correct to 3 significant figures. [9]

9 The complex number $3 - \mathrm{i}$ is denoted by u. Its complex conjugate is denoted by u^*.

(i) On an Argand diagram with origin O, show the points A, B and C representing the complex numbers u, u^* and $u^* - u$ respectively. What type of quadrilateral is $OABC$? [4]

(ii) Showing your working and without using a calculator, express $\dfrac{u^*}{u}$ in the form $x + \mathrm{i}y$, where x and y are real. [3]

(iii) By considering the argument of $\dfrac{u^*}{u}$, prove that

$$\tan^{-1}\left(\tfrac{3}{4}\right) = 2\tan^{-1}\left(\tfrac{1}{3}\right).$$ [3]

10



The diagram shows the curve $y = \dfrac{x^2}{1 + x^3}$ for $x \geqslant 0$, and its maximum point M. The shaded region R is enclosed by the curve, the x-axis and the lines $x = 1$ and $x = p$.

(i) Find the exact value of the x-coordinate of M. [4]

(ii) Calculate the value of p for which the area of R is equal to 1. Give your answer correct to 3 significant figures. [6]

© UCLES 2015

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2015**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

8940457897

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

© UCLES 2015

1 Sketch the graph of $y = e^{ax} - 1$ where a is a positive constant. [2]

2 Given that $\sqrt[3]{(1 + 9x)} \approx 1 + 3x + ax^2 + bx^3$ for small values of x, find the values of the coefficients a and b. [3]

3 A curve has equation

$$y = \frac{2 - \tan x}{1 + \tan x}.$$

Find the equation of the tangent to the curve at the point for which $x = \frac{1}{4}\pi$, giving the answer in the form $y = mx + c$ where c is correct to 3 significant figures. [6]

4 A curve has parametric equations

$$x = t^2 + 3t + 1, \qquad y = t^4 + 1.$$

The point P on the curve has parameter p. It is given that the gradient of the curve at P is 4.

(i) Show that $p = \sqrt[3]{(2p + 3)}$. [3]

(ii) Verify by calculation that the value of p lies between 1.8 and 2.0. [2]

(iii) Use an iterative formula based on the equation in part **(i)** to find the value of p correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

5 Use the substitution $u = 4 - 3\cos x$ to find the exact value of $\displaystyle\int_0^{\frac{1}{2}\pi} \frac{9\sin 2x}{\sqrt{(4 - 3\cos x)}}\,\mathrm{d}x$. [8]

6 The angles A and B are such that

$$\sin(A + 45°) = (2\sqrt{2})\cos A \quad \text{and} \quad 4\sec^2 B + 5 = 12\tan B.$$

Without using a calculator, find the exact value of $\tan(A - B)$. [8]

7 **(i)** Show that $(x + 1)$ is a factor of $4x^3 - x^2 - 11x - 6$. [2]

(ii) Find $\displaystyle\int \frac{4x^2 + 9x - 1}{4x^3 - x^2 - 11x - 6}\,\mathrm{d}x$. [8]

8 A plane has equation $4x - y + 5z = 39$. A straight line is parallel to the vector $\mathbf{i} - 3\mathbf{j} + 4\mathbf{k}$ and passes through the point A (0, 2, −8). The line meets the plane at the point B.

(i) Find the coordinates of B. [3]

(ii) Find the acute angle between the line and the plane. [4]

(iii) The point C lies on the line and is such that the distance between C and B is twice the distance between A and B. Find the coordinates of each of the possible positions of the point C. [3]

© UCLES 2015

9 **(a)** It is given that $(1 + 3i)w = 2 + 4i$. Showing all necessary working, prove that the exact value of $|w^2|$ is 2 and find $\arg(w^2)$ correct to 3 significant figures. [6]

(b) On a single Argand diagram sketch the loci $|z| = 5$ and $|z - 5| = |z|$. Hence determine the complex numbers represented by points common to both loci, giving each answer in the form $re^{i\theta}$. [4]

10 Naturalists are managing a wildlife reserve to increase the number of plants of a rare species. The number of plants at time t years is denoted by N, where N is treated as a continuous variable.

(i) It is given that the rate of increase of N with respect to t is proportional to $(N - 150)$. Write down a differential equation relating N, t and a constant of proportionality. [1]

(ii) Initially, when $t = 0$, the number of plants was 650. It was noted that, at a time when there were 900 plants, the number of plants was increasing at a rate of 60 per year. Express N in terms of t. [7]

(iii) The naturalists had a target of increasing the number of plants from 650 to 2000 within 15 years. Will this target be met? [2]

SAT PREP SP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2015**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

8254149865

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

© UCLES 2015

1 Use logarithms to solve the equation $2^{5x} = 3^{2x+1}$, giving the answer correct to 3 significant figures. [4]

2 Use the trapezium rule with three intervals to find an approximation to

$$\int_0^3 |3^x - 10| \, \mathrm{d}x.$$ [4]

3 Show that, for small values of x^2,

$$\left(1 - 2x^2\right)^{-2} - \left(1 + 6x^2\right)^{\frac{2}{3}} \approx kx^4,$$

where the value of the constant k is to be determined. [6]

4 The equation of a curve is

$$y = 3\cos 2x + 7\sin x + 2.$$

Find the x-coordinates of the stationary points in the interval $0 \leqslant x \leqslant \pi$. Give each answer correct to 3 significant figures. [7]

5 **(a)** Find $\int (4 + \tan^2 2x) \, \mathrm{d}x$. [3]

(b) Find the exact value of $\displaystyle\int_{\frac{1}{4}\pi}^{\frac{1}{2}\pi} \frac{\sin\left(x + \frac{1}{6}\pi\right)}{\sin x} \, \mathrm{d}x$. [5]

6 The straight line l_1 passes through the points (0, 1, 5) and (2, −2, 1). The straight line l_2 has equation $\mathbf{r} = 7\mathbf{i} + \mathbf{j} + \mathbf{k} + \mu(\mathbf{i} + 2\mathbf{j} + 5\mathbf{k})$.

(i) Show that the lines l_1 and l_2 are skew. [6]

(ii) Find the acute angle between the direction of the line l_2 and the direction of the x-axis. [3]

7 Given that $y = 1$ when $x = 0$, solve the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 4x(3y^2 + 10y + 3),$$

obtaining an expression for y in terms of x. [9]

8 The complex number w is defined by $w = \dfrac{22 + 4\mathrm{i}}{(2 - \mathrm{i})^2}$.

(i) Without using a calculator, show that $w = 2 + 4\mathrm{i}$. [3]

(ii) It is given that p is a real number such that $\frac{1}{4}\pi \leqslant \arg(w + p) \leqslant \frac{3}{4}\pi$. Find the set of possible values of p. [3]

(iii) The complex conjugate of w is denoted by w^*. The complex numbers w and w^* are represented in an Argand diagram by the points S and T respectively. Find, in the form $|z - a| = k$, the equation of the circle passing through S, T and the origin. [3]

© UCLES 2015

9



The diagram shows the curve $y = x^2e^{2-x}$ and its maximum point M.

(i) Show that the x-coordinate of M is 2. [3]

(ii) Find the exact value of $\int_0^2 x^2e^{2-x}\,dx$. [6]

10



The diagram shows part of the curve with parametric equations

$$x = 2\ln(t+2), \qquad y = t^3 + 2t + 3.$$

(i) Find the gradient of the curve at the origin. [5]

(ii) At the point P on the curve, the value of the parameter is p. It is given that the gradient of the curve at P is $\frac{1}{2}$.

(a) Show that $p = \dfrac{1}{3p^2+2} - 2$. [1]

(b) By first using an iterative formula based on the equation in part **(a)**, determine the coordinates of the point P. Give the result of each iteration to 5 decimal places and each coordinate of P correct to 2 decimal places. [4]

© UCLES 2015

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2015**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

2399507033

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

© UCLES 2015

1 Use the trapezium rule with three intervals to estimate the value of

$$\int_0^{\frac{1}{2}\pi} \ln(1 + \sin x)\,\mathrm{d}x,$$

giving your answer correct to 2 decimal places. [3]

2 Using the substitution $u = 4^x$, solve the equation $4^x + 4^2 = 4^{x+2}$, giving your answer correct to 3 significant figures. [4]

3 A curve has equation $y = \cos x \cos 2x$. Find the x-coordinate of the stationary point on the curve in the interval $0 < x < \frac{1}{2}\pi$, giving your answer correct to 3 significant figures. [6]

4 **(i)** Express $3\sin\theta + 2\cos\theta$ in the form $R\sin(\theta + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$, stating the exact value of R and giving the value of α correct to 2 decimal places. [3]

(ii) Hence solve the equation

$$3\sin\theta + 2\cos\theta = 1,$$

for $0° < \theta < 180°$. [3]

5



The diagram shows a circle with centre O and radius r. The tangents to the circle at the points A and B meet at T, and the angle AOB is $2x$ radians. The shaded region is bounded by the tangents AT and BT, and by the minor arc AB. The perimeter of the shaded region is equal to the circumference of the circle.

(i) Show that x satisfies the equation

$$\tan x = \pi - x.$$ [3]

(ii) This equation has one root in the interval $0 < x < \frac{1}{2}\pi$. Verify by calculation that this root lies between 1 and 1.3. [2]

(iii) Use the iterative formula

$$x_{n+1} = \tan^{-1}(\pi - x_n)$$

to determine the root correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

© UCLES 2015

6 Let $I = \int_0^1 \frac{\sqrt{x}}{2-\sqrt{x}}\,\mathrm{d}x$.

(i) Using the substitution $u = 2 - \sqrt{x}$, show that $I = \int_1^2 \frac{2(2-u)^2}{u}\,\mathrm{d}u$. [4]

(ii) Hence show that $I = 8\ln 2 - 5$. [4]

7 The complex number u is given by $u = -1 + (4\sqrt{3})\mathrm{i}$.

(i) Without using a calculator and showing all your working, find the two square roots of u. Give your answers in the form $a + \mathrm{i}b$, where the real numbers a and b are exact. [5]

(ii) On an Argand diagram, sketch the locus of points representing complex numbers z satisfying the relation $|z - u| = 1$. Determine the greatest value of $\arg z$ for points on this locus. [4]

8 Let $\mathrm{f}(x) = \frac{5x^2 + x + 6}{(3-2x)(x^2+4)}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

(ii) Hence obtain the expansion of $\mathrm{f}(x)$ in ascending powers of x, up to and including the term in x^2. [5]

9 The number of organisms in a population at time t is denoted by x. Treating x as a continuous variable, the differential equation satisfied by x and t is

$$\frac{\mathrm{d}x}{\mathrm{d}t} = \frac{x\mathrm{e}^{-t}}{k + \mathrm{e}^{-t}},$$

where k is a positive constant.

(i) Given that $x = 10$ when $t = 0$, solve the differential equation, obtaining a relation between x, k and t. [6]

(ii) Given also that $x = 20$ when $t = 1$, show that $k = 1 - \frac{2}{\mathrm{e}}$. [2]

(iii) Show that the number of organisms never reaches 48, however large t becomes. [2]

10 The points A and B have position vectors given by $\overrightarrow{OA} = 2\mathbf{i} - \mathbf{j} + 3\mathbf{k}$ and $\overrightarrow{OB} = \mathbf{i} + \mathbf{j} + 5\mathbf{k}$. The line l has equation $\mathbf{r} = \mathbf{i} + \mathbf{j} + 2\mathbf{k} + \mu(3\mathbf{i} + \mathbf{j} - \mathbf{k})$.

(i) Show that l does not intersect the line passing through A and B. [5]

(ii) Find the equation of the plane containing the line l and the point A. Give your answer in the form $ax + by + cz = d$. [6]

© UCLES 2015

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2015**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

9684619653

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

© UCLES 2015

1 Solve the equation $\ln(x+4) = 2\ln x + \ln 4$, giving your answer correct to 3 significant figures. [4]

2 Solve the inequality $|x-2| > 2x-3$. [4]

3 Solve the equation $\cot 2x + \cot x = 3$ for $0° < x < 180°$. [6]

4 The curve with equation $y = \dfrac{e^{2x}}{4+e^{3x}}$ has one stationary point. Find the exact values of the coordinates of this point. [6]

5 The parametric equations of a curve are

$$x = a\cos^4 t, \quad y = a\sin^4 t,$$

where a is a positive constant.

(i) Express $\dfrac{dy}{dx}$ in terms of t. [3]

(ii) Show that the equation of the tangent to the curve at the point with parameter t is

$$x\sin^2 t + y\cos^2 t = a\sin^2 t\cos^2 t.$$ [3]

(iii) Hence show that if the tangent meets the x-axis at P and the y-axis at Q, then

$$OP + OQ = a,$$

where O is the origin. [2]

6 It is given that $\displaystyle\int_0^a x\cos x\,dx = 0.5$, where $0 < a < \tfrac{1}{2}\pi$.

(i) Show that a satisfies the equation $\sin a = \dfrac{1.5-\cos a}{a}$. [4]

(ii) Verify by calculation that a is greater than 1. [2]

(iii) Use the iterative formula

$$a_{n+1} = \sin^{-1}\left(\frac{1.5-\cos a_n}{a_n}\right)$$

to determine the value of a correct to 4 decimal places, giving the result of each iteration to 6 decimal places. [3]

© UCLES 2015

7 The number of micro-organisms in a population at time t is denoted by M. At any time the variation in M is assumed to satisfy the differential equation

$$\frac{\mathrm{d}M}{\mathrm{d}t} = k(\sqrt{M})\cos(0.02t),$$

where k is a constant and M is taken to be a continuous variable. It is given that when $t = 0$, $M = 100$.

(i) Solve the differential equation, obtaining a relation between M, k and t. [5]

(ii) Given also that $M = 196$ when $t = 50$, find the value of k. [2]

(iii) Obtain an expression for M in terms of t and find the least possible number of micro-organisms. [2]

8 The complex number $1 - \mathrm{i}$ is denoted by u.

(i) Showing your working and without using a calculator, express

$$\frac{\mathrm{i}}{u}$$

in the form $x + \mathrm{i}y$, where x and y are real. [2]

(ii) On an Argand diagram, sketch the loci representing complex numbers z satisfying the equations $|z - u| = |z|$ and $|z - \mathrm{i}| = 2$. [4]

(iii) Find the argument of each of the complex numbers represented by the points of intersection of the two loci in part **(ii)**. [3]

9 Two planes have equations $x + 3y - 2z = 4$ and $2x + y + 3z = 5$. The planes intersect in the straight line l.

(i) Calculate the acute angle between the two planes. [4]

(ii) Find a vector equation for the line l. [6]

10 Let $\mathrm{f}(x) = \dfrac{11x + 7}{(2x - 1)(x + 2)^2}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

(ii) Show that $\displaystyle\int_1^2 \mathrm{f}(x)\,\mathrm{d}x = \tfrac{1}{4} + \ln\left(\tfrac{9}{4}\right)$. [5]

© UCLES 2015

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2014**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

0962340356

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

© UCLES 2014

1 Use logarithms to solve the equation $e^x = 3^{x-2}$, giving your answer correct to 3 decimal places. [3]

2 **(i)** Use the trapezium rule with 3 intervals to estimate the value of

$$\int_{\frac{1}{6}\pi}^{\frac{2}{3}\pi} \operatorname{cosec} x \, \mathrm{d}x,$$

giving your answer correct to 2 decimal places. [3]

(ii) Using a sketch of the graph of $y = \operatorname{cosec} x$, explain whether the trapezium rule gives an overestimate or an underestimate of the true value of the integral in part **(i)**. [2]

3 The polynomial $ax^3 + bx^2 + x + 3$, where a and b are constants, is denoted by $\mathrm{p}(x)$. It is given that $(3x + 1)$ is a factor of $\mathrm{p}(x)$, and that when $\mathrm{p}(x)$ is divided by $(x - 2)$ the remainder is 21. Find the values of a and b. [5]

4 The parametric equations of a curve are

$$x = \frac{1}{\cos^3 t}, \quad y = \tan^3 t,$$

where $0 \leqslant t < \frac{1}{2}\pi$.

(i) Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \sin t$. [4]

(ii) Hence show that the equation of the tangent to the curve at the point with parameter t is $y = x \sin t - \tan t$. [3]

5 **Throughout this question the use of a calculator is not permitted.**

The complex numbers w and z satisfy the relation

$$w = \frac{z + \mathrm{i}}{\mathrm{i}z + 2}.$$

(i) Given that $z = 1 + \mathrm{i}$, find w, giving your answer in the form $x + \mathrm{i}y$, where x and y are real. [4]

(ii) Given instead that $w = z$ and the real part of z is negative, find z, giving your answer in the form $x + \mathrm{i}y$, where x and y are real. [4]

© UCLES 2014

6 It is given that $\int_1^a \ln(2x)\,\mathrm{d}x = 1$, where $a > 1$.

(i) Show that $a = \frac{1}{2}\exp\left(1 + \frac{\ln 2}{a}\right)$, where $\exp(x)$ denotes e^x. [6]

(ii) Use the iterative formula

$$a_{n+1} = \tfrac{1}{2}\exp\left(1 + \frac{\ln 2}{a_n}\right)$$

to determine the value of a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

7 In a certain country the government charges tax on each litre of petrol sold to motorists. The revenue per year is R million dollars when the rate of tax is x dollars per litre. The variation of R with x is modelled by the differential equation

$$\frac{\mathrm{d}R}{\mathrm{d}x} = R\left(\frac{1}{x} - 0.57\right),$$

where R and x are taken to be continuous variables. When $x = 0.5$, $R = 16.8$.

(i) Solve the differential equation and obtain an expression for R in terms of x. [6]

(ii) This model predicts that R cannot exceed a certain amount. Find this maximum value of R. [3]

8 **(i)** By first expanding $\sin(2\theta + \theta)$, show that

$$\sin 3\theta = 3\sin\theta - 4\sin^3\theta.$$ [4]

(ii) Show that, after making the substitution $x = \dfrac{2\sin\theta}{\sqrt{3}}$, the equation $x^3 - x + \frac{1}{6}\sqrt{3} = 0$ can be written in the form $\sin 3\theta = \frac{3}{4}$. [1]

(iii) Hence solve the equation

$$x^3 - x + \tfrac{1}{6}\sqrt{3} = 0,$$

giving your answers correct to 3 significant figures. [4]

9 Let $\mathrm{f}(x) = \dfrac{x^2 - 8x + 9}{(1-x)(2-x)^2}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

(ii) Hence obtain the expansion of $\mathrm{f}(x)$ in ascending powers of x, up to and including the term in x^2. [5]

10 The line l has equation $\mathbf{r} = 4\mathbf{i} - 9\mathbf{j} + 9\mathbf{k} + \lambda(-2\mathbf{i} + \mathbf{j} - 2\mathbf{k})$. The point A has position vector $3\mathbf{i} + 8\mathbf{j} + 5\mathbf{k}$.

(i) Show that the length of the perpendicular from A to l is 15. [5]

(ii) The line l lies in the plane with equation $ax + by - 3z + 1 = 0$, where a and b are constants. Find the values of a and b. [5]

© UCLES 2014

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2014**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

9231113471

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

www.satprep.org

This document consists of **3** printed pages and **1** blank page.

© UCLES 2014

1 Use logarithms to solve the equation $e^x = 3^{x-2}$, giving your answer correct to 3 decimal places. [3]

2 **(i)** Use the trapezium rule with 3 intervals to estimate the value of

$$\int_{\frac{1}{6}\pi}^{\frac{2}{3}\pi} \operatorname{cosec} x \, \mathrm{d}x,$$

giving your answer correct to 2 decimal places. [3]

(ii) Using a sketch of the graph of $y = \operatorname{cosec} x$, explain whether the trapezium rule gives an overestimate or an underestimate of the true value of the integral in part **(i)**. [2]

3 The polynomial $ax^3 + bx^2 + x + 3$, where a and b are constants, is denoted by $\mathrm{p}(x)$. It is given that $(3x + 1)$ is a factor of $\mathrm{p}(x)$, and that when $\mathrm{p}(x)$ is divided by $(x - 2)$ the remainder is 21. Find the values of a and b. [5]

4 The parametric equations of a curve are

$$x = \frac{1}{\cos^3 t}, \quad y = \tan^3 t,$$

where $0 \leqslant t < \frac{1}{2}\pi$.

(i) Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \sin t$. [4]

(ii) Hence show that the equation of the tangent to the curve at the point with parameter t is $y = x \sin t - \tan t$. [3]

5 **Throughout this question the use of a calculator is not permitted.**

The complex numbers w and z satisfy the relation

$$w = \frac{z + \mathrm{i}}{\mathrm{i}z + 2}.$$

(i) Given that $z = 1 + \mathrm{i}$, find w, giving your answer in the form $x + \mathrm{i}y$, where x and y are real. [4]

(ii) Given instead that $w = z$ and the real part of z is negative, find z, giving your answer in the form $x + \mathrm{i}y$, where x and y are real. [4]

© UCLES 2014

6 It is given that $\int_1^a \ln(2x)\,\mathrm{d}x = 1$, where $a > 1$.

(i) Show that $a = \frac{1}{2}\exp\left(1 + \frac{\ln 2}{a}\right)$, where $\exp(x)$ denotes e^x. [6]

(ii) Use the iterative formula

$$a_{n+1} = \tfrac{1}{2}\exp\left(1 + \frac{\ln 2}{a_n}\right)$$

to determine the value of a correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

7 In a certain country the government charges tax on each litre of petrol sold to motorists. The revenue per year is R million dollars when the rate of tax is x dollars per litre. The variation of R with x is modelled by the differential equation

$$\frac{\mathrm{d}R}{\mathrm{d}x} = R\left(\frac{1}{x} - 0.57\right),$$

where R and x are taken to be continuous variables. When $x = 0.5$, $R = 16.8$.

(i) Solve the differential equation and obtain an expression for R in terms of x. [6]

(ii) This model predicts that R cannot exceed a certain amount. Find this maximum value of R. [3]

8 **(i)** By first expanding $\sin(2\theta + \theta)$, show that

$$\sin 3\theta = 3\sin\theta - 4\sin^3\theta.$$ [4]

(ii) Show that, after making the substitution $x = \dfrac{2\sin\theta}{\sqrt{3}}$, the equation $x^3 - x + \frac{1}{6}\sqrt{3} = 0$ can be written in the form $\sin 3\theta = \frac{3}{4}$. [1]

(iii) Hence solve the equation

$$x^3 - x + \tfrac{1}{6}\sqrt{3} = 0,$$

giving your answers correct to 3 significant figures. [4]

9 Let $\mathrm{f}(x) = \dfrac{x^2 - 8x + 9}{(1-x)(2-x)^2}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

(ii) Hence obtain the expansion of $\mathrm{f}(x)$ in ascending powers of x, up to and including the term in x^2. [5]

10 The line l has equation $\mathbf{r} = 4\mathbf{i} - 9\mathbf{j} + 9\mathbf{k} + \lambda(-2\mathbf{i} + \mathbf{j} - 2\mathbf{k})$. The point A has position vector $3\mathbf{i} + 8\mathbf{j} + 5\mathbf{k}$.

(i) Show that the length of the perpendicular from A to l is 15. [5]

(ii) The line l lies in the plane with equation $ax + by - 3z + 1 = 0$, where a and b are constants. Find the values of a and b. [5]

© UCLES 2014

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2014**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

8698806885

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

© UCLES 2014

1 Solve the inequality $|3x - 1| < |2x + 5|$. [4]

2 A curve is defined for $0 < \theta < \frac{1}{2}\pi$ by the parametric equations

$$x = \tan\theta, \quad y = 2\cos^2\theta \sin\theta.$$

Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = 6\cos^5\theta - 4\cos^3\theta$. [5]

3 The polynomial $4x^3 + ax^2 + bx - 2$, where a and b are constants, is denoted by $\mathrm{p}(x)$. It is given that $(x + 1)$ and $(x + 2)$ are factors of $\mathrm{p}(x)$.

(i) Find the values of a and b. [4]

(ii) When a and b have these values, find the remainder when $\mathrm{p}(x)$ is divided by $(x^2 + 1)$. [3]

4 **(i)** Show that $\cos(\theta - 60°) + \cos(\theta + 60°) \equiv \cos\theta$. [3]

(ii) Given that $\dfrac{\cos(2x - 60°) + \cos(2x + 60°)}{\cos(x - 60°) + \cos(x + 60°)} = 3$, find the exact value of $\cos x$. [4]

5 The complex numbers w and z are defined by $w = 5 + 3\mathrm{i}$ and $z = 4 + \mathrm{i}$.

(i) Express $\dfrac{\mathrm{i}w}{z}$ in the form $x + \mathrm{i}y$, showing all your working and giving the exact values of x and y. [3]

(ii) Find wz and hence, by considering arguments, show that

$$\tan^{-1}\left(\tfrac{3}{5}\right) + \tan^{-1}\left(\tfrac{1}{4}\right) = \tfrac{1}{4}\pi.$$ [4]

6 It is given that $I = \displaystyle\int_0^{0.3} \left(1 + 3x^2\right)^{-2} \mathrm{d}x$.

(i) Use the trapezium rule with 3 intervals to find an approximation to I, giving the answer correct to 3 decimal places. [3]

(ii) For small values of x, $\left(1 + 3x^2\right)^{-2} \approx 1 + ax^2 + bx^4$. Find the values of the constants a and b. Hence, by evaluating $\displaystyle\int_0^{0.3} (1 + ax^2 + bx^4)\,\mathrm{d}x$, find a second approximation to I, giving the answer correct to 3 decimal places. [5]

© UCLES 2014

7 The equations of two straight lines are

$$\mathbf{r} = \mathbf{i} + 4\mathbf{j} - 2\mathbf{k} + \lambda(\mathbf{i} + 3\mathbf{k}) \quad \text{and} \quad \mathbf{r} = a\mathbf{i} + 2\mathbf{j} - 2\mathbf{k} + \mu(\mathbf{i} + 2\mathbf{j} + 3a\mathbf{k}),$$

where a is a constant.

(i) Show that the lines intersect for all values of a. [4]

(ii) Given that the point of intersection is at a distance of 9 units from the origin, find the possible values of a. [4]

8 The variables x and y are related by the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \tfrac{1}{5}xy^{\frac{1}{2}} \sin\left(\tfrac{1}{3}x\right).$$

(i) Find the general solution, giving y in terms of x. [6]

(ii) Given that $y = 100$ when $x = 0$, find the value of y when $x = 25$. [3]

9 **(i)** Sketch the curve $y = \ln(x + 1)$ and hence, by sketching a second curve, show that the equation

$$x^3 + \ln(x + 1) = 40$$

has exactly one real root. State the equation of the second curve. [3]

(ii) Verify by calculation that the root lies between 3 and 4. [2]

(iii) Use the iterative formula

$$x_{n+1} = \sqrt[3]{\left(40 - \ln(x_n + 1)\right)},$$

with a suitable starting value, to find the root correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]

(iv) Deduce the root of the equation

$$(\mathrm{e}^y - 1)^3 + y = 40,$$

giving the answer correct to 2 decimal places. [2]

10 By first using the substitution $u = \mathrm{e}^x$, show that

$$\int_0^{\ln 4} \frac{\mathrm{e}^{2x}}{\mathrm{e}^{2x} + 3\mathrm{e}^x + 2}\,\mathrm{d}x = \ln\left(\tfrac{8}{5}\right). \quad [10]$$

© UCLES 2014

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2014**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

* 5 3 7 4 6 3 8 5 8 2 *

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

www.satprep.org

This document consists of **3** printed pages and **1** blank page.

© UCLES 2014

1 **(i)** Simplify $\sin 2\alpha \sec \alpha$. [2]

(ii) Given that $3\cos 2\beta + 7\cos \beta = 0$, find the exact value of $\cos \beta$. [3]

2 Use the substitution $u = 1 + 3\tan x$ to find the exact value of

$$\int_0^{\frac{1}{4}\pi} \frac{\sqrt{(1 + 3\tan x)}}{\cos^2 x}\, \mathrm{d}x.$$ [5]

3 The parametric equations of a curve are

$$x = \ln(2t + 3), \quad y = \frac{3t + 2}{2t + 3}.$$

Find the gradient of the curve at the point where it crosses the y-axis. [6]

4 The variables x and y are related by the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{6y\mathrm{e}^{3x}}{2 + \mathrm{e}^{3x}}.$$

Given that $y = 36$ when $x = 0$, find an expression for y in terms of x. [6]

5 The complex number z is defined by $z = \dfrac{9\sqrt{3} + 9\mathrm{i}}{\sqrt{3} - \mathrm{i}}$. Find, showing all your working,

(i) an expression for z in the form $r\mathrm{e}^{\mathrm{i}\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$, [5]

(ii) the two square roots of z, giving your answers in the form $r\mathrm{e}^{\mathrm{i}\theta}$, where $r > 0$ and $-\pi < \theta \leqslant \pi$. [3]

6 It is given that $2\ln(4x - 5) + \ln(x + 1) = 3\ln 3$.

(i) Show that $16x^3 - 24x^2 - 15x - 2 = 0$. [3]

(ii) By first using the factor theorem, factorise $16x^3 - 24x^2 - 15x - 2$ completely. [4]

(iii) Hence solve the equation $2\ln(4x - 5) + \ln(x + 1) = 3\ln 3$. [1]

7 The straight line l has equation $\mathbf{r} = 4\mathbf{i} - \mathbf{j} + 2\mathbf{k} + \lambda(2\mathbf{i} - 3\mathbf{j} + 6\mathbf{k})$. The plane p passes through the point $(4, -1, 2)$ and is perpendicular to l.

(i) Find the equation of p, giving your answer in the form $ax + by + cz = d$. [2]

(ii) Find the perpendicular distance from the origin to p. [3]

(iii) A second plane q is parallel to p and the perpendicular distance between p and q is 14 units. Find the possible equations of q. [3]

© UCLES 2014

8 **(i)** By sketching each of the graphs $y = \operatorname{cosec} x$ and $y = x(\pi - x)$ for $0 < x < \pi$, show that the equation

$$\operatorname{cosec} x = x(\pi - x)$$

has exactly two real roots in the interval $0 < x < \pi$. [3]

(ii) Show that the equation $\operatorname{cosec} x = x(\pi - x)$ can be written in the form $x = \dfrac{1 + x^2 \sin x}{\pi \sin x}$. [2]

(iii) The two real roots of the equation $\operatorname{cosec} x = x(\pi - x)$ in the interval $0 < x < \pi$ are denoted by α and β, where $\alpha < \beta$.

(a) Use the iterative formula

$$x_{n+1} = \frac{1 + x_n^2 \sin x_n}{\pi \sin x_n}$$

to find α correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

(b) Deduce the value of β correct to 2 decimal places. [1]

9 **(i)** Express $\dfrac{4 + 12x + x^2}{(3 - x)(1 + 2x)^2}$ in partial fractions. [5]

(ii) Hence obtain the expansion of $\dfrac{4 + 12x + x^2}{(3 - x)(1 + 2x)^2}$ in ascending powers of x, up to and including the term in x^2. [5]

10



The diagram shows the curve $y = 10e^{-\frac{1}{2}x} \sin 4x$ for $x \geqslant 0$. The stationary points are labelled T_1, T_2, T_3, … as shown.

(i) Find the x-coordinates of T_1 and T_2, giving each x-coordinate correct to 3 decimal places. [6]

(ii) It is given that the x-coordinate of T_n is greater than 25. Find the least possible value of n. [4]

© UCLES 2014

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2014**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

1929888867

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

JC14 06_9709_32/RP
© UCLES 2014

1 Find the set of values of x satisfying the inequality

$$|x + 2a| > 3|x - a|,$$

where a is a positive constant. [4]

2 Solve the equation

$$2\ln(5 - e^{-2x}) = 1,$$

giving your answer correct to 3 significant figures. [4]

3 Solve the equation

$$\cos(x + 30°) = 2\cos x,$$

giving all solutions in the interval $-180° < x < 180°$. [5]

4 The parametric equations of a curve are

$$x = t - \tan t, \quad y = \ln(\cos t),$$

for $-\frac{1}{2}\pi < t < \frac{1}{2}\pi$.

(i) Show that $\dfrac{dy}{dx} = \cot t$. [5]

(ii) Hence find the x-coordinate of the point on the curve at which the gradient is equal to 2. Give your answer correct to 3 significant figures. [2]

5 **(i)** The polynomial $f(x)$ is of the form $(x - 2)^2 g(x)$, where $g(x)$ is another polynomial. Show that $(x - 2)$ is a factor of $f'(x)$. [2]

(ii) The polynomial $x^5 + ax^4 + 3x^3 + bx^2 + a$, where a and b are constants, has a factor $(x - 2)^2$. Using the factor theorem and the result of part **(i)**, or otherwise, find the values of a and b. [5]

© UCLES 2014

6



In the diagram, A is a point on the circumference of a circle with centre O and radius r. A circular arc with centre A meets the circumference at B and C. The angle OAB is equal to x radians. The shaded region is bounded by AB, AC and the circular arc with centre A joining B and C. The perimeter of the shaded region is equal to half the circumference of the circle.

(i) Show that $x = \cos^{-1}\left(\dfrac{\pi}{4+4x}\right)$. [3]

(ii) Verify by calculation that x lies between 1 and 1.5. [2]

(iii) Use the iterative formula

$$x_{n+1} = \cos^{-1}\left(\frac{\pi}{4+4x_n}\right)$$

to determine the value of x correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

7 **(a)** It is given that $-1 + (\sqrt{5})\mathrm{i}$ is a root of the equation $z^3 + 2z + a = 0$, where a is real. Showing your working, find the value of a, and write down the other complex root of this equation. [4]

(b) The complex number w has modulus 1 and argument 2θ radians. Show that $\dfrac{w-1}{w+1} = \mathrm{i}\tan\theta$. [4]

8



The diagram shows the curve $y = x\cos\frac{1}{2}x$ for $0 \leqslant x \leqslant \pi$.

(i) Find $\dfrac{\mathrm{d}y}{\mathrm{d}x}$ and show that $4\dfrac{\mathrm{d}^2y}{\mathrm{d}x^2} + y + 4\sin\frac{1}{2}x = 0$. [5]

(ii) Find the exact value of the area of the region enclosed by this part of the curve and the x-axis. [5]

© UCLES 2014

9 The population of a country at time t years is N millions. At any time, N is assumed to increase at a rate proportional to the product of N and $(1 - 0.01N)$. When $t = 0$, $N = 20$ and $\frac{\mathrm{d}N}{\mathrm{d}t} = 0.32$.

(i) Treating N and t as continuous variables, show that they satisfy the differential equation

$$\frac{\mathrm{d}N}{\mathrm{d}t} = 0.02N(1 - 0.01N). \quad [1]$$

(ii) Solve the differential equation, obtaining an expression for t in terms of N. [8]

(iii) Find the time at which the population will be double its value at $t = 0$. [1]

10 Referred to the origin O, the points A, B and C have position vectors given by

$$\overrightarrow{OA} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k}, \quad \overrightarrow{OB} = 2\mathbf{i} + 4\mathbf{j} + \mathbf{k} \quad \text{and} \quad \overrightarrow{OC} = 3\mathbf{i} + 5\mathbf{j} - 3\mathbf{k}.$$

(i) Find the exact value of the cosine of angle BAC. [4]

(ii) Hence find the exact value of the area of triangle ABC. [3]

(iii) Find the equation of the plane which is parallel to the y-axis and contains the line through B and C. Give your answer in the form $ax + by + cz = d$. [5]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Cambridge International A Level

Cambridge International Examinations
Cambridge International Advanced Level

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2014**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

7019745292

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

JC14 06_9709_33/FP
© UCLES 2014

1 Solve the equation $\log_{10}(x + 9) = 2 + \log_{10} x$. [3]

2 Expand $(1 + 3x)^{-\frac{1}{3}}$ in ascending powers of x, up to and including the term in x^3, simplifying the coefficients. [4]

3 **(i)** Show that the equation

$$\tan(x - 60°) + \cot x = \sqrt{3}$$

can be written in the form

$$2\tan^2 x + (\sqrt{3})\tan x - 1 = 0.$$ [3]

(ii) Hence solve the equation

$$\tan(x - 60°) + \cot x = \sqrt{3},$$

for $0° < x < 180°$. [3]

4 The equation $x = \dfrac{10}{e^{2x} - 1}$ has one positive real root, denoted by α.

(i) Show that α lies between $x = 1$ and $x = 2$. [2]

(ii) Show that if a sequence of positive values given by the iterative formula

$$x_{n+1} = \tfrac{1}{2}\ln\left(1 + \frac{10}{x_n}\right)$$

converges, then it converges to α. [2]

(iii) Use this iterative formula to determine α correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

5 The variables x and θ satisfy the differential equation

$$2\cos^2\theta \frac{dx}{d\theta} = \sqrt{(2x + 1)},$$

and $x = 0$ when $\theta = \frac{1}{4}\pi$. Solve the differential equation and obtain an expression for x in terms of θ. [7]

© UCLES 2014

6



The diagram shows the curve $(x^2 + y^2)^2 = 2(x^2 - y^2)$ and one of its maximum points M. Find the coordinates of M. [7]

7 **(a)** The complex number $\dfrac{3 - 5\mathrm{i}}{1 + 4\mathrm{i}}$ is denoted by u. Showing your working, express u in the form $x + \mathrm{i}y$, where x and y are real. [3]

(b) **(i)** On a sketch of an Argand diagram, shade the region whose points represent complex numbers satisfying the inequalities $|z - 2 - \mathrm{i}| \leqslant 1$ and $|z - \mathrm{i}| \leqslant |z - 2|$. [4]

(ii) Calculate the maximum value of $\arg z$ for points lying in the shaded region. [2]

8 Let $\mathrm{f}(x) = \dfrac{6 + 6x}{(2 - x)(2 + x^2)}$.

(i) Express $\mathrm{f}(x)$ in the form $\dfrac{A}{2 - x} + \dfrac{Bx + C}{2 + x^2}$. [4]

(ii) Show that $\displaystyle\int_{-1}^{1} \mathrm{f}(x)\,\mathrm{d}x = 3\ln 3$. [5]

9



The diagram shows the curve $y = \mathrm{e}^{2\sin x}\cos x$ for $0 \leqslant x \leqslant \frac{1}{2}\pi$, and its maximum point M.

(i) Using the substitution $u = \sin x$, find the exact value of the area of the shaded region bounded by the curve and the axes. [5]

(ii) Find the x-coordinate of M, giving your answer correct to 3 decimal places. [6]

© UCLES 2014

10 The line l has equation $\mathbf{r} = \mathbf{i} + 2\mathbf{j} - \mathbf{k} + \lambda(3\mathbf{i} - 2\mathbf{j} + 2\mathbf{k})$ and the plane p has equation $2x + 3y - 5z = 18$.

(i) Find the position vector of the point of intersection of l and p. [3]

(ii) Find the acute angle between l and p. [4]

(iii) A second plane q is perpendicular to the plane p and contains the line l. Find the equation of q, giving your answer in the form $ax + by + cz = d$. [5]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Advanced Level

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2013**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

5238158802

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a soft pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

www.satprep.co.in

This document consists of **4** printed pages.

© UCLES 2013

1 The equation of a curve is $y = \dfrac{1+x}{1+2x}$ for $x > -\frac{1}{2}$. Show that the gradient of the curve is always negative. [3]

2 Solve the equation $2|3^x - 1| = 3^x$, giving your answers correct to 3 significant figures. [4]

3 Find the exact value of $\displaystyle\int_1^4 \frac{\ln x}{\sqrt{x}}\,\mathrm{d}x$. [5]

4 The parametric equations of a curve are

$$x = \mathrm{e}^{-t}\cos t, \quad y = \mathrm{e}^{-t}\sin t.$$

Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \tan\left(t - \frac{1}{4}\pi\right)$. [6]

5 **(i)** Prove that $\cot\theta + \tan\theta \equiv 2\,\mathrm{cosec}\,2\theta$. [3]

(ii) Hence show that $\displaystyle\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \mathrm{cosec}\,2\theta\,\mathrm{d}\theta = \tfrac{1}{2}\ln 3$. [4]

6



In the diagram, A is a point on the circumference of a circle with centre O and radius r. A circular arc with centre A meets the circumference at B and C. The angle OAB is θ radians. The shaded region is bounded by the circumference of the circle and the arc with centre A joining B and C. The area of the shaded region is equal to half the area of the circle.

(i) Show that $\cos 2\theta = \dfrac{2\sin 2\theta - \pi}{4\theta}$. [5]

(ii) Use the iterative formula

$$\theta_{n+1} = \tfrac{1}{2}\cos^{-1}\left(\frac{2\sin 2\theta_n - \pi}{4\theta_n}\right),$$

with initial value $\theta_1 = 1$, to determine θ correct to 2 decimal places, showing the result of each iteration to 4 decimal places. [3]

© UCLES 2013

7 Let $\mathrm{f}(x) = \dfrac{2x^2 - 7x - 1}{(x-2)(x^2+3)}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

(ii) Hence obtain the expansion of $\mathrm{f}(x)$ in ascending powers of x, up to and including the term in x^2. [5]

8 Throughout this question the use of a calculator is not permitted.

(a) The complex numbers u and v satisfy the equations

$$u + 2v = 2\mathrm{i} \quad \text{and} \quad \mathrm{i}u + v = 3.$$

Solve the equations for u and v, giving both answers in the form $x + \mathrm{i}y$, where x and y are real. [5]

(b) On an Argand diagram, sketch the locus representing complex numbers z satisfying $|z + \mathrm{i}| = 1$ and the locus representing complex numbers w satisfying $\arg(w - 2) = \frac{3}{4}\pi$. Find the least value of $|z - w|$ for points on these loci. [5]

9



The diagram shows three points A, B and C whose position vectors with respect to the origin O are given by $\overrightarrow{OA} = \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix}$, $\overrightarrow{OB} = \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix}$ and $\overrightarrow{OC} = \begin{pmatrix} 3 \\ 0 \\ 4 \end{pmatrix}$. The point D lies on BC, between B and C, and is such that $CD = 2DB$.

(i) Find the equation of the plane ABC, giving your answer in the form $ax + by + cz = d$. [6]

(ii) Find the position vector of D. [1]

(iii) Show that the length of the perpendicular from A to OD is $\frac{1}{3}\sqrt{(65)}$. [4]

[Question 10 is printed on the next page.]

© UCLES 2013

10



A tank containing water is in the form of a cone with vertex C. The axis is vertical and the semi-vertical angle is 60°, as shown in the diagram. At time $t = 0$, the tank is full and the depth of water is H. At this instant, a tap at C is opened and water begins to flow out. The volume of water in the tank decreases at a rate proportional to $\sqrt{h}$, where h is the depth of water at time t. The tank becomes empty when $t = 60$.

(i) Show that h and t satisfy a differential equation of the form

$$\frac{\mathrm{d}h}{\mathrm{d}t} = -Ah^{-\frac{3}{2}},$$

where A is a positive constant. [4]

(ii) Solve the differential equation given in part **(i)** and obtain an expression for t in terms of h and H. [6]

(iii) Find the time at which the depth reaches $\frac{1}{2}H$. [1]

[The volume V of a cone of vertical height h and base radius r is given by $V = \frac{1}{3}\pi r^2 h$.]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Advanced Level

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2013**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

4229833005

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a soft pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

www.satprep.co.in

This document consists of **4** printed pages.

© UCLES 2013

1 The equation of a curve is $y = \dfrac{1+x}{1+2x}$ for $x > -\frac{1}{2}$. Show that the gradient of the curve is always negative. [3]

2 Solve the equation $2|3^x - 1| = 3^x$, giving your answers correct to 3 significant figures. [4]

3 Find the exact value of $\displaystyle\int_1^4 \frac{\ln x}{\sqrt{x}}\,\mathrm{d}x$. [5]

4 The parametric equations of a curve are

$$x = \mathrm{e}^{-t}\cos t, \quad y = \mathrm{e}^{-t}\sin t.$$

Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \tan\left(t - \frac{1}{4}\pi\right)$. [6]

5 **(i)** Prove that $\cot\theta + \tan\theta \equiv 2\,\mathrm{cosec}\,2\theta$. [3]

(ii) Hence show that $\displaystyle\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \mathrm{cosec}\,2\theta\,\mathrm{d}\theta = \tfrac{1}{2}\ln 3$. [4]

6



In the diagram, A is a point on the circumference of a circle with centre O and radius r. A circular arc with centre A meets the circumference at B and C. The angle OAB is θ radians. The shaded region is bounded by the circumference of the circle and the arc with centre A joining B and C. The area of the shaded region is equal to half the area of the circle.

(i) Show that $\cos 2\theta = \dfrac{2\sin 2\theta - \pi}{4\theta}$. [5]

(ii) Use the iterative formula

$$\theta_{n+1} = \tfrac{1}{2}\cos^{-1}\left(\frac{2\sin 2\theta_n - \pi}{4\theta_n}\right),$$

with initial value $\theta_1 = 1$, to determine θ correct to 2 decimal places, showing the result of each iteration to 4 decimal places. [3]

© UCLES 2013

7 Let $\mathrm{f}(x) = \dfrac{2x^2 - 7x - 1}{(x-2)(x^2+3)}$.

(i) Express $\mathrm{f}(x)$ in partial fractions. [5]

(ii) Hence obtain the expansion of $\mathrm{f}(x)$ in ascending powers of x, up to and including the term in x^2. [5]

8 Throughout this question the use of a calculator is not permitted.

(a) The complex numbers u and v satisfy the equations

$$u + 2v = 2\mathrm{i} \quad \text{and} \quad \mathrm{i}u + v = 3.$$

Solve the equations for u and v, giving both answers in the form $x + \mathrm{i}y$, where x and y are real. [5]

(b) On an Argand diagram, sketch the locus representing complex numbers z satisfying $|z + \mathrm{i}| = 1$ and the locus representing complex numbers w satisfying $\arg(w - 2) = \frac{3}{4}\pi$. Find the least value of $|z - w|$ for points on these loci. [5]

9



The diagram shows three points A, B and C whose position vectors with respect to the origin O are given by $\overrightarrow{OA} = \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix}$, $\overrightarrow{OB} = \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix}$ and $\overrightarrow{OC} = \begin{pmatrix} 3 \\ 0 \\ 4 \end{pmatrix}$. The point D lies on BC, between B and C, and is such that $CD = 2DB$.

(i) Find the equation of the plane ABC, giving your answer in the form $ax + by + cz = d$. [6]

(ii) Find the position vector of D. [1]

(iii) Show that the length of the perpendicular from A to OD is $\frac{1}{3}\sqrt{(65)}$. [4]

[Question 10 is printed on the next page.]

10



A tank containing water is in the form of a cone with vertex C. The axis is vertical and the semi-vertical angle is 60°, as shown in the diagram. At time $t = 0$, the tank is full and the depth of water is H. At this instant, a tap at C is opened and water begins to flow out. The volume of water in the tank decreases at a rate proportional to $\sqrt{h}$, where h is the depth of water at time t. The tank becomes empty when $t = 60$.

(i) Show that h and t satisfy a differential equation of the form

$$\frac{\mathrm{d}h}{\mathrm{d}t} = -Ah^{-\frac{3}{2}},$$

where A is a positive constant. [4]

(ii) Solve the differential equation given in part **(i)** and obtain an expression for t in terms of h and H. [6]

(iii) Find the time at which the depth reaches $\frac{1}{2}H$. [1]

[The volume V of a cone of vertical height h and base radius r is given by $V = \frac{1}{3}\pi r^2 h$.]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2013

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Advanced Level

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **October/November 2013**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

6391576228

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a soft pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

www.satprep.co.in

This document consists of **3** printed pages and **1** blank page.

© UCLES 2013

1 Given that $2\ln(x+4) - \ln x = \ln(x+a)$, express x in terms of a. [4]

2 Use the substitution $u = 3x + 1$ to find $\int \frac{3x}{3x+1}\,\mathrm{d}x$. [4]

3 The polynomial $\mathrm{f}(x)$ is defined by

$$\mathrm{f}(x) = x^3 + ax^2 - ax + 14,$$

where a is a constant. It is given that $(x + 2)$ is a factor of $\mathrm{f}(x)$.

(i) Find the value of a. [2]

(ii) Show that, when a has this value, the equation $\mathrm{f}(x) = 0$ has only one real root. [3]

4 A curve has equation $3\mathrm{e}^{2x}y + \mathrm{e}^{x}y^3 = 14$. Find the gradient of the curve at the point $(0, 2)$. [5]

5 It is given that $\int_0^p 4x\mathrm{e}^{-\frac{1}{2}x}\,\mathrm{d}x = 9$, where p is a positive constant.

(i) Show that $p = 2\ln\left(\frac{8p+16}{7}\right)$. [5]

(ii) Use an iterative process based on the equation in part **(i)** to find the value of p correct to 3 significant figures. Use a starting value of 3.5 and give the result of each iteration to 5 significant figures. [3]

6 Two planes have equations $3x - y + 2z = 9$ and $x + y - 4z = -1$.

(i) Find the acute angle between the planes. [3]

(ii) Find a vector equation of the line of intersection of the planes. [6]

7 **(i)** Given that $\sec\theta + 2\,\mathrm{cosec}\,\theta = 3\,\mathrm{cosec}\,2\theta$, show that $2\sin\theta + 4\cos\theta = 3$. [3]

(ii) Express $2\sin\theta + 4\cos\theta$ in the form $R\sin(\theta + \alpha)$ where $R > 0$ and $0° < \alpha < 90°$, giving the value of α correct to 2 decimal places. [3]

(iii) Hence solve the equation $\sec\theta + 2\,\mathrm{cosec}\,\theta = 3\,\mathrm{cosec}\,2\theta$ for $0° < \theta < 360°$. [4]

8 **(i)** Express $\frac{7x^2+8}{(1+x)^2(2-3x)}$ in partial fractions. [5]

(ii) Hence expand $\frac{7x^2+8}{(1+x)^2(2-3x)}$ in ascending powers of x up to and including the term in x^2, simplifying the coefficients. [5]

© UCLES 2013

9 **(a)** Without using a calculator, use the formula for the solution of a quadratic equation to solve

$$(2 - \mathrm{i})z^2 + 2z + 2 + \mathrm{i} = 0.$$

Give your answers in the form $a + b\mathrm{i}$. [5]

(b) The complex number w is defined by $w = 2\mathrm{e}^{\frac{1}{4}\pi\mathrm{i}}$. In an Argand diagram, the points A, B and C represent the complex numbers w, w^3 and w^* respectively (where w^* denotes the complex conjugate of w). Draw the Argand diagram showing the points A, B and C, and calculate the area of triangle ABC. [5]

10



A particular solution of the differential equation

$$3y^2 \frac{\mathrm{d}y}{\mathrm{d}x} = 4(y^3 + 1)\cos^2 x$$

is such that $y = 2$ when $x = 0$. The diagram shows a sketch of the graph of this solution for $0 \leqslant x \leqslant 2\pi$; the graph has stationary points at A and B. Find the y-coordinates of A and B, giving each coordinate correct to 1 decimal place. [10]

© UCLES 2013

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Advanced Level

MATHEMATICS **9709/31**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2013**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

* 4 0 6 4 0 1 4 4 9 0 *

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a soft pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

www.satprep.co.in

This document consists of **3** printed pages and **1** blank page.

© UCLES 2013

1 Find the quotient and remainder when $2x^2$ is divided by $x + 2$. [3]

2 Expand $\dfrac{1 + 3x}{\sqrt{(1 + 2x)}}$ in ascending powers of x up to and including the term in x^2, simplifying the coefficients. [4]

3 Express $\dfrac{7x^2 - 3x + 2}{x(x^2 + 1)}$ in partial fractions. [5]

4 **(i)** Solve the equation $|4x - 1| = |x - 3|$. [3]

(ii) Hence solve the equation $\left|4^{y+1} - 1\right| = |4^y - 3|$ correct to 3 significant figures. [3]

5 For each of the following curves, find the gradient at the point where the curve crosses the y-axis:

(i) $y = \dfrac{1 + x^2}{1 + \mathrm{e}^{2x}}$; [3]

(ii) $2x^3 + 5xy + y^3 = 8$. [4]

6 The points P and Q have position vectors, relative to the origin O, given by

$$\overrightarrow{OP} = 7\mathbf{i} + 7\mathbf{j} - 5\mathbf{k} \quad \text{and} \quad \overrightarrow{OQ} = -5\mathbf{i} + \mathbf{j} + \mathbf{k}.$$

The mid-point of PQ is the point A. The plane Π is perpendicular to the line PQ and passes through A.

(i) Find the equation of Π, giving your answer in the form $ax + by + cz = d$. [4]

(ii) The straight line through P parallel to the x-axis meets Π at the point B. Find the distance AB, correct to 3 significant figures. [5]

7 **(a)** Without using a calculator, solve the equation

$$3w + 2\mathrm{i}w^* = 17 + 8\mathrm{i},$$

where w^* denotes the complex conjugate of w. Give your answer in the form $a + b\mathrm{i}$. [4]

(b) In an Argand diagram, the loci

$$\arg(z - 2\mathrm{i}) = \tfrac{1}{6}\pi \quad \text{and} \quad |z - 3| = |z - 3\mathrm{i}|$$

intersect at the point P. Express the complex number represented by P in the form $r\mathrm{e}^{\mathrm{i}\theta}$, giving the exact value of θ and the value of r correct to 3 significant figures. [5]

8 **(a)** Show that $\displaystyle\int_2^4 4x \ln x \, \mathrm{d}x = 56 \ln 2 - 12$. [5]

(b) Use the substitution $u = \sin 4x$ to find the exact value of $\displaystyle\int_0^{\frac{1}{24}\pi} \cos^3 4x \, \mathrm{d}x$. [5]

© UCLES 2013

9 **(i)** Express $4\cos\theta + 3\sin\theta$ in the form $R\cos(\theta - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. Give the value of α correct to 4 decimal places. [3]

(ii) Hence

(a) solve the equation $4\cos\theta + 3\sin\theta = 2$ for $0 < \theta < 2\pi$, [4]

(b) find $\displaystyle\int \frac{50}{(4\cos\theta + 3\sin\theta)^2}\,\mathrm{d}\theta$. [3]

10 Liquid is flowing into a small tank which has a leak. Initially the tank is empty and, t minutes later, the volume of liquid in the tank is $V\,\mathrm{cm}^3$. The liquid is flowing into the tank at a constant rate of $80\,\mathrm{cm}^3$ per minute. Because of the leak, liquid is being lost from the tank at a rate which, at any instant, is equal to $kV\,\mathrm{cm}^3$ per minute where k is a positive constant.

(i) Write down a differential equation describing this situation and solve it to show that

$$V = \frac{1}{k}(80 - 80\mathrm{e}^{-kt}).$$ [7]

(ii) It is observed that $V = 500$ when $t = 15$, so that k satisfies the equation

$$k = \frac{4 - 4\mathrm{e}^{-15k}}{25}.$$

Use an iterative formula, based on this equation, to find the value of k correct to 2 significant figures. Use an initial value of $k = 0.1$ and show the result of each iteration to 4 significant figures. [3]

(iii) Determine how much liquid there is in the tank 20 minutes after the liquid started flowing, and state what happens to the volume of liquid in the tank after a long time. [2]

SATPREP www.satprep.co.in

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Advanced Level

MATHEMATICS **9709/32**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2013**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

* 0 1 5 7 9 2 6 8 8 0 *

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a soft pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

www.satprep.co.in

This document consists of **3** printed pages and **1** blank page.

© UCLES 2013

1 Solve the equation $|x - 2| = \left|\frac{1}{3}x\right|$. [3]

2 The sequence of values given by the iterative formula

$$x_{n+1} = \frac{x_n(x_n^3 + 100)}{2(x_n^3 + 25)},$$

with initial value $x_1 = 3.5$, converges to α.

(i) Use this formula to calculate α correct to 4 decimal places, showing the result of each iteration to 6 decimal places. [3]

(ii) State an equation satisfied by α and hence find the exact value of α. [2]

3



The variables x and y satisfy the equation $y = Ae^{-kx^2}$, where A and k are constants. The graph of $\ln y$ against x^2 is a straight line passing through the points (0.64, 0.76) and (1.69, 0.32), as shown in the diagram. Find the values of A and k correct to 2 decimal places. [5]

4 The polynomial $ax^3 - 20x^2 + x + 3$, where a is a constant, is denoted by $\mathrm{p}(x)$. It is given that $(3x + 1)$ is a factor of $\mathrm{p}(x)$.

(i) Find the value of a. [3]

(ii) When a has this value, factorise $\mathrm{p}(x)$ completely. [3]

5



The diagram shows the curve with equation

$$x^3 + xy^2 + ay^2 - 3ax^2 = 0,$$

where a is a positive constant. The maximum point on the curve is M. Find the x-coordinate of M in terms of a. [6]

© UCLES 2013

6 **(i)** By differentiating $\frac{1}{\cos x}$, show that the derivative of $\sec x$ is $\sec x \tan x$. Hence show that if $y = \ln(\sec x + \tan x)$ then $\frac{\mathrm{d}y}{\mathrm{d}x} = \sec x$. [4]

(ii) Using the substitution $x = (\sqrt{3}) \tan \theta$, find the exact value of

$$\int_1^3 \frac{1}{\sqrt{(3 + x^2)}} \, \mathrm{d}x,$$

expressing your answer as a single logarithm. [4]

7 **(i)** By first expanding $\cos(x + 45°)$, express $\cos(x + 45°) - (\sqrt{2}) \sin x$ in the form $R \cos(x + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$. Give the value of R correct to 4 significant figures and the value of α correct to 2 decimal places. [5]

(ii) Hence solve the equation

$$\cos(x + 45°) - (\sqrt{2}) \sin x = 2,$$

for $0° < x < 360°$. [4]

8 **(i)** Express $\frac{1}{x^2(2x + 1)}$ in the form $\frac{A}{x^2} + \frac{B}{x} + \frac{C}{2x + 1}$. [4]

(ii) The variables x and y satisfy the differential equation

$$y = x^2(2x + 1)\frac{\mathrm{d}y}{\mathrm{d}x},$$

and $y = 1$ when $x = 1$. Solve the differential equation and find the exact value of y when $x = 2$. Give your value of y in a form not involving logarithms. [7]

9 **(a)** The complex number w is such that $\mathrm{Re}\, w > 0$ and $w + 3w^* = \mathrm{i}w^2$, where w^* denotes the complex conjugate of w. Find w, giving your answer in the form $x + \mathrm{i}y$, where x and y are real. [5]

(b) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z which satisfy both the inequalities $|z - 2\mathrm{i}| \leqslant 2$ and $0 \leqslant \arg(z + 2) \leqslant \frac{1}{4}\pi$. Calculate the greatest value of $|z|$ for points in this region, giving your answer correct to 2 decimal places. [6]

10 The points A and B have position vectors $2\mathbf{i} - 3\mathbf{j} + 2\mathbf{k}$ and $5\mathbf{i} - 2\mathbf{j} + \mathbf{k}$ respectively. The plane p has equation $x + y = 5$.

(i) Find the position vector of the point of intersection of the line through A and B and the plane p. [4]

(ii) A second plane q has an equation of the form $x + by + cz = d$, where b, c and d are constants. The plane q contains the line AB, and the acute angle between the planes p and q is 60°. Find the equation of q. [7]

© UCLES 2013

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Advanced Level

MATHEMATICS **9709/33**

Paper 3 Pure Mathematics 3 **(P3)** **May/June 2013**

1 hour 45 minutes

Additional Materials: Answer Booklet/Paper
Graph Paper
List of Formulae (MF9)

9836805872

READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.
Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a soft pencil for any diagrams or graphs.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

www.satprep.co.in

This document consists of **3** printed pages and **1** blank page.

© UCLES 2013

1 Solve the inequality $|4x + 3| > |x|$. [4]

2 It is given that $\ln(y + 1) - \ln y = 1 + 3\ln x$. Express y in terms of x, in a form not involving logarithms. [4]

3 Solve the equation $\tan 2x = 5\cot x$, for $0° < x < 180°$. [5]

4 **(i)** Express $(\sqrt{3})\cos x + \sin x$ in the form $R\cos(x - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$, giving the exact values of R and α. [3]

(ii) Hence show that

$$\int_{\frac{1}{6}\pi}^{\frac{1}{2}\pi} \frac{1}{\left((\sqrt{3})\cos x + \sin x\right)^2}\,\mathrm{d}x = \tfrac{1}{4}\sqrt{3}.$$ [4]

5 The polynomial $8x^3 + ax^2 + bx + 3$, where a and b are constants, is denoted by $\mathrm{p}(x)$. It is given that $(2x + 1)$ is a factor of $\mathrm{p}(x)$ and that when $\mathrm{p}(x)$ is divided by $(2x - 1)$ the remainder is 1.

(i) Find the values of a and b. [5]

(ii) When a and b have these values, find the remainder when $\mathrm{p}(x)$ is divided by $2x^2 - 1$. [3]

6



The diagram shows the curves $y = \mathrm{e}^{2x-3}$ and $y = 2\ln x$. When $x = a$ the tangents to the curves are parallel.

(i) Show that a satisfies the equation $a = \frac{1}{2}(3 - \ln a)$. [3]

(ii) Verify by calculation that this equation has a root between 1 and 2. [2]

(iii) Use the iterative formula $a_{n+1} = \frac{1}{2}(3 - \ln a_n)$ to calculate a correct to 2 decimal places, showing the result of each iteration to 4 decimal places. [3]

© UCLES 2013

7 The complex number z is defined by $z = a + \mathrm{i}b$, where a and b are real. The complex conjugate of z is denoted by z^*.

(i) Show that $|z|^2 = zz^*$ and that $(z - k\mathrm{i})^* = z^* + k\mathrm{i}$, where k is real. [2]

In an Argand diagram a set of points representing complex numbers z is defined by the equation $|z - 10\mathrm{i}| = 2|z - 4\mathrm{i}|$.

(ii) Show, by squaring both sides, that

$$zz^* - 2\mathrm{i}z^* + 2\mathrm{i}z - 12 = 0.$$

Hence show that $|z - 2\mathrm{i}| = 4$. [5]

(iii) Describe the set of points geometrically. [1]

8 The variables x and t satisfy the differential equation

$$t\frac{\mathrm{d}x}{\mathrm{d}t} = \frac{k - x^3}{2x^2},$$

for $t > 0$, where k is a constant. When $t = 1$, $x = 1$ and when $t = 4$, $x = 2$.

(i) Solve the differential equation, finding the value of k and obtaining an expression for x in terms of t. [9]

(ii) State what happens to the value of x as t becomes large. [1]

9



The diagram shows the curve $y = \sin^2 2x \cos x$ for $0 \leqslant x \leqslant \frac{1}{2}\pi$, and its maximum point M.

(i) Find the x-coordinate of M. [6]

(ii) Using the substitution $u = \sin x$, find by integration the area of the shaded region bounded by the curve and the x-axis. [4]

10 The line l has equation $\mathbf{r} = \mathbf{i} + \mathbf{j} + \mathbf{k} + \lambda(a\mathbf{i} + 2\mathbf{j} + \mathbf{k})$, where a is a constant. The plane p has equation $x + 2y + 2z = 6$. Find the value or values of a in each of the following cases.

(i) The line l is parallel to the plane p. [2]

(ii) The line l intersects the line passing through the points with position vectors $3\mathbf{i} + 2\mathbf{j} + \mathbf{k}$ and $\mathbf{i} + \mathbf{j} - \mathbf{k}$. [4]

(iii) The acute angle between the line l and the plane p is $\tan^{-1} 2$. [5]

© UCLES 2013

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.