

Subject – Math AA(Higher Level)
Topic - Algebra
Year - May 2021 – Nov 2022
Paper -1
Questions

Question 1

[Maximum mark: 18]

- (a) Express $-3 + \sqrt{3}i$ in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leq \pi$. [5]

Let the roots of the equation $z^3 = -3 + \sqrt{3}i$ be u , v and w .

- (b) Find u , v and w expressing your answers in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leq \pi$. [5]

On an Argand diagram, u , v and w are represented by the points U , V and W respectively.

- (c) Find the area of triangle UVW . [4]

- (d) By considering the sum of the roots u , v and w , show that

$$\cos \frac{5\pi}{18} + \cos \frac{7\pi}{18} + \cos \frac{17\pi}{18} = 0. \quad [4]$$

Question 2

[Maximum mark: 8]

- (a) Show that $\log_9(\cos 2x + 2) = \log_3 \sqrt{\cos 2x + 2}$. [3]

- (b) Hence or otherwise solve $\log_3(2 \sin x) = \log_9(\cos 2x + 2)$ for $0 < x < \frac{\pi}{2}$. [5]

Question 3

[Maximum mark: 5]

- (a) Show that $(2n - 1)^2 + (2n + 1)^2 = 8n^2 + 2$, where $n \in \mathbb{Z}$. [2]

- (b) Hence, or otherwise, prove that the sum of the squares of any two consecutive odd integers is even. [3]

Question 4

[Maximum mark: 8]

A farmer has six sheep pens, arranged in a grid with three rows and two columns as shown in the following diagram.



Five sheep called Amber, Brownie, Curly, Daisy and Eden are to be placed in the pens. Each pen is large enough to hold all of the sheep. Amber and Brownie are known to fight.

Find the number of ways of placing the sheep in the pens in each of the following cases:

- (a) Each pen is large enough to contain five sheep. Amber and Brownie must not be placed in the same pen. [4]
- (b) Each pen may only contain one sheep. Amber and Brownie must not be placed in pens which share a boundary. [4]

Question 5

[Maximum mark: 8]

Consider the quartic equation $z^4 + 4z^3 + 8z^2 + 80z + 400 = 0$, $z \in \mathbb{C}$.

Two of the roots of this equation are $a + bi$ and $b + ai$, where $a, b \in \mathbb{Z}$.

Find the possible values of a .

Question 6


[Maximum mark: 5]

Consider an arithmetic sequence where $u_8 = S_8 = 8$. Find the value of the first term, u_1 , and the value of the common difference, d .

Question 7

[Maximum mark: 19]

The following diagram shows the graph of $y = \arctan(2x+1) + \frac{\pi}{4}$ for $x \in \mathbb{R}$, with asymptotes at $y = -\frac{\pi}{4}$ and $y = \frac{3\pi}{4}$.



- (a) Describe a sequence of transformations that transforms the graph of $y = \arctan x$ to the graph of $y = \arctan(2x+1) + \frac{\pi}{4}$ for $x \in \mathbb{R}$. [3]
- (b) Show that $\arctan p + \arctan q \equiv \arctan\left(\frac{p+q}{1-pq}\right)$ where $p, q > 0$ and $pq < 1$. [4]
- (c) Verify that $\arctan(2x+1) = \arctan\left(\frac{x}{x+1}\right) + \frac{\pi}{4}$ for $x \in \mathbb{R}, x > 0$. [3]
- (d) Using mathematical induction and the result from part (b), prove that $\sum_{r=1}^n \arctan\left(\frac{1}{2r^2}\right) = \arctan\left(\frac{n}{n+1}\right)$ for $n \in \mathbb{Z}^+$. [9]

Question 8

[Maximum mark: 5]

The cubic equation $x^3 - kx^2 + 3k = 0$ where $k > 0$ has roots α, β and $\alpha + \beta$.

Given that $\alpha\beta = -\frac{k^2}{4}$, find the value of k .

Question 9

[Maximum mark: 5]

In the expansion of $(x + k)^7$, where $k \in \mathbb{R}$, the coefficient of the term in x^5 is 63.

Find the possible values of k .

Question 10

[Maximum mark: 4]

Consider two consecutive positive integers, n and $n + 1$.

Show that the difference of their squares is equal to the sum of the two integers.

Question 11

[Maximum mark: 22]

Consider the equation $(z - 1)^3 = i$, $z \in \mathbb{C}$. The roots of this equation are ω_1 , ω_2 and ω_3 , where $\text{Im}(\omega_2) > 0$ and $\text{Im}(\omega_3) < 0$.

- (a) (i) Verify that $\omega_1 = 1 + e^{i\frac{\pi}{6}}$ is a root of this equation.
- (ii) Find ω_2 and ω_3 , expressing these in the form $a + e^{i\theta}$, where $a \in \mathbb{R}$ and $\theta > 0$. [6]

The roots ω_1 , ω_2 and ω_3 are represented by the points A, B and C respectively on an Argand diagram.

- (b) Plot the points A, B and C on an Argand diagram. [4]
- (c) Find AC. [3]

Consider the equation $(z - 1)^3 = iz^3$, $z \in \mathbb{C}$.

- (d) By using de Moivre's theorem, show that $\alpha = \frac{1}{1 - e^{i\frac{\pi}{6}}}$ is a root of this equation. [3]
- (e) Determine the value of $\text{Re}(\alpha)$. [6]

Question 12

[Maximum mark: 7]

Consider the expression $\frac{1}{\sqrt{1+ax}} - \sqrt{1-x}$ where $a \in \mathbb{Q}$, $a \neq 0$.

The binomial expansion of this expression, in ascending powers of x , as far as the term in x^2 is $4bx + bx^2$, where $b \in \mathbb{Q}$.

(a) Find the value of a and the value of b . [6]

(b) State the restriction which must be placed on x for this expansion to be valid. [1]

Question 13

[Maximum mark: 7]

The equation $3px^2 + 2px + 1 = p$ has two real, distinct roots.

(a) Find the possible values for p . [5]

(b) Consider the case when $p = 4$. The roots of the equation can be expressed in the form $x = \frac{a \pm \sqrt{13}}{6}$, where $a \in \mathbb{Z}$. Find the value of a . [2]

Question 14

[Maximum mark: 5]

Solve the equation $\log_3 \sqrt{x} = \frac{1}{2\log_2 3} + \log_3(4x^3)$, where $x > 0$.

Question 15

[Maximum mark: 18]

Consider the series $\ln x + p \ln x + \frac{1}{3} \ln x + \dots$, where $x \in \mathbb{R}, x > 1$ and $p \in \mathbb{R}, p \neq 0$.

(a) Consider the case where the series is geometric.

(i) Show that $p = \pm \frac{1}{\sqrt{3}}$.

(ii) Hence or otherwise, show that the series is convergent.

(iii) Given that $p > 0$ and $S_\infty = 3 + \sqrt{3}$, find the value of x . [6]

(b) Now consider the case where the series is arithmetic with common difference d .

(i) Show that $p = \frac{2}{3}$.

(ii) Write down d in the form $k \ln x$, where $k \in \mathbb{Q}$.

(iii) The sum of the first n terms of the series is $\ln\left(\frac{1}{x^3}\right)$.

Find the value of n . [12]

Question 16

[Maximum mark: 6]

Consider the complex numbers $z_1 = 1 + bi$ and $z_2 = (1 - b^2) - 2bi$, where $b \in \mathbb{R}, b \neq 0$.

(a) Find an expression for $z_1 z_2$ in terms of b . [3]

(b) Hence, given that $\arg(z_1 z_2) = \frac{\pi}{4}$, find the value of b . [3]

Question 17

[Maximum mark: 6]

Consider integers a and b such that $a^2 + b^2$ is exactly divisible by 4. Prove by contradiction that a and b cannot both be odd.

Question 18

[Maximum mark: 5]


Consider the expansion of $\left(8x^3 - \frac{1}{2x}\right)^n$ where $n \in \mathbb{Z}^+$. Determine all possible values of n for

which the expansion has a non-zero constant term.

Question 19

[Maximum mark: 18]

In the following Argand diagram, the points Z_1 , O and Z_2 are the vertices of triangle Z_1OZ_2 described anticlockwise.



The point Z_1 represents the complex number $z_1 = r_1 e^{i\alpha}$, where $r_1 > 0$. The point Z_2 represents the complex number $z_2 = r_2 e^{i\theta}$, where $r_2 > 0$.

Angles α , θ are measured anticlockwise from the positive direction of the real axis such that $0 \leq \alpha, \theta < 2\pi$ and $0 < \alpha - \theta < \pi$.

(a) Show that $z_1 z_2^* = r_1 r_2 e^{i(\alpha - \theta)}$ where z_2^* is the complex conjugate of z_2 . [2]

(b) Given that $\operatorname{Re}(z_1 z_2^*) = 0$, show that Z_1OZ_2 is a right-angled triangle. [2]

In parts (c), (d) and (e), consider the case where Z_1OZ_2 is an equilateral triangle.

(c) (i) Express z_1 in terms of z_2 . [2]

(ii) Hence show that $z_1^2 + z_2^2 = z_1 z_2$. [6]

Let z_1 and z_2 be the distinct roots of the equation $z^2 + az + b = 0$ where $z \in \mathbb{C}$ and $a, b \in \mathbb{R}$.

(d) Use the result from part (c)(ii) to show that $a^2 - 3b = 0$. [5]

Consider the equation $z^2 + az + 12 = 0$, where $z \in \mathbb{C}$ and $a \in \mathbb{R}$.

(e) Given that $0 < \alpha - \theta < \pi$, deduce that only one equilateral triangle Z_1OZ_2 can be formed from the point O and the roots of this equation. [3]

Question 20

[Maximum mark: 5]

Prove by contradiction that the equation $2x^3 + 6x + 1 = 0$ has no integer roots.

Question 21

[Maximum mark: 7]

Consider the binomial expansion $(x + 1)^7 = x^7 + ax^6 + bx^5 + 35x^4 + \dots + 1$ where $x \neq 0$ and $a, b \in \mathbb{Z}^+$.

(a) Show that $b = 21$. [2]

The third term in the expansion is the mean of the second term and the fourth term in the expansion.

(b) Find the possible values of x . [5]

Question 22

[Maximum mark: 6]

Consider any three consecutive integers, $n - 1$, n and $n + 1$.

(a) Prove that the sum of these three integers is always divisible by 3. [2]

(b) Prove that the sum of the squares of these three integers is never divisible by 3. [4]

Question 23

[Maximum mark: 5]

The n^{th} term of an arithmetic sequence is given by $u_n = 15 - 3n$.

(a) State the value of the first term, u_1 . [1]

(b) Given that the n^{th} term of this sequence is -33 , find the value of n . [2]

(c) Find the common difference, d . [2]

Question 24

[Maximum mark: 18]

Let z_n be the complex number defined as $z_n = (n^2 + n + 1) + i$ for $n \in \mathbb{N}$.

- (a) (i) Find $\arg(z_0)$.
- (ii) Write down an expression for $\arg(z_n)$ in terms of n . [3]

Let $w_n = z_0 z_1 z_2 z_3 \dots z_{n-1} z_n$ for $n \in \mathbb{N}$.

- (b) (i) Show that $\arctan(a) + \arctan(b) = \arctan\left(\frac{a+b}{1-ab}\right)$ for $a, b \in \mathbb{R}^+$, $ab < 1$.
- (ii) Hence or otherwise, show that $\arg(w_1) = \arctan(2)$. [5]
- (c) Prove by mathematical induction that $\arg(w_n) = \arctan(n+1)$ for $n \in \mathbb{N}$. [10]

Question 25

[Maximum mark: 16]

Consider a three-digit code abc , where each of a , b and c is assigned one of the values 1, 2, 3, 4 or 5.

- (a) Find the total number of possible codes
- (i) assuming that each value can be repeated (for example, 121 or 444);
- (ii) assuming that no value is repeated. [4]

Let $P(x) = x^3 + ax^2 + bx + c$, where each of a , b and c is assigned one of the values 1, 2, 3, 4 or 5. Assume that no value is repeated.

Consider the case where $P(x)$ has a factor of $(x^2 + 3x + 2)$.

- (b) (i) Find an expression for b in terms of a .
- (ii) Hence show that the only way to assign the values is $a = 4$, $b = 5$ and $c = 2$.
- (iii) Express $P(x)$ as a product of linear factors.
- (iv) Hence or otherwise, sketch the graph of $y = P(x)$, clearly showing the coordinates of any intercepts with the axes. [12]

Question 26

[Maximum mark: 7]

Consider the equation $z^4 + pz^3 + 54z^2 - 108z + 80 = 0$ where $z \in \mathbb{C}$ and $p \in \mathbb{R}$.

Three of the roots of the equation are $3 + i$, α and α^2 , where $\alpha \in \mathbb{R}$.

- (a) By considering the product of all the roots of the equation, find the value of α . [4]
- (b) Find the value of p . [3]

