

Subject – Math AA(Standard Level)
Topic - Calculus
Year - May 2021 – Nov 2022
Paper -1
Questions

Question 1

[Maximum mark: 16]

Let $f(x) = \frac{\ln 5x}{kx}$ where $x > 0, k \in \mathbb{R}^+$.

(a) Show that $f'(x) = \frac{1 - \ln 5x}{kx^2}$. [3]


The graph of f has exactly one maximum point P.

(b) Find the x -coordinate of P. [3]

The second derivative of f is given by $f''(x) = \frac{2 \ln 5x - 3}{kx^3}$. The graph of f has exactly one point of inflexion Q.

(c) Show that the x -coordinate of Q is $\frac{1}{5}e^{\frac{3}{2}}$. [3]

The region R is enclosed by the graph of f , the x -axis, and the vertical lines through the maximum point P and the point of inflexion Q.



(d) Given that the area of R is 3, find the value of k . [7]

Question 2

[Maximum mark: 16]

Let $f(x) = \frac{1}{3}x^3 + x^2 - 15x + 17$.

(a) Find $f'(x)$. [2]

The graph of f has horizontal tangents at the points where $x = a$ and $x = b$, $a < b$.

(b) Find the value of a and the value of b . [3]

(c) (i) Sketch the graph of $y = f'(x)$.

(ii) Hence explain why the graph of f has a local maximum point at $x = a$. [2]

(d) (i) Find $f''(b)$.

(ii) Hence, use your answer to part (d)(i) to show that the graph of f has a local minimum point at $x = b$. [4]

The normal to the graph of f at $x = a$ and the tangent to the graph of f at $x = b$ intersect at the point (p, q) .

(e) Find the value of p and the value of q . [5]

Question 3

[Maximum mark: 5]

Let $f'(x) = \frac{8x}{\sqrt{2x^2 + 1}}$. Given that $f(0) = 5$, find $f(x)$.

Question 4

[Maximum mark: 16]

Let $y = \frac{\ln x}{x^4}$ for $x > 0$.

(a) Show that $\frac{dy}{dx} = \frac{1 - 4 \ln x}{x^5}$. [3]

Consider the function defined by $f(x) = \frac{\ln x}{x^4}$ for $x > 0$ and its graph $y = f(x)$.

(b) The graph of f has a horizontal tangent at point P. Find the coordinates of P. [5]

(c) Given that $f''(x) = \frac{20 \ln x - 9}{x^6}$, show that P is a local maximum point. [3]

(d) Solve $f(x) > 0$ for $x > 0$. [2]

(e) Sketch the graph of f , showing clearly the value of the x -intercept and the approximate position of point P. [3]

Question 5

[Maximum mark: 7]

Consider the functions $f(x) = -(x - h)^2 + 2k$ and $g(x) = e^{x-2} + k$ where $h, k \in \mathbb{R}$.

(a) Find $f'(x)$. [1]

The graphs of f and g have a common tangent at $x = 3$.


(b) Show that $h = \frac{e+6}{2}$. [3]

(c) Hence, show that $k = e + \frac{e^2}{4}$. [3]

Question 6

[Maximum mark: 14]

Particle A travels in a straight line such that its displacement, s metres, from a fixed origin after t seconds is given by $s(t) = 8t - t^2$, for $0 \leq t \leq 10$, as shown in the following diagram.



Particle A starts at the origin and passes through the origin again when $t = p$.

- (a) Find the value of p . [2]

Particle A changes direction when $t = q$.

- (b) (i) Find the value of q .
(ii) Find the displacement of particle A from the origin when $t = q$. [4]

- (c) Find the distance of particle A from the origin when $t = 10$. [2]

The total distance travelled by particle A is given by d .

- (d) Find the value of d . [2]

A second particle, particle B, travels along the same straight line such that its velocity is given by $v(t) = 14 - 2t$, for $t \geq 0$.

When $t = k$, the distance travelled by particle B is equal to d .


- (e) Find the value of k . [4]

Question 7

[Maximum mark: 15]

Consider the function f defined by $f(x) = 6 + 6 \cos x$, for $0 \leq x \leq 4\pi$.

The following diagram shows the graph of $y = f(x)$.




The graph of f touches the x -axis at points A and B, as shown. The shaded region is enclosed by the graph of $y = f(x)$ and the x -axis, between the points A and B.

- (a) Find the x -coordinates of A and B. [3]
- (b) Show that the area of the shaded region is 12π . [5]

The right cone in the following diagram has a total surface area of 12π , equal to the shaded area in the previous diagram.

The cone has a base radius of 2, height h , and slant height l .

diagram not to scale




- (c) Find the value of l . [3]
- (d) Hence, find the volume of the cone. [4]

Question 8

[Maximum mark: 9]

Consider the function f defined by $f(x) = \ln(x^2 - 16)$ for $x > 4$.

The following diagram shows part of the graph of f which crosses the x -axis at point A, with coordinates $(a, 0)$. The line L is the tangent to the graph of f at the point B.




- (a) Find the exact value of a . [3]
- (b) Given that the gradient of L is $\frac{1}{3}$, find the x -coordinate of B. [6]

Question 9

[Maximum mark: 14]

Consider a function f with domain $a < x < b$. The following diagram shows the graph of f' , the derivative of f .



The graph of f' , the derivative of f , has x -intercepts at $x = p$, $x = 0$ and $x = t$. There are local maximum points at $x = q$ and $x = t$ and a local minimum point at $x = r$.

- (a) Find all the values of x where the graph of f is increasing. Justify your answer. [2]
- (b) Find the value of x where the graph of f has a local maximum. [1]
- (c) (i) Find the value of x where the graph of f has a local minimum. Justify your answer.
- (ii) Find the values of x where the graph of f has points of inflexion. Justify your answer. [5]
- (d) The total area of the region enclosed by the graph of f' , the derivative of f , and the x -axis is 20.
- Given that $f(p) + f(t) = 4$, find the value of $f(0)$. [6]

Question 10

[Maximum mark: 16]

A particle P moves along the x -axis. The velocity of P is $v \text{ m s}^{-1}$ at time t seconds, where $v(t) = 4 + 4t - 3t^2$ for $0 \leq t \leq 3$. When $t = 0$, P is at the origin O .

- (a) (i) Find the value of t when P reaches its maximum velocity. [7]
- (ii) Show that the distance of P from O at this time is $\frac{88}{27}$ metres. [7]
- (b) Sketch a graph of v against t , clearly showing any points of intersection with the axes. [4]
- (c) Find the total distance travelled by P . [5]

Question 11

[Maximum mark: 7]

The function f is defined for all $x \in \mathbb{R}$. The line with equation $y = 6x - 1$ is the tangent to the graph of f at $x = 4$.

- (a) Write down the value of $f'(4)$. [1]
- (b) Find $f(4)$. [1]

The function g is defined for all $x \in \mathbb{R}$ where $g(x) = x^2 - 3x$ and $h(x) = f(g(x))$.

- (c) Find $h(4)$. [2]
- (d) Hence find the equation of the tangent to the graph of h at $x = 4$. [3]

Question 12

[Maximum mark: 4]

Given that $\frac{dy}{dx} = \cos\left(x - \frac{\pi}{4}\right)$ and $y = 2$ when $x = \frac{3\pi}{4}$, find y in terms of x .

Question 13

[Maximum mark: 15]

(a) (i) Expand and simplify $(1 - a)^3$ in ascending powers of a .

(ii) By using a suitable substitution for a , show that
 $1 - 3 \cos 2x + 3 \cos^2 2x - \cos^3 2x = 8 \sin^6 x$.

[6]

Consider $f(x) = 4 \cos x(1 - 3 \cos 2x + 3 \cos^2 2x - \cos^3 2x)$.

(b) (i) Show that $\int_0^m f(x) dx = \frac{32}{7} \sin^7 m$, where m is a positive real constant.

(ii) It is given that $\int_m^{\frac{\pi}{2}} f(x) dx = \frac{127}{28}$, where $0 \leq m \leq \frac{\pi}{2}$. Find the value of m .

[9]

Question 14

[Maximum mark: 5]

Consider the curve with equation $y = (2x - 1)e^{kx}$, where $x \in \mathbb{R}$ and $k \in \mathbb{Q}$.

The tangent to the curve at the point where $x = 1$ is parallel to the line $y = 5e^k x$.

Find the value of k .

Question 15

[Maximum mark: 5]

(a) The expression $\frac{3\sqrt{x} - 5}{\sqrt{x}}$ can be written as $3 - 5x^p$. Write down the value of p .

[1]

(b) Hence, find the value of $\int_1^9 \left(\frac{3\sqrt{x} - 5}{\sqrt{x}} \right) dx$.


[4]

Question 16

[Maximum mark: 15]

Consider the functions $f(x) = \frac{1}{x-4} + 1$, for $x \neq 4$, and $g(x) = x - 3$ for $x \in \mathbb{R}$.

The following diagram shows the graphs of f and g .




The graphs of f and g intersect at points A and B. The coordinates of A are $(3, 0)$.

(a) Find the coordinates of B.

[5]


In the following diagram, the shaded region is enclosed by the graph of f , the graph of g , the x -axis, and the line $x = k$, where $k \in \mathbb{Z}$.



The area of the shaded region can be written as $\ln(p) + 8$, where $p \in \mathbb{Z}$.

(b) Find the value of k and the value of p .

[10]




Question 17

[Maximum mark: 15]

The following diagram shows part of the graph of a quadratic function f .

The graph of f has its vertex at $(3, 4)$, and it passes through point Q as shown.



- (a) Write down the equation of the axis of symmetry. [1]
- (b) The function can be written in the form $f(x) = a(x - h)^2 + k$.
- (i) Write down the values of h and k .
- (ii) Point Q has coordinates $(5, 12)$. Find the value of a . [4]

The line L is tangent to the graph of f at Q.

- (c) Find the equation of L . [4]

Now consider another function $y = g(x)$. The derivative of g is given by $g'(x) = f(x) - d$, where $d \in \mathbb{R}$.

- (d) Find the values of d for which g is an increasing function. [3]
- (e) Find the values of x for which the graph of g is concave-up. [3]

Question 18

[Maximum mark: 15]

The function f is defined by $f(x) = \cos^2 x - 3 \sin^2 x$, $0 \leq x \leq \pi$.

- (a) Find the roots of the equation $f(x) = 0$. [5]
- (b) (i) Find $f'(x)$.
- (ii) Hence find the coordinates of the points on the graph of $y = f(x)$ where $f'(x) = 0$. [7]
- (c) Sketch the graph of $y = f(x)$, clearly showing the coordinates of any points where $f'(x) = 0$ and any points where the graph meets the coordinate axes. [3]

Question 19

[Maximum mark: 5]

The derivative of the function f is given by $f'(x) = \frac{6x}{x^2 + 1}$.

The graph of $y = f(x)$ passes through the point $(1, 5)$. Find an expression for $f(x)$.

Question 20

[Maximum mark: 4]

The function g is defined by $g(x) = e^{x^2+1}$, where $x \in \mathbb{R}$.

Find $g'(-1)$.